

SNH (2017). Recommended bird survey methods to inform impact assessment of onshore wind farms. Scottish Natural Heritage, Inverness, Scotland. Available at: <https://www.nature.scot/sites/default/files/2018-06/Guidance%20Note%20-%20Recommended%20bird%20survey%20methods%20to%20inform%20impact%20assessment%20of%20onshore%20windfarms.pdf>

SNH (2018a) Avoidance rates for the onshore SNH wind farm collision risk model. Scottish Natural Heritage, Inverness, Scotland. Available at: <https://www.nature.scot/sites/default/files/2018-09/Wind%20farm%20impacts%20on%20birds%20-%20Use%20of%20Avoidance%20Rates%20in%20the%20SNH%20Wind%20Farm%20Collision%20Risk%20Model.pdf>

SNH (2018b). Assessing the cumulative impacts of onshore wind farms on birds. Scottish Natural Heritage, Inverness, Scotland. Available at: <https://www.nature.scot/sites/default/files/2018-08/Guidance%20-%20Assessing%20the%20cumulative%20impacts%20of%20onshore%20wind%20farms%20on%20birds.pdf>

SNH (2018c). Assessing significance of impacts from onshore wind farms outwith designated areas. Scottish Natural Heritage, Inverness, Scotland. Available at: <https://www.nature.scot/doc/guidance-assessing-significance-impacts-bird-populations-onshore-wind-farms-do-not-affect-protected>

Wildlife Act 1976, No. 39/1976, Dublin: Stationary Office. Available at: <http://www.irishstatutebook.ie/eli/1976/act/39/enacted/en/html?q=wildlife+act>

Alerstam, T., Rosén, M., Bäckman, J., Ericson, P.G.P. and Hellgren, O. (2007). Flight speeds among bird species: allometric and phylogenetic effects. *PLoS Biology*, 5(8): e197. <https://doi.org/10.1371/journal.pbio.0050197>

Duffy, K. and Steward, M. (2008) Turbine search methods and carcass removal trials at the Braes of Doune Windfarm. Natural Research Information Note 4. Natural Research Ltd, Banchory, UK. Available at: <https://ethys.pnnl.gov/sites/default/files/publications/Duffy-2008.pdf>

Evans P.R. and Pienkowski M.W. (1984) "Population Dynamics of Shorebirds". In: Burger J., Olla B.L. (eds) *Shorebirds*. Boston, MA: Springer

Fossitt, J.A. (2000) A guide to habitats in Ireland. The Heritage Council, Kilkenny, Ireland.

Gillings, S. and Fuller, R.J. (1999). Winter ecology of Golden Plover and Lapwings: a review and consideration of extensive survey methods. BTO Research Report No. 224, The National Centre for Ornithology, Norfolk, UK. Available at: https://www.bto.org/sites/default/files/shared_documents/publications/research-reports/1999/r224.pdf

Madders, M. and Whitfield, D.P. (2006) Upland raptors and the assessment of windfarm impacts. *Ibis*, 148(s1): 43-56. <https://doi.org/10.1111/j.1474-919X.2006.00506.x>

Orta, J., Boesman, P.F.D. and Marks, J.S. (2020). "Eurasian Kestrel (*Falco tinnunculus*)". In: Billerman, S.M., Keeney, B.K., Rodewald, P.G. and Schulenberg, T.S. (eds). *Birds of the World*. Ithaca, NY, USA: Cornell Lab of Ornithology.

Peach, W.J., Thompson, P.S. and Coulson, J.C. (1994). Annual and long-term variation in the survival rates of British Lapwings *Vanellus vanellus*. *Journal of Animal Ecology*, 63(1): 60-70. <https://doi.org/10.2307/5583>

Pearce-Higgins, J.W., Stephen, L., Douse, A. and Langston, R.H.W. (2012). Greater impacts of wind farms on bird populations during construction than subsequent operation: results of a multi-site and multi-species analysis. *Journal of Applied Ecology*, 49: 386–394. <https://doi.org/10.1111/j.1365-2656.2012.02110.x>

Pearce-Higgins, J.W., Stephen, L., Langston, R.H.W., Bainbridge, I.P. and Bullman, R. (2009) The distribution of breeding birds around upland wind farms. *Journal of Applied Ecology*, 46(6): 1323–1331. <https://doi.org/10.1111/j.1365-2656.2009.01715.x>

[REDACTED]

<https://www.npws.ie/sites/default/files/publications/pdf/rw/m93.pdf>

Spence, I.M. (1987). Mortality of snipe estimated from a mark and recapture Study. *Ring and Migration* 9(1)27-31. <https://doi.org/10.1080/03078698.1988.9673918>

Thaxter, C. B., Lascelles, B., Sugar, K., Cook, A. S. C. P., Roos, S. Bolton, M., Langston, R. H. W. and Burton, N. H. K. (2012). Seabird foraging ranges as a preliminary tool for identifying candidate Marine Protection Areas. *Biological Conservation*, 156: 53-61. <https://doi.org/10.1016/j.biocon.2011.12.009>

Village, A. (1990) *The Kestrel*. London: T. & A. D. Poyser.