APPENDIX No. 16

Conservation Objectives for Additional Habitats contained in Compensatory Area

1150 Coastal lagoons

To restore the favourable conservation condition of Coastal lagoons in Galway Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable, subject to slight natural variation. Favourable reference area 76.7ha. See map 4	Areas calculated from spatial data derived from Oliver, 2007. Site codes IL037, IL038, IL039, IL046, IL047, IL048, IL049, IL050, IL051, IL052. NB there may be more, as yet unmapped, lagoons within this SAC. See lagoon supporting document for further details
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 4 for mapped lagoons	Sites IL037, IL038, IL039, IL046, IL047, IL048, IL049, IL050, IL051, IL052 in Oliver, 2007. NB there may be more, as yet unmapped, lagoons within this SAC. See lagoon supporting document for further details
Salinity regime	Practical salinity units (psu)	Median annual salinity and temporal variation within natural ranges	The lagoons in the site vary from oligohaline to euhaline. See lagoon supporting document for further details
Hydrological regime	Metres	Annual water level fluctuations and minima within natural ranges	Most of the lagoons listed for this site are considered to be shallow; however, Aughinish lagoon and Lough Atalia do have deeper (at least 3m) parts. See lagoon supporting document for further details
Barrier: connectivity between lagoon and sea	Permeability	Appropriate hydrological connections between lagoons and sea, including where necessary, appropriate management	The lagoons within this site exhibit a variety of barrier types including cobble/shingle, karst and artificial embankment/causeway. Several are recorded as having sluices. See lagoon supporting document for further details
Water quality: Chlorophyll *a*	µg/L	Annual median chlorophyll *a* within natural ranges and less than 5µg/L	Target based on Roden and Oliver (2010). See lagoon supporting document for further details
Water quality: Molybdate Reactive Phosphorus (MRP)	mg/L	Annual median MRP within natural ranges 0.1mg/L	Target based on Roden and Oliver (2010). See lagoon supporting document for further details
Water quality: Dissolved Inorganic Nitrogen (DIN)	mg/L	Annual median DIN within natural ranges and less than 0.15mg/L	Target based on Roden and Oliver (2010). See lagoon supporting document for further details
Depth of macrophyte colonisation	Metres	Macrophyte colonisation to at least 2m depth	For shallow lagoons, it is expected that macrophytes should extend to their deepest points. See lagoon supporting document for further details
Typical plant species	Number and m²	Maintain number and extent of listed lagoonal specialists, subject to natural variation	Species listed in Oliver, 2007. See lagoon supporting document for further details
Typical animal species	Number	Maintain listed lagoon specialists, subject to natural variation	Species listed in Oliver, 2007. See lagoon supporting document for further details
Negative indicator species	Number and % cover	Negative indicator species absent or under control	Low salinity, shallow water and elevated nutrient levels increase the threat of accelerated encroachment by reedbeds. See lagoon supporting document for further details

1310 Salicornia and other annuals colonising mud and sand

To maintain the favourable conservation condition of *Salicornia* and other annuals colonizing mud and sand in Galway Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For sub-sites mapped: Barna House - 0.067ha, Seaweed Point - 0.003ha, Roscam West and South - 0.023ha, Kilcaimin - 0.015, Kileenaran - 0.007ha, Kinvara West - 0.017ha, Scanlan's Island - 0.117ha, Tawin Island - 1.098ha. See map 9	Based on data from Saltmarsh Monitoring Project (SMP) (McCorry and Ryle, 2009). Habitat recorded at eight of the ten sub-sites surveyed and mapped, giving a total estimated area of 1.347ha. N.B. Further unsurveyed areas may be present within this site. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, or change in habitat distribution, subject to natural processes. See map 9 for known distribution	Based on data from SMP (McCorry, 2007; McCorry and Ryle, 2009). *Salicornia* is an annual species, so its distribution can vary significantly from year to year. See coastal habitats supporting document for further details
Physical structure: sediment supply	Presence/ absence of physical barriers	Maintain/restore, natural circulation of sediments and organic matter, without any physical obstructions	Sediment supply is particularly important for pioneer saltmarsh community, as the distribution of this habitat depends on accretion rates. See coastal habitats supporting document for further details
Physical structure: creeks and pans	Occurrence	Maintain, or where necessary restore creek and pan structure, subject to natural processes, including erosion and succession	Based on data from SMP (McCorry, 2007; McCorry and Ryle, 2009). Creeks deliver sediment throughout saltmarsh system. Creeks and pan structures well developed at Kileenaran and Tawin Island. See coastal habitats supporting document for further details
Physical structure: flooding regime	Hectares flooded; frequency	Maintain natural tidal regime	This pioneer saltmarsh community requires regular tidal inundation. See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession.	Based on data from SMP (McCorry, 2007; McCorry and Ryle, 2009). See coastal habitats supporting document for further details
Vegetation structure: vegetation height	Centimetres	Maintain structural variation within sward	Based on data from SMP (McCorry, 2007; McCorry and Ryle, 2009). See coastal habitats supporting document for details
Vegetation structure: vegetation cover	Percentage cover at a representative sample of monitoring stops	Maintain more than 90% of area outside creeks vegetated	Based on data from SMP (McCorry, 2007; McCorry and Ryle, 2009). See coastal habitats supporting document for details
Vegetation composition: typical species and sub-communities	Percentage cover	Maintain the range of species-poor communities with typical species listed in SMP (McCorry and Ryle, 2009)	Based on data from SMP (McCorry, 2007; McCorry and Ryle, 2009). See coastal habitats supporting document for further details
Vegetation structure: negative indicator species - *Spartina anglica*	Hectares	There is currently no common cordgrass (*Spartina anglica*) in this SAC. Prevent establishment of cordgrass	Based on data from SMP (McCorry, 2007; McCorry and Ryle, 2009). See coastal habitats supporting document for further details

8240 Limestone pavements

To maintain the favourable conservation condition of Limestone pavements in Black Head-Poulsallagh Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable, subject to natural processes	Limestone pavements occurs in intimate association with other Annex I habitats in this SAC: Alpine and Boreal heaths (4060); *Juniperus communis* formations on heaths or calcareous grasslands (5130); Semi-natural dry grasslands and scrubland facies on calcareous substrates (Festuco-Brometalia) (6210); Petrifying springs with tufa formation (Cratoneurion) (7220). Therefore, these habitats cannot easily be mapped or considered separately. Conservation objectives for all these habitats should be used in conjunction with each other as appropriate. Wilson and Fernandez (2013) mapped the indicative area of limestone pavement, including associated habitats as 5,572ha (map 6). Four sites within the SAC were surveyed in detail as part of the national survey of limestone pavement and associated habitats (Wilson and Fernandez, 2013). This survey should be consulted for further details
Habitat distribution	Occurrence	No decline. Map 6 shows indicative distribution, including associated habitats	See notes for area above. Based on data from Wilson and Fernandez (2013). This habitat is split into exposed pavement and wooded pavement. In this SAC, the area of wooded pavement is relatively small
Vegetation composition: typical species	Number at a representative number of monitoring stops	At least seven positive indicator species present	Positive indicator species for exposed and wooded pavement are listed in Wilson and Fernandez (2013)
Vegetation composition: bryophyte layer	Percentage at a representative number of monitoring stops	Bryophyte cover at least 50% on wooded pavement	Attribute and target based on Wilson and Fernandez (2013)
Vegetation composition: negative indicator species	Percentage at a representative number of monitoring stops	Collective cover of negative indicator species on exposed pavement not more than 1%	Negative indicator species listed in Wilson and Fernandez (2013). Negative indicator species for wooded pavement overlap with non-native species (below)
Vegetation composition: non-native species	Percentage at a representative number of monitoring stops	Cover of non-native species not more than 1% on exposed pavement; on wooded pavement not more than 10% with no regeneration	Attribute and target based on Wilson and Fernandez (2013)
Vegetation composition: scrub	Percentage at a representative number of monitoring stops	Scrub cover no more than 25% of exposed pavement	Attribute and target based on Wilson and Fernandez (2013). This SAC has very little scrub cover compared with areas further inland
Vegetation composition: bracken cover	Percentage at a representative number of monitoring stops	Bracken (*Pteridium aquilinum*) cover no more than 10% on exposed pavement	Attribute and target based on Wilson and Fernandez (2013)
Vegetation structure: woodland canopy	Percentage at a representative number of monitoring stops	Canopy cover on wooded pavement at least 30%	Wooded limestone pavement in this SAC is mostly low-growing hazel (*Corylus avellana*) woodland, some of which can be classified as Atlantic hazel woodland, an internationally rare woodland type. Despite its low stature it is nonetheless an important habitat for woodland species. Attribute and target based on Wilson and Fernandez (2013)
Vegetation structure: dead wood	Occurrence in a representative number of monitoring stops	Sufficient quantity of dead wood on wooded pavement to provide habitat for saproxylic organisms	Dead wood is a valuable resource and an integral part of a healthy, functioning woodland ecosystem

Physical structure: disturbance	Occurrence in a representative number of monitoring stops	No evidence of grazing pressure on wooded pavement	Attribute and target based on Wilson and Fernandez (2013)
Indicators of local distinctiveness	Occurrence	Indicators of local distinctiveness are maintained	Includes red-data and other rare or localised species as well as archaeological and geological features, which often support distinctive species