APPENDIX No. 3

Conservation Objectives for Galway Bay Complex cSAC (000268)

- 1140 Mudflatsand Sandflats not covered by seawater at low tide
- 1170 Reefs
- 1220 Perennial Vegetation of Stony Banks
- 1330 Atlantic Salt Meadows
- 1410 Mediterranean Salt Meadows

National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,

7 Ely Place, Dublin 2, Ireland.

Web: www.npws.ie
E-mail: nature.conservation@ahg.gov.ie

Citation:

NPWS (2013) Conservation Objectives: Galway Bay Complex SAC 000268. Version 1. National Parks and Wildlife Service, Department of Arts, Heritage and the Gaeltacht.

Series Editor: Rebecca Jeffrey
ISSN 2009-4086

1140 Mudflats and sandflats not covered by seawater at low tide

To maintain the favourable conservation condition of Mudflats and sandflats not covered by seawater at low tide in Galway Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	The permanent habitat area is stable or increasing, subject to natural processes. See map 3	Habitat area was estimated using OSi data as 744ha
Community distribution	Hectares	Conserve the following community types in a natural condition: Intertidal sandy mud community complex; and Intertidal sand community complex. See map 7	Based on intertidal surveys undertaken in 2009 and 2010 (RPS, 2012). See marine supporting document for further information

1170 Reefs

To maintain the favourable conservation condition of Reefs in Galway Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Occurrence	The distribution of reefs is stable or increasing, subject to natural processes. See map 6 for mapped distribution	Based on information from 2009 and 2010 intertidal survey data and 2009 subtidal survey data (Aquafact, 2010a, b; RPS, 2012). See marine supporting document for further details
Habitat area	Hectares	The permanent habitat area is stable, subject to natural processes. See map 6	Habitat area estimated as 2773ha using 2009 and 2010 intertidal survey data and 2009 subtidal survey data (Aquafact, 2010a, b; RPS, 2012)
Community extent	Hectares	Maintain the extent of the *Mytilus*-dominated reef community, subject to natural processes. See map 7	Area established from 2009 intertidal survey (RPS, 2012)
Community structure: *Mytilus* density	Individuals per m^2	Conserve the high quality of the *Mytilus*-dominated reef community, subject to natural processes	Based on intertidal survey 2009 (RPS, 2012) and intertidal walkover 2012
Community structure	Biological composition	Conserve the following community types in a natural condition: Fucoid-dominated community complex; *Laminaria*-dominated community complex; and Shallow sponge-dominated community complex See map 7	Reef mapping based on information from 2009 subtidal reef survey (Aquafact, 2010b) and 2009 and 2010 intertidal surveys (RPS, 2012). See marine supporting document for further details

1220 Perennial vegetation of stony banks

To maintain the favourable conservation condition of Perennial vegetation of stony banks in Galway Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession	Current area unknown. It was recorded from Rinville Point, Tawin Point and coastline from Blackhead to Carrickada during the National Shingle Beach Survey (Moore and Wilson, 1999), but the extent was not mapped. Two areas of vegetated shingle were recorded during the Coastal Monitoring Project (Ryle et al., 2009): Bishopsquarter - 0.18ha and Barna (Whitestrand) - 0.45ha. NB further unsurveyed areas maybe present within the site. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, or change in habitat distribution, subject to natural processes. See map 8 for mapped locations	Full distribution unmapped at present, although the habitat has been recorded at Rinville Point, Tawin Point and coastline from Blackhead to Carrickada (Moore and Wilson, 1999). It has also been recorded from Barna and Bishopquarter by Ryle et al. (2009). See coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/ absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	The Galway Bay shoreline supports good examples of shingle beaches along the more exposed shores to the south and west of Galway city and to the north-east of Finnavara, County Clare. Shingle features are relatively stable in the longterm (Moore and Wilson, 1999). See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Moore and Wilson (1999). See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative sample of monitoring stops	Maintain the typical vegetated shingle flora including the range of sub-communities within the different zones. Typical species include sea sandwort (*Honckenya peploides*), sea beet (*Beta vulgaris* ssp *maritima*), rock samphire (*Crithmum maritimum*), sea mayweed (*Tripleurospermum maritimum*), yellow-horned poppy (*Glaucium flavum*) and sea campion (*Silene uniflora*)	Based on data from Moore and Wilson (1999). See coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-natives) to represent less than 5% cover	Based on data from Moore and Wilson (1999). Negative indicators include non-native species indicative of changes in nutrient status and species not considered characteristic of the habitat. See coastal habitats supporting document for further details

1330 Atlantic salt meadows (Glauco-Puccinellietalia maritimae)

To restore the favourable conservation condition of Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*) in Galway Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area increasing, subject to natural processes, including erosion and succession. For sub-sites mapped: Barna House - 2.33ha, Seaweed Point - 1.41ha, Roscam West and South - 3.30ha, Oranmore North - 4.24ha, Kilcaimin - 6.82ha, Tawin Island - 53.85ha, Tyrone House-Dunbulcaun Bay - 9.83ha, Kileenaran - 15.37ha, Kinvara West - 13.33ha, Scanlan's Island - 4.13ha. See map 9	Based on data from Saltmarsh monitoring Project (SMP) (McCorry, 2007; McCorry and Ryle, 2009). Ten sub-sites that supported Atlantic salt meadow were mapped (114.612ha) and additional areas of potential saltmarsh (149.18ha) were identified by an examination of aerial photographs, giving a total estimated area of 263.80ha. NB further unsurveyed areas maybe present within the site. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline or change in habitat distribution, subject to natural processes. See map 9 for known distribution	Based on data from SMP (McCorry, 2007; McCorry and Ryle, 2009). See coastal habitats supporting document for further details
Physical structure: sediment supply	Presence/ absence of physical barriers	Maintain/restore natural circulation of sediments and organic matter, without any physical obstructions	See coastal habitats supporting document for further details
Physical structure: creeks and pans	Occurrence	Maintain creek and pan structure, subject to natural processes, including erosion and succession	Based on data from SMP (McCorry, 2007; McCorry and Ryle, 2009). The efficiency of sediment circulation throughout a saltmarsh depends on the creek pattern. Creeks and pans are well developed at both Tawin Island and Kileenaran. See coastal habitats supporting document for further details
Physical structure: flooding regime	Hectares flooded; frequency	Maintain natural tidal regime	See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from SMP (McCorry, 2007; McCorry and Ryle, 2009). See coastal habitats supporting document for further details
Vegetation structure: vegetation height	Centimetres	Maintain structural variation within sward	Based on data from SMP (McCorry, 2007; McCorry and Ryle, 2009). See coastal habitats supporting document for further details
Vegetation structure: vegetation cover	Percentage cover at a representative sample of monitoring stops	Maintain more than 90% area outside creeks vegetated	Based on data from SMP (McCorry, 2007; McCorry and Ryle, 2009). See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative sample of monitoring stops	Maintain range of sub-communities with typical species listed in SMP (McCorry and Ryle, 2009)	Based on data from SMP (McCorry, 2007; McCorry and Ryle, 2009). See coastal habitats supporting document for further details
Vegetation structure: negative indicator species - *Spartina anglica*	Hectares	There is currently no common cordgrass (*Spartina anglica*) in this SAC. Prevent establishment of cordgrass	Based on data from SMP (McCorry, 2007; McCorry and Ryle, 2009). McCorry and Ryle, 2009). See coastal habitats supporting document for further details

1410 Mediterranean salt meadows (Juncetalia maritimi)

To restore the favourable conservation condition of Mediterranean salt meadows (*Juncetalia maritimi*) in Galway Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For sub-sites mapped: Barna House - 0.282ha, Seaweed Point - 0.931ha, Kilcaimin - 0.005ha, Tawin Island - 1.799ha. Tyrone House- Dunbulcan Bay - 8.184ha, Kileenaran - 0.271ha. See map 9	Based on data from the Saltmarsh Monitoring Project (SMP) (McCorry, 2007; McCorry and Ryle, 2009). Six sub-sites that support Mediterranean salt meadow were mapped (11.472ha) and additional areas of potential saltmarsh (8.415ha) were identified from an examination of aerial photographs, giving a total estimated area of 19.887ha. NB further unsurveyed areas maybe present within the site. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 9 for known distribution	See coastal habitats supporting document for further details
Physical structure: sediment supply	Presence/absence of physical barriers	Maintain/restore natural circulation of sediments and organic matter, without any physical obstructions	See coastal habitats supporting document for further details
Physical structure: creeks and pans	Occurrence	Maintain creek and pan structure, subject to natural processes, including erosion and succession	Based on data from the SMP (McCorry, 2007; McCorry and Ryle, 2009). [Site-specific info.]. See coastal habitats supporting document for further details
Physical structure: flooding regime	Hectares flooded; frequency	Maintain natural tidal regime	Mediterranean salt meadows is found high up in the saltmarsh but requires occasional tidal inundation. [Site-specific info.] See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from SMP (McCorry, 2007; McCorry and Ryle, 2009). See coastal habitats supporting document for further details
Vegetation structure: vegetation height	Centimetres	Maintain structural variation in the sward	Based on data from SMP (McCorry, 2007; McCorry and Ryle, 2009). See coastal habitats supporting document for further details
Vegetation structure: vegetation cover	Percentage cover at a representative sample of monitoring stops	Maintain more than 90% of area outside creeks vegetated	Based on data from SMP (McCorry, 2007; McCorry and Ryle (2009). See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative sample of monitoring stops	Maintain range of sub-communities with typical species listed in SMP (McCorry and Ryle, 2009)	Based on data from SMP (McCorry, 2007; McCorry and Ryle (2009). See coastal habitats supporting document for further details
Vegetation structure: negative indicator species - *Spartina anglica*	Hectares	There is currently no common cordgrass (*Spartina anglica*) in this SAC. Prevent establishment of cordgrass	Based on data from SMP (McCorry, 2007; McCorry and Ryle, 2009). See coastal habitats supporting document for further details