



Application for Substitute Consent for an Existing Sand and Gravel Pit at Roscat, Tullow, Co. Carlow

Remedial Natura Impact Statement

April 2019

Document Details

Publisher: Earth Science Partnership Ireland Ltd.
Tonranny,
Westport,
Co. Mayo

T: +353 (0) 98 28999
E: earthsciencepartnership@gmail.com
W: <http://www.earthsciencepartnership.ie/>



Earth Science Partnership Ireland Ltd.
Consulting Engineers, Geologists & Environmental Scientists

Client: Kilcarrig Quarries Ltd.

Document Title: Application for Substitute Consent at an Existing Sand and Gravel Pit at Roscat, Tullow, Co. Carlow – Remedial Natura Impact Statement

Document Author: Ms. Sarah Ingham MSc (Biodiv. Con.) BSc (Hons.) (Zoology) ACIEEM
Senior Ecologist, Earth Science Partnership Ireland Ltd.
<https://www.linkedin.com/in/sarah-ingham-8b782043/>

Project Location: Roscat, Tullow, Co. Carlow

Document Issue:

Revision	Status	Issue Date	Author	Approved by
Rev 1.0	Final	11/04/19	SI	POD

Notice

This report was produced by Earth Science Partnership Ireland on behalf of Kilcarrig Quarries Ltd, the client, for the specific purpose of an application for Substitute Consent at an existing sand and gravel pit at Roscat, Tullow, Co. Carlow, with all reasonable skill, care and due diligence within the terms of the contract with the client, incorporating our terms and conditions and taking account of the resources devoted to it by agreement with the client.

This report may not be used by any person other than Kilcarrig Quarries Ltd, the client, without the client's express permission. In any event, Earth Science Partnership Ireland accepts no liability for any costs, liabilities or losses arising as a result of the use of or reliance upon the contents of this report by any person other than the client.

This report is confidential to the client and Earth Science Partnership Ireland accepts no responsibility of whatsoever nature to third parties to whom this report, or any part thereof, is made known. Any such party relies upon the report at their own risk.

© Earth Science Partnership Ireland, 2019

Table of Contents

1	INTRODUCTION	1
1.1	Project Background and Report Objective.....	1
2	METHODOLOGY.....	3
2.1	Legislative Context.....	3
2.1.1	<i>Article 6 of the Habitats Directive</i>	3
2.1.2	<i>Appropriate Assessment Process</i>	3
2.2	Relevant Guidance.....	4
2.3	Consultation	5
3	PROJECT DESCRIPTION	6
3.1	Existing Substitute Consent Site.....	6
4	RECEIVING ENVIRONMENT	6
4.1	Natura 2000 Sites	6
4.1.1	<i>Identification of Relevant Natura 2000 Sites</i>	6
4.1.2	<i>River Barrow and River Nore SAC (Site Code: 002162)</i>	12
5	STAGE 1 – SCREENING FOR APPROPRIATE ASSESSMENT	14
5.1	Introduction.....	14
5.2	Potential Risks to Natura 2000 Sites	14
5.2.1	<i>Potential Risks as a Result of the Existing Project</i>	14
5.3	Evaluation of Likely Effects.....	15
5.4	Potential In-combination Effects.....	18
5.4.1	<i>Capped Landfill Site</i>	18
5.4.2	<i>Existing Ardristan Sand and Gravel Pit (QY 27)</i>	18
5.5	Screening for Appropriate Assessment Matrix.....	20
5.6	Screening Statement and Conclusions.....	21
6	STAGE 2 – APPROPRIATE ASSESSMENT REPORT	22
6.1	Assessment of the Effects of the Project or Plan on the Integrity of the Sites	22
6.2	The Conservation Objectives of the Natura 2000 Sites	22
6.3	Potential Impacts on Key Species and Key Habitats	23
6.4	Potential Impacts on the Integrity of the Sites	24
6.5	Remedial Measures	26
6.6	Efficacy of the proposed Mitigation Measures.....	29
6.7	Conclusion	30
7	REFERENCES	31
	APPENDIX I – NATURA 2000 SITE SYNOPSES AND CONSERVATION OBJECTIVES.....	32
	APPENDIX II – RESPONSE TO SCOPING CONSULTATION	33

1 INTRODUCTION

1.1 Project Background and Report Objective

Earth Science Partnership Ireland (ESPI) was commissioned by Kilcarrig Quarries to produce a remedial Natura Impact Statement (rNIS) to accompany an application for Substitute Consent regarding the historical development of an existing sand and gravel pit located at Roscat, Tullow, Co. Carlow (Figure 1.1).

This Substitute Consent application consists of an existing 4.7 hectare sand and gravel pit.

The site is situated on the eastern side of a minor valley which is defined by a small north-south trending ridge that peaks at 94 mOD, 1 km northeast of the site at Ellengrove Crossroads, and a wider area of raised ground which reaches 93 mOD, 1.8 km southwest of the site. This valley flattens out to less than 70 mOD a short distance southwest of the site. OS Discovery maps indicate the site elevation to be in the range 68 – 74 mOD.

The area immediately around the site is sparsely populated, with individual farmsteads and scattered houses along the road network. A series of irregular third class roads run around the lands, serving a number of dwellings and farms.

The existing sand and gravel pit area has been registered with Carlow County Council in accordance with the requirements of Section 261 of the Planning and Development Act, 2000 (Quarry Ref. No. QY12/28). An area of 6.02ha of the landholding was registered in this process which is comprised of the 4.7ha Substitute Consent application area and an area of 1.3ha which is subject to an existing planning permission (Planning Ref: CW7850).


This document screens and evaluates likely significant effects of the existing development upon European designated sites (Natura 2000) during its operation. A scientific and objective approach has been used to identify these Natura 2000 sites, such as hydrological and/or ecological connections, and to whether their conservation objectives are at risk from the development for specific reasons. This report is conducted in line with the requirements of Article 6(3) of the EU habitats Directive (92/43/EEC) and the National Parks and Wildlife Service (NPWS) Guidance for Planning Authorities (2009).

It is intended that the information contained within this document will form the basis for the Article 6(3) Appropriate Assessment process which will be completed by the Competent Authority.

THIS DRAWING AND ANY DESIGN HEREON IS THE COPYRIGHT OF THE CONSULTANT AND MAY NOT BE COPIED, REPRODUCED OR TRANSMITTED IN ANY FORM WITHOUT THE WRITTEN CONSENT OF THE CONSULTANT. ALL DIMENSIONS TO BE CHECKED ON SITE AND ANY DISCREPANCIES TO BE REPORTED TO THE CONSULTANT BEFORE WORK PROCEEDS

Legend

Site Location



ITM Coordinates: 683027 E, 670998 N

Rev	Description	Date

EARTH SCIENCE PARTNERSHIP

CONSULTING ENGINEERS, GEOLOGISTS & ENVIRONMENTAL SCIENTISTS
 Tonranny Westport Co. Mayo
 tel: +353(0)98 28999 fax: +353(0)98 24727
 e-mail: earthsciencepartnership@gmail.com
 Ordnance Survey Ireland Copyright Licence No: EN 0021419

Client: Kilcarrig Quarries

Project: Environmental Impact
 Assessment Report to
 Accompany a Substitute
 Consent Application for Quarry
 Located at Roscat, Tullow,
 Co. Carlow

Title: Site Location Map

Drawn By: Sean O' Donnell

Checked By: Patrick O' Donnell

Scale: 1:50,000 @ A4 Date: Jan. 2019

Job No: EI 061 Rev: 0

Figure 1.1

2 METHODOLOGY

2.1 Legislative Context

It should also be noted that, although the legislation described below refers to proposed (future) developments, this rNIS is a retrospective assessment and as such, the same relevant methodology and assessment guidelines have been adhered to as are adhered to when assessing a proposed (future) development. To remain in context however, the text within the project description and assessment of the report is presented in the past tense.

2.1.1 Article 6 of the Habitats Directive

Article 6(3) of the Habitats Directive requires that, in relation to European designated sites (i.e. SACs and SPAs that form the Natura 2000 network), *"any plan or project not directly connected with or necessary to the management of the site but likely to have a significant effect thereon, either individually or in combination with other plans or projects, shall be subject to **appropriate assessment** of its implications for the site in view of the site's conservation objectives"*.

A competent authority (e.g. a Local Authority) can only agree to a plan or project after having determined that it will not adversely affect the integrity of the site concerned. Under Article 2(2) of the Directive, a network of sites of nature conservation importance have been identified by each Member State as containing specified habitats or species requiring to be maintained or returned to favourable conservation status. In Ireland and Northern Ireland, the network consists of Special Areas of Conservation (SACs) and Special Protection Areas (SPAs), and also candidate sites, which form the Natura 2000 network.

Under Article 6(4) of the Directive, if adverse impacts are likely, and in the absence of alternative options, a plan or project must nevertheless proceed for imperative reasons of overriding public interest (IROPI), including social or economic reasons, a Member State is required to take all compensatory measures necessary to ensure the overall integrity of the Natura 2000 site. The European Commission are required to be informed of any compensatory measures adopted, unless a priority habitat type or species is present and in which case, an opinion from the European Commission is required in advance (unless for human health or public safety reasons, or of benefit to the environment).

2.1.2 Appropriate Assessment Process

Guidance on the Appropriate Assessment (AA) process was produced by the European Commission in 2002, which was subsequently developed into guidance specifically for Ireland by the Department of Environment, Heritage and Local Government (DEHLG, 2009). These guidance documents identify a staged approach to conducting an AA, as shown Figure 2.1.




Figure 2.1: The Appropriate Assessment Process (from: Appropriate Assessment of Plans and Projects in Ireland - Guidance for Planning Authorities, DEHLG, 2009)

2.1.2.1 Screening for Appropriate Assessment

This stage examines the likely effects of a project either alone or in combination with other projects upon a Natura 2000 site and considers whether it can be objectively concluded that these effects will not be significant. In the light of the recent case-law ruling [C-323/17], “it is not appropriate, at the screening stage, to take account of the measures intended to avoid or reduce the harmful effects of the plan or project on that site”. As such, the current screening for appropriate assessment does not take account of measures, such as “best practice construction methods”, intended to avoid or reduce the project’s harmful effects on relevant Natura 2000 sites. This evaluation is presented in a Screening for Appropriate Assessment.

2.1.2.2 Appropriate Assessment

Should the screening stage find that the proposed project or plan has the potential to cause harmful effects on Natura 2000 sites, the impact of the project on the integrity of the Natura 2000 site is considered with respect to the conservation objectives of the site and to its structure and function. Mitigation measures should be applied to the point where no adverse impacts on the site(s) remain. This evaluation is presented in a Natura Impact Statement (NIS).

2.1.2.3 Alternative Solutions

Should the Appropriate Assessment determine that adverse impacts are likely upon a Natura 2000 site, this stage examines alternative ways of implementing the project that, where possible, avoid these adverse impacts. For the avoidance of doubt, the developer does not purport to place reliance on this stage.

2.1.2.4 Imperative Reasons of Overriding Public Interest (IROPI)

Assessment where no alternative solutions exist and where adverse impacts remain: Where imperative reasons of overriding public interest (IROPI) exist, an assessment to consider whether compensatory measures will or will not effectively offset the damage to the Natura site will be necessary. European case law highlights that consideration must be given to alternatives outside the project area in carrying out the IROPI test. It is a rigorous test which projects are generally considered unlikely to pass. In any event, the developer does not purport to place any reliance on Stage 4.

2.2 Relevant Guidance

In the preparation of this rNIS, regard has been given to the EU Habitats Directive (1), Part XAB of the Planning and Development Act 2000, and to the relevant guidance, in particular:

- European Commission (2001). *Assessment of Plans and Projects significantly affecting Natura 2000 Sites: Methodological guidance on the provisions of Article 6(3) and (4) of the Habitats Directive 92/43/EEC*.
- European Commission (2000). *Managing Natura 2000 Sites: The Provisions of Article 6 of the 'Habitats Directive' 92/43/EEC*.
- National Parks and Wildlife Service, Department of the Environment, Heritage and Local Government, Dublin (2009). *Appropriate Assessment of Plans and Projects in Ireland: Guidance for Planning Authorities*.

Throughout the Stage One screening report (Section 5), the line items in *italics* refer to suggested instructions for information to be contained in a screening assessment, from the guidance document '*Assessment of Plans and Projects significantly affecting Natura 2000 Sites: Methodological guidance on the provisions of Article 6(3) and (4) of the Habitats Directive 92/43/EEC*', European Commission, 2001.

In order to identify potential 'In Combination Effects', other key plans and projects were identified for this area. The following plans, projects or data sources have been considered:

- Carlow County Council planning database (<http://www.carlow.ie/en/services/planning/onlineplanningsystems/>)
- NPWS website (<http://webgis.npws.ie/npwsviewer/>)
- EPA website (<https://gis.epa.ie/EPAMaps/>)
- National Biodiversity Data Centre (<http://maps.biodiversityireland.ie/#/Home>)

2.3 Consultation

A Screening and Scoping document in relation to the full remedial EIAR, outlining the complete project description and potential effects of the historic development, was forwarded to the DAU 26th October 2018, which in turn disseminated the information to the NPWS. At the time of writing (Jan 2019), no formal written response to consultation has been received from NPWS.

The Screening and Scoping document was also sent to a number other relevant statutory and non-statutory bodies with an ecological interest such as Inland Fisheries Ireland (IFI) and An Taisce. A formal response was received from IFI and is available for reference in Appendix II. No formal response to consultation has yet been received from An Taisce.

3 PROJECT DESCRIPTION

3.1 Existing Substitute Consent Site

The existing site is located in the townland of Roscat approximately 3 km south west of Tullow and 2 km east of Rathoe. The N81 national road which connects Tullow with the N80 at Ballon passes in a north-south direction 1.5 km east of the site.

Vehicular access is off a local road and via a c.1km long gated laneway that provides access to the pit and surrounding lands. The site is surrounded by agricultural fields and a farm is located nearby.

This Substitute Consent application consists of an existing 4.7 hectare sand and gravel pit.

Material was extracted by excavators and exported to a processing plant offsite. On occasion, material was passed through mobile processing plant onsite, which processed the material into various grades of aggregate, as and when required. Material was processed into various grades depending on market demand for aggregate and was stockpiled close to the processing plant or at designated stockpiles.

Plant and machinery which operated in the application area consisted of tracked excavators, wheel loaders, trucks and mobile processing plant. Ancillary plant such as a water bowser for dust suppression was deployed where required.

As all production was a dry process, surface water percolated through the pit floor or flows to the settlement lagoons located in the south eastern corner of the pit where water is retained and used for dust suppression. Excess water is discharged off-site.

4 RECEIVING ENVIRONMENT

4.1 Natura 2000 Sites

4.1.1 Identification of Relevant Natura 2000 Sites

The DEHLG (2009) guidance identifies that the Appropriate Assessment of a plan or project should consider the following Natura 2000 sites:

- Any Natura 2000 sites within or adjacent to the plan or project area.
- Any Natura 2000 sites within the likely zone of impact of the plan or project. This is dependent on the nature and scale of the plan, with 15km generally recommended for plans, but potentially much less for projects.
- Any Natura 2000 sites that are more than 15km from the plan or project area, but may potentially be impacted upon, for example, through a hydrological connection.

A 15km buffer zone has been applied for the search criteria for the historical development, including those with a surface and/or ground water hydrological connection.

A total of 3 Natura 2000 sites were identified during the desk study as being located within 15km (from closest to farthest distance). The sites are as follows (also see Figure 4.1):

1. Slaney River Valley SAC (Site Code: 000781)
2. River Barrow and River Nore SAC (Site Code: 002162)
3. Blackstairs Mountains SAC (Site Code: 000770)

The pathways for potential historical effects are identified in Table 4.1.

Earth Science Partnership
Tonranny, Westport, Co. Mayo
Tel: +353 98 28999
Email: earthsciencepartnership@gmail.com

Client: Kilcarrig Quarries Ltd

Project: Substitute Consent application for Roscat Quarry,
Tullow, Co. Carlow

Title: Natura 2000 within 15km of the application area

Drawn by: Sarah Ingham

Checked by: Patrick O'Donnell

Scale: 1:18k @ A4

Date: 03/01/2019

Job No.: EI 061

Rev.: 1.0

Figure: 4.1

Legend

- Substitute Consent Application Area Location
- ▨ SAC
- ⋯ 5, 10 and 15km Buffer



THIS DRAWING AND ANY DESIGN HEREON IS THE COPYRIGHT
OF THE CONSULTANT AND MAY NOT BE COPIED, REPRODUCED
OR TRANSMITTED IN ANY FORM WITHOUT THE WRITTEN
CONSENT OF THE CONSULTANT.




Table 4.1: Identification of pathways for likely significant historical effects on Natura 2000 sites within 15km of the existing pit location and/or where a hydrological connection exists between the existing development and the Natura 2000 site.

Natura 2000 Site	Primary reason(s) for selection of site	Approximate distance from proposed application area	Further Screening Required (Y/N)
Slaney River Valley SAC (Site Code: 000781)	<p>Priority Annex I Habitats [91E0] Alluvial Forests*</p> <p>Annex I Habitats [1130] Estuaries [1140] Tidal Mudflats and Sandflats [1330] Atlantic salt meadows (<i>Glauco-Puccinellietalia maritima</i>) [1410] Mediterranean salt meadows (<i>Juncetalia maritimi</i>) [3260] Floating River Vegetation [91A0] Old Oak Woodlands</p> <p>Annex II Species [1029] Freshwater Pearl Mussel (<i>Margaritifera margaritifera</i>) [1095] Sea Lamprey (<i>Petromyzon marinus</i>) [1096] Brook Lamprey (<i>Lampetra planeri</i>) [1099] River Lamprey (<i>Lampetra fluviatilis</i>) [1103] Twaite Shad (<i>Alosa fallax</i>) [1106] Atlantic Salmon (<i>Salmo salar</i>) [1355] Otter (<i>Lutra lutra</i>) [1365] Common (Harbour) Seal (<i>Phoca vitulina</i>)</p>	<p>This SAC is located approximately 1.8km east of the proposed application area.</p>	<p>No. The Substitute Consent application area lies outside the River Slaney catchment area with no groundwater or hydrological pathways to the Slaney River Valley SAC. At a distance of 1.8km, this European site is also outside the maximum extent where it would be anticipated that there would have been any likely measureable significant effects from disturbance (i.e. noise, vibration and human and visual disturbance) and from any deposition of dust.</p> <p>It is considered therefore that this European site can be screened out from any further evaluation as the existing development is not likely to have had any significant effect on the integrity of the Slaney River Valley SAC during its operation, or on any of the qualifying habitats and/or species for which this site has been designated as being of European importance.</p> <p>Therefore, further screening is not required.</p>
River Barrow and River Nore SAC (Site Code: 002162)	<p>Priority Annex I Habitats [7220] Petrifying Springs* [91E0] Alluvial Forests*</p>	<p>This SAC is located approximately 12.7km west (direct distance) and 15.5km (hydrological distance) north west of the proposed</p>	<p>Yes. A hydrological connection exists between the River Barrow and River Nore SAC and the Substitute Consent application area via the Roscat Stream which is located approximately</p>

Natura 2000 Site	Primary reason(s) for selection of site	Approximate distance from proposed application area	Further Screening Required (Y/N)
	<p>Annex I Habitats</p> <p>[1130] Estuaries [1140] Tidal Mudflats and Sandflats [1170] Reefs [1310] Salicornia Mud [1330] Atlantic Salt Meadows [1410] Mediterranean Salt Meadows [3260] Floating River Vegetation [4030] Dry Heath [6430] Hydrophilous Tall Herb Communities [91A0] Old Oak Woodlands</p> <p>Annex II Species</p> <p>[1016] Desmoulin's Whorl Snail (<i>Vertigo moulinsiana</i>) [1029] Freshwater Pearl Mussel (<i>Margaritifera margaritifera</i>) [1092] White-clawed Crayfish (<i>Austropotamobius pallipes</i>) [1095] Sea Lamprey (<i>Petromyzon marinus</i>) [1096] Brook Lamprey (<i>Lampetra planeri</i>) [1099] River Lamprey (<i>Lampetra fluviatilis</i>) [1103] Twaite Shad (<i>Alosa fallax</i>) [1106] Atlantic Salmon (<i>Salmo salar</i>) [1355] Otter (<i>Lutra lutra</i>) [1421] Killarney Fern (<i>Trichomanes speciosum</i>) [1990] Nore Freshwater Pearl Mussel (<i>Margaritifera durrovensis</i>)</p>	application area.	<p>520m to the west of the proposed application area. The Roscat Stream flows in a north-westerly direction towards its confluence with the Burren River, which flows into the River Barrow in Carlow town. The total hydrological distance between the application area and the SAC is 15.5km, which provides a potential, albeit weak, source-pathway-receptor link between the two.</p> <p>As such, potential effects of the historical works on this SAC are evaluated further in the following screening assessment.</p>

Natura 2000 Site	Primary reason(s) for selection of site	Approximate distance from proposed application area	Further Screening Required (Y/N)
Blackstairs Mountains SAC (Site Code: 000770)	Annex I Habitats [4010] Wet Heath [4030] Dry Heath	This SAC is located approximately 14.2km south of the proposed application area.	No. Owing to the considerable distance and absence of a hydrological connection between this SAC and the Substitute Consent application area, there are no pathways identified for likely significant direct or indirect effects which could have occurred on any of the qualifying interests of this Natura 2000 site during the operation of the pit. Therefore, further screening is not required.

Given that two of the three Natura 2000 sites identified in this scoping study are neither hydrologically nor ecologically connected with the Substitute Consent application area, no pathways for direct or indirect likely significant effects exist. As such, the following sites have been excluded at the scoping stage and are therefore no longer considered in this report:

1. Slaney River Valley SAC (Site Code: 000781)
2. Blackstairs Mountains SAC (Site Code: 000770)

The following Natura 2000 site will undergo further Stage 1 Screening for Appropriate Assessment:

1. River Barrow and River Nore SAC (Site Code: 002162)

4.1.2 River Barrow and River Nore SAC (Site Code: 002162)

This site consists of the freshwater stretches of the Barrow and Nore River catchments as far upstream as the Slieve Bloom Mountains, and it also includes the tidal elements and estuary as far downstream as Creadun Head in Waterford. The site passes through eight counties – Offaly, Kildare, Laois, Carlow, Kilkenny, Tipperary, Wexford and Waterford.

The site is of considerable conservation significance for the occurrence of good examples of habitats and of populations of plant and animal species that are listed on Annexes I and II of the E.U. Habitats Directive. Furthermore it is of high conservation value for the populations of bird species that use it. The occurrence of several Red Data Book plant species including three rare plants in the salt meadows and the population of the hard water form of the Freshwater Pearl Mussel, which is limited to a 10 km stretch of the Nore, add further interest to this site.

Land use at the site consists mainly of agricultural activities – mostly intensive in nature and principally grazing and silage production. The main threats to the site and current damaging activities include high inputs of nutrients into the river system from agricultural run-off and several sewage plants, over-grazing within the woodland areas, and invasion by non-native species, for example Cherry Laurel (*Prunus laurocerasus*) and Rhododendron (*Rhododendron ponticum*). The water quality of the site remains vulnerable.

Qualifying Interests of River Barrow and River Nore SAC

The features of qualifying interest of the SAC are outlined in Table 4.2.

Table 4.2: Qualifying interests for River Barrow and River Nore SAC (* denotes priority Annex I habitats)

	Natura Code	Item Description
Annex I Habitats	1130	Estuaries
	1140	Tidal Mudflats and Sandflats
	1170	Reefs
	1310	Salicornia Mud
	1330	Atlantic Salt Meadows
	1410	Mediterranean Salt Meadows
	3260	Floating River Vegetation
	4030	Dry Heath
	6430	Hydrophilous Tall Herb Communities
	7220	Petrifying Springs*
	91A0	Old Oak Woodlands
	91E0	Alluvial Forests*
Annex II Species	1016	Desmoulin's Whorl Snail (<i>Vertigo moulinsiana</i>)
	1029	[Freshwater Pearl Mussel (<i>Margaritifera margaritifera</i>)
	1092	White-clawed Crayfish (<i>Austropotamobius pallipes</i>)
	1095	Sea Lamprey (<i>Petromyzon marinus</i>)
	1096	Brook Lamprey (<i>Lampetra planeri</i>)
	1099	River Lamprey (<i>Lampetra fluviatilis</i>)
	1103	Twaite Shad (<i>Alosa fallax</i>)
	1106	Atlantic Salmon (<i>Salmo salar</i>)
	1355	Otter (<i>Lutra lutra</i>)
	1421	Killarney Fern (<i>Trichomanes speciosum</i>)
1990	Nore Freshwater Pearl Mussel (<i>Margaritifera durrovensis</i>)	

Conservation Objectives

Site-specific conservation objectives aim to define favourable conservation condition for the particular habitat or species at that site (NPWS, 2011)¹. Please see Appendix I for the complete conservation objectives document. The generic conservation objectives for the River Barrow and River Nore SAC are:

Objective 1: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and/or the Annex II species for which the SAC has been selected (as listed in Table 4.4).

Objective 2: To maintain the extent, species richness and biodiversity of the entire site.

Objective 3: To establish effective liaison and co-operation with landowners, legal users and relevant authorities.

¹ NPWS (2011) Conservation Objectives: River Barrow and River Nore SAC 002162. Version 1.0. National Parks and Wildlife Service, Department of Arts, Heritage and the Gaeltacht.

5 STAGE 1 – SCREENING FOR APPROPRIATE ASSESSMENT

5.1 Introduction

The existing sand and gravel pit development at Roscat, Tullow Co. Carlow is not connected with, or necessary for the nature conservation management of River Barrow and River Nore SAC. Consequently, the screening will focus on evaluating the likely effects of the historical works during the operational period at the pit on the designated Natura site identified.

5.2 Potential Risks to Natura 2000 Sites

This section identifies the potential risk which may have arisen as result of the historical works and then goes on to identify how these could have potentially effected the qualifying interests within the identified Natura 2000 site. The pathways and mechanisms of exposure to each of the identified risks are evaluated. The significance of potential effects is evaluated, with any potential in-combination effects also identified.

5.2.1 Potential Risks as a Result of the Existing Project

The following are potential risks to the River Barrow and River Nore SAC that may have arisen as a result of the historical sand and gravel extraction works:

- ***Deterioration in water quality*** – although the hydrological distance between the Substitute Consent development site and this SAC is considerable (15.5km), the precautionary approach has been followed and as such, the existing development had the potential to cause indirect impacts to the qualifying interests of the identified SAC, especially freshwater aquatic species such as Otter and salmonid species, through the contamination of surface water within the hydrologically connected Roscat Stream by virtue of the following:
 1. The release and percolation of hydrocarbons into the groundwater system as a result of excavation activities.
 2. The creation of suspended solids in runoff waters.

5.3 Evaluation of Likely Effects

Table 5.1: Evaluation of potential risks and their level of potential effect on features of qualifying interest for the River Barrow and River Nore SAC

Qualifying Interests	Evaluation of Potential Risks	Potential Exposure to Risk and Mechanism of Effect, if known
Estuaries	These primarily coastal habitats occur within the boundaries of the SAC, which is located at a hydrological distance of 15.5km at its closest point to the Substitute Consent application area. Owing to the dilution effect at this distance, there would have been no pathways for significant direct or indirect effects on these habitats as a result of the historical project.	The risk to these coastal habitats is not considered to have been significant given the distances involved and that the surface hydrological connection between the SAC and the Substitute Consent application area is weak. Therefore, no pathway for likely significant effects exists. Conclusion: <i>No likely significant effect</i>
Tidal Mudflats and Sandflats		
Reefs		
Salicornia Mud		
Atlantic Salt Meadows		
Mediterranean Salt Meadows		
Dry Heath	These primarily terrestrial habitats occur within the boundaries of the SAC, which, as stated above, is located at a hydrological distance of 15.5km at its closest point to the Substitute Consent application area. Owing to the dilution effect at this distance, there would have been no pathways for significant direct or indirect effects on these habitats as a result of the historical project.	The risk to these terrestrial habitats is not considered to have been significant given the distances involved and that the surface hydrological connection between the SAC and the Substitute Consent application area is weak. Therefore, no pathway for likely significant effects exists. Conclusion: <i>No likely significant effect</i>
Hydrophilous Tall Herb Communities		
Petrifying Springs*		
Old Oak Woodlands		
Alluvial Forests*		
Floating River Vegetation	Although a river corridor survey was not undertaken on the Roscat Stream as part of this evaluation, given the Roscat Stream has a current Q-Value of 4-5 (High Quality) (EPA, 2016), it is possible that this stream could have contained floating river vegetation during the time of the operation of the pit. As such, following the precautionary principle, pathways for potentially significant indirect effects on this qualifying interest of the SAC as a result of the historical works exist by virtue of	The risk to this aquatic habitat is considered to have been significant given the surface water connectivity between the Substitute Consent application area and the Roscat Stream and its subsequent connectivity with the SAC. Conclusion: <i>Likely significant effect</i>

Qualifying Interests	Evaluation of Potential Risks	Potential Exposure to Risk and Mechanism of Effect, if known
	risks to the surface water system at the Substitute Consent application area.	
Desmoulin's Whorl Snail (<i>Vertigo moulinsiana</i>) Killarney Fern (<i>Trichomanes speciosum</i>)	These terrestrial hydrology dependant species occur within the boundaries of the SAC, which is at direct distance 12.7km at its closest point to the Substitute Consent application area and a hydrological distance of 15.5km. Owing to the dilution effect at this distance, there would have been no pathways for significant direct or indirect effects on these habitats as a result of the historical project.	The risk to these terrestrial species is not considered to have been significant given the distances involved and that the surface hydrological connection between the SAC and the Substitute Consent application area is weak. Therefore, no pathway for likely significant effects exists. Conclusion: No likely significant effect
Freshwater Pearl Mussel (<i>Margaritifera margaritifera</i>) Nore Freshwater Pearl Mussel (<i>Margaritifera durrovensis</i>) White-clawed Crayfish (<i>Austropotamobius pallipes</i>) Sea Lamprey (<i>Petromyzon marinus</i>) Brook Lamprey (<i>Lampetra planeri</i>) River Lamprey (<i>Lampetra fluviatilis</i>) Twaite Shad (<i>Alosa fallax</i>) Atlantic Salmon (<i>Salmo salar</i>) Otter (<i>Lutra lutra</i>)	Given the Roscat Stream's current Q-Value rating of 4-5 (High Quality) (EPA, 2016), it is possible that these aquatic species of qualifying interest occurred within the stream during the time of the operation of the pit, which is hydrologically connected (albeit at distance) with the SAC. The Substitute Consent application area is also located within the Barrow <i>Margaritifera</i> Sensitive Area (NPWS, 2017). The Barrow <i>Margaritifera</i> Sensitive Area is categorised as a "catchment with previous records of <i>Margaritifera</i> but its current status is unknown". The closest <i>Margaritifera</i> Sensitive Area designated for catchments of SAC populations listed under S.I. 296 of 2009 is the Slaney-Derreen catchment which is located approximately 1.5km to the east, however, is not hydrologically connected to the Substitute Consent application area.	The risk to these aquatic species is considered to have been significant given the surface water connectivity between the Substitute Consent application area and the Roscat Stream. Conclusion: Likely significant effect

Qualifying Interests	Evaluation of Potential Risks	Potential Exposure to Risk and Mechanism of Effect, if known
	<p>Following the precautionary principle, pathways for potentially significant indirect effects on these qualifying interests of the SAC which could have occurred as a result of the historical works existed by virtue of risks to the surface water system at the Substitute Consent application area.</p> <p>These risks also had the potential to cause indirect effects on Otter through potential effects on prey species via deterioration in water quality.</p>	

5.4 Potential In-combination Effects

It is a requirement of Article 6(3) of the Habitats Directive that an assessment is conducted to determine if significant effects of a plan or project are likely alone and also in combination with other plans or projects. The following section discusses potential in-combination effects that may have arisen during the period of operation of the existing pit.

5.4.1 Capped Landfill Site

An unauthorised capped landfill site of approximately 1.6ha in area and 8m in depth is located approximately 1.2km to the south of the Substitute application area. A review of aerial mapping from 1995 shows that this landfill site was in existence at that time (<http://map.geohive.ie/>). Given its unauthorised status, a Tier 3 Risk Assessment and Restoration Plan was commissioned by Carlow County Council and undertaken by RPS in 2011, in accordance with the “*EPA Code of Practice Environmental Risk Assessment for Unregulated Waste Disposal Sites*”.

A detailed qualitative risk assessment was carried out to determine the potential risks to human health and controlled waters for the contaminants identified as being of concern (e.g. heavy metals, hydrocarbons, ammonia, volatile organic compounds and ground gases). Groundwater monitoring was undertaken as part of the risk assessment. The assessment concluded that the site then represented a moderate risk to groundwater quality via leachate being produced at the site. However, it has since been shown that the installation of the infiltration barrier led to long term improvements in groundwater quality to an acceptable level.

From the assessment undertaken here, it is considered that it can be clearly demonstrated that the existing development at Roscat Sand and Gravel Pit did not act in-combination with the capped landfill site, as described above, to result in any effects on the River Barrow and River Nore SAC or on any of its qualifying features for which this site is of European importance.

5.4.2 Existing Ardristan Sand and Gravel Pit (QY 27)

Ardristan Sand and Gravel Pit is located approximately 1.5km to the south of the Substitute Consent application area. Permission for the development of the sand and gravel pit to include continuation of existing extraction, screening crushing, washing and ancillary activities and phased restoration of the pit to agricultural afteruse at Ardristan, Tullow, Co. Carlow was granted with a number of planning conditions in 2007 under PL 07/769; ABP Ref: 01.232014. As such, this sand and gravel pit was in existence during the period operation of the Substitute Consent area. Of particular relevance to the current application is Condition 11, which directs the implementation of ground water monitoring scheme at the consented site and the submission of the results of same to Carlow County Council.

Groundwater quality monitoring was carried out on behalf of the developer by trained personnel employed by SLR Consulting Ireland (formerly John Barnett and Associates). Groundwater quality was monitored at 3 no. locations. The report concluded results indicate that the operation does not adversely impact on water quality in the locality.

In 2013, a remedial Natura Impact Statement (rNIS) was undertaken on this site, which considered the potential effects associated with the extraction of sand and gravel at the existing Ardristan sand and gravel pit on the River Barrow and River Nore SAC. Although 2013 is 5 years after the last operation of the Roscat pit, the results of the rNIS are still considered to be relevant to the current Substitute Consent application from a cumulative impact perspective.

The rNIS concluded that the extraction of sand and gravel had not had any significant effects on the River Barrow and River Nore SAC, nor is it likely to have any effects through any continuation of quarrying operations to affect the integrity of this relevant European site, or on any of its qualifying features for which this site is of European importance, in light of its conservation objectives.

More recently a further planning application has been submitted to Carlow Co. Co. for permission to import inert waste consisting of soil and stone material to improve the land for agricultural use at this site (PL Ref: 18/220). At the time of writing (Jan 2019), this application is currently at the clarification of further information stage. The Screening for Appropriate Assessment which was undertaken as part of the application was reviewed and concludes that there no significant direct or indirect effects on the integrity of the Natura 2000 network would occur as a result of the proposal.

As such, it is considered that it can be clearly demonstrated that the existing pit at Roscat Sand and Gravel Pit did not act in-combination with the previous operations at the Ardristan Sand and Gravel Pit, as described above, to result in any effects on the River Barrow and River Nore SAC or on any of its qualifying features for which this site is of European importance.

5.5 Screening for Appropriate Assessment Matrix

The following matrix summarises the screening process that has been undertaken as part of this evaluation.

Table 5.2: Screening for AA Matrix

Screening Criterion	Screening Result
<i>Brief description of the project or plan</i>	This Substitute Consent application consists of the extraction of material over an area of 4.7 hectares down to a depth of approximately 63 mOD at a sand and gravel pit at Roscat, Tullow, Co. Carlow.
<i>Brief description of the Natura 2000 sites</i>	<ul style="list-style-type: none"> • River Barrow and River Nore SAC (Site Code: 002162) <p><i>Relevant features of the sites are described in Section 5.1.1.</i></p>
<i>Describe the individual elements of the project (either alone or in combination with other plans or projects) likely to give rise to impacts on the Natura 2000 site.</i>	<p><u>Indirect effects on water quality</u></p> <p>Given the Roscat Stream's Q-Value rating of 4-5 (High Quality) (EPA, 2016), it is possible that aquatic species and habitats of qualifying interest occurred within the stream or in the rivers downstream of the existing development during its period of operation, which are hydrologically connected with the SAC. As such, pathways for potentially significant indirect effects on the surface water dependent freshwater aquatic qualifying interests of the SAC as a result of the Substitute Consent development existed during its period of operation by virtue of risks to the surface and groundwater system at the application area. These risks also had the potential to cause indirect effects on Otter through potential effects on prey species via deterioration in water quality.</p> <p>As such, where remedial measures are not applied (See Section 6.1), previous extraction works may have the potential to result in significant indirect effects through the release of sediment in surface water run-off affecting water dependent species and habitats associated with the River Barrow and River Nore SAC.</p> <p>Conclusion: <i>Potential for significant effects</i></p>

5.6 Screening Statement and Conclusions

The scoping assessment detailed in Section 4.1.1 examined potential effects on the Natura 2000 designated SACs and SPAs within a 15km radius of the Substitute Consent application area. The following 3 designated sites were scoped as part of this evaluation:

1. Slaney River Valley SAC (Site Code: 000781)
2. River Barrow and River Nore SAC (Site Code: 002162)
3. Blackstairs Mountains SAC (Site Code: 000770)

Following consideration of the nature and scale of the project, the hydrological/ecological connectivity between the application area location and the designated sites, the site specifics of the Substitute Consent application area location and the particular qualifying interests/special conservation interests of the Natura 2000 sites, **the following 2 no. designated sites were excluded at the scoping stage** of the screening:

1. Slaney River Valley SAC (Site Code: 000781)
2. Blackstairs Mountains SAC (Site Code: 000770)

Consequently, likely significant effects on the qualifying interests of the **River Barrow and River Nore SAC (Site Code: 002162)** were evaluated within this screening report.

Following the screening process, it has been determined that the **potential for significant effects on the River Barrow and River Nore SAC remain by virtue of indirect effects resulting in a reduction in water quality within the SAC** and by consequence, the potential for significant effects on water dependant habitats and species as a result of works undertaken during the period of operation of the pit exists.

The recommendation of the screening process is therefore to proceed to Stage 2 Appropriate Assessment for River Barrow and River Nore SAC (See Section 5.4) in terms of the potential for significant indirect effects.

6 STAGE 2 – APPROPRIATE ASSESSMENT REPORT

This section of the remedial Natura Impact Statement addresses the possibility of there having been a significant effect or effects on the following European site which was identified during Stage One screening (Section 5):

- River Barrow and River Nore SAC (Site Code: 002162)

It should be noted here that there is no specific guidance for remedial Appropriate Assessment for the Substitute Consent process. As such, the guidance for Appropriate Assessment (European Commission (2018)) is adhered to. This guidance is set out in the future tense to deal with proposed plans or projects, however, in order to accommodate the retrospective nature of the Substitute Consent process, the guidance questions (in *italics* below) have been altered to the past tense.

6.1 Assessment of the Effects of the Project or Plan on the Integrity of the Sites

‘Describe the elements of the project or plan (alone or in combination with other projects or plans) that are likely to have given rise to significant effects on the site (from screening assessment)’

The historical works had the potential to cause a reduction in water quality within the River Barrow and River Nore SAC and by consequence, the potential for significant effects on water dependant habitats and species existed.

6.2 The Conservation Objectives of the Natura 2000 Sites

‘Set out the conservation objectives of the site’.

The conservation objectives of the European site concerned are to maintain the favourable conservation status of the key species and habitats for which the site has been designated. These are laid out in Table 6.1 below.

Table 6.1: Key surface and groundwater dependant species and habitats of qualifying interest of the Natura 2000 site which could have been impacted upon by the existing development

Designated Site	Conservation Objectives
River Barrow and River Nore SAC	<p>Objective: To maintain or restore the favourable conservation condition of the Annex I habitat(s) and the Annex II species for which the SAC has been selected:</p> <ul style="list-style-type: none"> • Floating River Vegetation • Freshwater Pearl Mussel (<i>Margaritifera margaritifera</i>) • White-clawed Crayfish (<i>Austropotamobius pallipes</i>) • Sea Lamprey (<i>Petromyzon marinus</i>) • Brook Lamprey (<i>Lampetra planeri</i>) • River Lamprey (<i>Lampetra fluviatilis</i>) • Twaite Shad (<i>Alosa fallax</i>)

Designated Site	Conservation Objectives
	<ul style="list-style-type: none"> • Atlantic Salmon (<i>Salmo salar</i>) • Otter (<i>Lutra lutra</i>) • Nore Freshwater Pearl Mussel (<i>Margaritifera durrovensis</i>) <p><small>NPWS (2011) Conservation Objectives: River Barrow and River Nore SAC 002162. Version 1.0. National Parks and Wildlife Service, Department of Arts, Heritage and the Gaeltacht. [accessed online on 04/01/2019]</small></p>

6.3 Potential Impacts on Key Species and Key Habitats

‘Describe how the project or plan would have affected key species and key habitats. Acknowledge uncertainties and any gaps in information’.

In the scenario of a release of hydrocarbons from vehicles and machinery and stored fuels during the extraction works within the existing pit during its period of operation, there would have been a potential for significant indirect effects on the water dependant habitats and species of qualifying interest within the River Barrow and River Nore SAC.

A reduction in water quality due to hydrocarbons has the potential result in effects on salmonids and plant species within Roscat stream.

In addition, although Otter was not recorded during the site survey, given their ubiquitous nature in the Irish landscape, it is likely that they would have been present in Roscat stream during the period of operation of the pit. Otters may have been secondarily affected by a reduction in water quality through a reduction in potential prey should fish have been impacted upon as a result of hydrocarbon runoff to groundwater.

Furthermore, the potential for increased silt content in runoff could have degraded local surface water quality. Silt has the potential to clog salmonid spawning beds and juvenile salmonids are particularly sensitive to siltation of gill structures. Similarly, plant and macro-invertebrate communities could have been blanketed over and this could have led to loss or degradation of valuable habitat.

6.4 Potential Impacts on the Integrity of the Sites

‘Describe how the integrity of the site (determined by structure and function and conservation objectives) would have been likely to have been affected by the project and plan (e.g. loss of habitat, disturbance, disruption, chemical changes, hydrological changes and geological changes etc.). Acknowledge uncertainties and any gaps in information’.

The integrity of the River Barrow and River Nore SAC could have been indirectly affected by the existing development through a reduction in water quality and foraging potential for aquatic species. This could in turn have led to reduced numbers or reduced breeding success of these species which are a qualifying interest of the SAC. Table 6.2 presents specific impacts detailing their importance, magnitude and level of significance.

Table 6.2: Potential impacts of the existing development on the aquatic environment which is hydrologically connected with the River Barrow and River Nore SAC via surface and groundwater

Activity	Attribute	Character of potential impact	Potential consequential impact on aquatic ecology	Importance of attribute	Magnitude of potential impact	Term	Significance of potential impact
Compaction of pit floor and hardstanding pad	Roscat stream (River Barrow and River Nore SAC)	Silt-laden runoff from pit floor. The increased silt content in runoff had potential to degrade local surface water quality.	Silt has the potential to clog salmonid spawning beds and juvenile salmonids are particularly sensitive to siltation of gill structures. Similarly, plant and macro-invertebrate communities can be blanketed over and this could have led to loss or degradation of valuable habitat.	Very High	Small	Long-term	Moderate

Activity	Attribute	Character of potential impact	Potential consequential impact on aquatic ecology	Importance of attribute	Magnitude of potential impact	Term	Significance of potential impact
Stockpiling of topsoil/subsoil/screenings	Roscat stream (River Barrow and River Nore SAC)	Silt-laden runoff from stockpiles. The increased silt content in runoff had potential to degrade local surface water quality.	As above.	Very High	Small	Medium-term	Moderate
Removal of overburden	Roscat stream/Aquifer (River Barrow and River Nore SAC)	Increase in vulnerability of underlying aquifer.	As above.	Very High	Small	Long-term	Moderate
Storage of hydrocarbons; leakages from machinery; spillages during refuelling	Roscat stream/Aquifer (River Barrow and River Nore SAC)	Runoff/recharge may have contained hydrocarbons.	<p>A reduction in water quality due to hydrocarbons had the potential result in effects on salmonids and plant species within Roscat stream.</p> <p>In addition, although Otter was not recorded during the site survey, given their ubiquitous nature in the Irish landscape, it is likely that they were present in Roscat stream during the period of operation of the pit. Otters may have been secondarily</p>	Very high	Small	Temporary	Moderate

Activity	Attribute	Character of potential impact	Potential consequential impact on aquatic ecology	Importance of attribute	Magnitude of potential impact	Term	Significance of potential impact
			affected by a reduction in water quality through a reduction in potential prey should fish be impacted upon as a result of hydrocarbon runoff to groundwater.				

6.5 Mitigation/Remedial Measures

‘Describe what remedial measures were or will be introduced to avoid or reduce the adverse effects on the integrity of the site. Acknowledge uncertainties and any gaps in information’

- *List measures which were or will be introduced*
- *Explain how the measures avoided or will avoid the adverse effects on the integrity of the site*
- *Explain how the measures reduced or will reduce the adverse effects on the integrity of the site*
- *Provide evidence of how they were or will be implemented and by whom.*

Table 6.3: Mitigation/Remedial Measures to reduce or avoid adverse effects on the River Barrow and River Nore SAC

Activity	Attribute	Character of potential impact	measure	Residual impact
Compaction of pit floor and hardstanding pad	Roscat stream (River Barrow and River Nore SAC)	Silt-laden runoff from pit floor. The increased silt content in runoff had potential to degrade local surface water quality.	Runoff passes through a series of settlement ponds which have been shown to clarify the water prior to it crossing the site boundary. Results from sampling show that water quality in the fen has not been negatively impacted upon. Compacted areas of the pit floor will be broken up using a ripper.	Imperceptible
Stockpiling of topsoil/subsoil/screenings	Roscat stream (River Barrow and River Nore SAC)	Silt-laden runoff from stockpiles. The increased silt content in runoff had potential to degrade local surface water quality.	Stockpiles have become vegetated which has enhanced stability. Establishment of rooting restricts surface erosion. Stockpiled material shall be re-used in the restoration process.	Imperceptible
Storage of hydrocarbons; leakages from machinery; spillages during refuelling	Roscat stream/Aquifer (River Barrow and River Nore SAC)	Runoff/recharge may contain hydrocarbons.	Potentially contaminating substances were stored in designated areas isolated from surface water drains or open waters. Hazardous wastes such as waste oil, chemicals and preservatives were stored in sealed containers. Fuelling, lubrication and storage areas were in a designated area, not within 50 m of drainage ditches or surface waters. All waste containers were stored within a secondary containment system (e.g. a bund for static tanks or a drip tray for mobile stores and drums). Bunds were capable of storing 110% of tank	Imperceptible

Activity	Attribute	Character of potential impact	measure	Residual impact
			<p>capacity, plus a minimum 30 mm rainwater allowance where the bund was uncovered.</p> <p>Refuelling was carried out by a licensed third party using a mobile bunded bowser. Haulage vehicles were refuelled off-site.</p> <p>Refuelling and parking was on the existing hardstanding pad.</p>	

6.6 Efficacy of the proposed Mitigation Measures

Provide evidence of the degree of confidence in the likely success of the remedial measures

Remedial measures were devised in consideration of the following guidelines:

- Water Framework Directive (2000/60/EC)
- UK Pollution Prevention Guidelines (PPG):
 - PPG1: Good Environmental Practices (2013)
 - PPG2: Above ground oil storage tanks (2011)
 - PPG3: Use and design of oil separators in surface water drainage systems (2006)
 - PPG4: The disposal of sewage where no foul sewer is available (2006)
 - PPG5: Works and maintenance in or near water (2007)
 - PPG6: Working at construction and demolition sites (2012)
 - PPG7: The safe operation of refuelling facilities (2011)
 - PPG8: Safe storage and disposal of used oil (2004)
 - PPG21: Incident response planning (2009)
 - PPG22: Dealing with Spills (2011)
 - PPG26: Drums and Intermediate Bulk Containers (2011)
- CIRIA (2001) "Control of Water Pollution from Construction Sites. Guidance for Consultants and Contractors" (C532);
- CIRIA (2006): Control of Water Pollution from Construction Sites - Guidance for Consultants and Contractors. CIRIA C532. London;
- Environmental Protection Agency (2006). "Environmental Management Guidelines: Environmental Management in the Extractive Industry (Non-Scheduled Minerals)";
- Inland Fisheries Ireland (2016). "Guidelines on Protection of Fisheries During Construction Works In and Adjacent to Waters";
- National Parks and Wildlife Service (Notice Nature) (no year). "Wildlife, Habitats and the Extractive Industry: Guidelines for the Protection of Biodiversity within the Extractive Industry";
- National Roads Authority (2005): Guidelines on Procedures for Assessment and Treatment of Geology, Hydrology and Hydrogeology for National Road Schemes.

Accordingly, given the provenance of the measures as set out above, those measures (when correctly implemented) were successful in ensuring that the European site was preserved at a

favourable conservation status by ensuring the lasting preservation of the constitutive characteristics of this site.

In circumstances where the remedial measures were developed in the light of the best scientific knowledge, no reasonable scientific doubt remains to the *absence* of any adverse effects caused by the existing development on the integrity of the European Site under consideration in circumstances where those remedial measures were implemented.

6.7 Conclusion

In summary, it can be concluded that in light of the conservation objectives and rationale for designation of the River Barrow and River Nore SAC, the potential for significant indirect impacts existed as a result of aspects of the historical excavation operation at Roscat Sand and Gravel Pit, Tullow, Co. Carlow. These potentially significant impacts have been evaluated and with the implementation of the remedial measures, it is concluded that the Substitute Consent development did not result in any impacts that adversely affected the River Barrow and River Nore SAC, having regard to the sites' respective conservation objectives, in circumstances where "no reasonable scientific doubt" remains as to the absence of such adverse effects.

7 REFERENCES

Department of Arts, Heritage and the Gaeltacht (2016) Site Synopsis: River Barrow and River Nore SAC [002162]

Available at:

<https://www.npws.ie/sites/default/files/protected-sites/synopsis/SY0002162.pdf> [Accessed on 04.01.19]

Environment and Heritage Service (2002) The Habitats Regulations: A Guide for Competent Authorities. Environment and Heritage Service, Belfast.

European Commission (2018) Managing Natura 2000 Sites: The Provisions of Article 6 of the Habitats Directive 92/43/EEC.

Fossitt J.A (2000) A Guide to Habitats in Ireland. The Heritage Council, Dublin.

NPWS (2011) Conservation Objectives: River Barrow and River Nore SAC 002162. Version 1. National Parks and Wildlife Service, Department of Arts, Heritage, Regional, Rural and Gaeltacht Affairs.

Available at:

https://www.npws.ie/sites/default/files/protectedsites/conservation_objectives/CO0002162.pdf [Accessed on 04.01.19]

Office for Official Publications of the European Communities (2000) Managing Natura 2000 Sites: The provisions of Article 6 of the 'Habitats' Directive 92/43/EEC. Office for Official Publications of the European Communities, Luxembourg

Office for Official Publications of the European Communities (2001) Assessment of Plans and Projects Significantly Affecting Natura 2000 Sites: Methodological guidance on the provisions of Article 6(3) and (4) of the Habitats Directive 92/43/EEC. Office for Official Publications of the European Communities.

APPENDIX I – NATURA 2000 SITE SYNOPSES AND CONSERVATION OBJECTIVES

Site Name: River Barrow and River Nore SAC

Site Code: 002162

This site consists of the freshwater stretches of the Barrow and Nore River catchments as far upstream as the Slieve Bloom Mountains, and it also includes the tidal elements and estuary as far downstream as Creadun Head in Waterford. The site passes through eight counties – Offaly, Kildare, Laois, Carlow, Kilkenny, Tipperary, Wexford and Waterford. Major towns along the edge of the site include Mountmellick, Portarlinton, Monasterevin, Stradbally, Athy, Carlow, Leighlinbridge, Graiguenamanagh, New Ross, Inistioge, Thomastown, Callan, Bennettsbridge, Kilkenny and Durrow. The larger of the many tributaries include the Lerr, Fushoge, Mountain, Aughavaud, Owenass, Boherbaun and Stradbally Rivers of the Barrow, and the Delour, Dinin, Erkina, Owveg, Munster, Arrigle and King’s Rivers on the Nore.

Both rivers rise in the Old Red Sandstone of the Slieve Bloom Mountains before passing through a band of Carboniferous shales and sandstones. The Nore, for a large part of its course, traverses limestone plains and then Old Red Sandstone for a short stretch below Thomastown. Before joining the Barrow it runs over intrusive rocks poor in silica. The upper reaches of the Barrow also run through limestone. The middle reaches and many of the eastern tributaries, sourced in the Blackstairs Mountains, run through Leinster Granite. The southern end, like the Nore runs over intrusive rocks poor in silica. Waterford Harbour is a deep valley excavated by glacial floodwaters when the sea level was lower than today. The coast shelves quite rapidly along much of the shore.

The site is a Special Area of Conservation (SAC) selected for the following habitats and/or species listed on Annex I / II of the E.U. Habitats Directive (* = priority; numbers in brackets are Natura 2000 codes):

- [1130] Estuaries
- [1140] Tidal Mudflats and Sandflats
- [1170] Reefs
- [1310] *Salicornia* Mud
- [1330] Atlantic Salt Meadows
- [1410] Mediterranean Salt Meadows
- [3260] Floating River Vegetation
- [4030] Dry Heath
- [6430] Hydrophilous Tall Herb Communities
- [7220] Petrifying Springs*
- [91A0] Old Oak Woodlands

[91E0] Alluvial Forests*

- [1016] Desmoulin's Whorl Snail (*Vertigo moulinsiana*)
- [1029] Freshwater Pearl Mussel (*Margaritifera margaritifera*)
- [1092] White-clawed Crayfish (*Austropotamobius pallipes*)
- [1095] Sea Lamprey (*Petromyzon marinus*)
- [1096] Brook Lamprey (*Lampetra planeri*)
- [1099] River Lamprey (*Lampetra fluviatilis*)
- [1103] Twaite Shad (*Alosa fallax*)
- [1106] Atlantic Salmon (*Salmo salar*)
- [1355] Otter (*Lutra lutra*)
- [1421] Killarney Fern (*Trichomanes speciosum*)
- [1990] Nore Freshwater Pearl Mussel (*Margaritifera durrovensis*)

Good examples of alluvial forest (a priority habitat on Annex I of the E.U. Habitats Directive) are seen at Rathsnagadan, Murphy's of the River, in Abbeyleix estate and along other shorter stretches of both the tidal and freshwater elements of the site. Typical species seen include Almond Willow (*Salix triandra*), White Willow (*S. alba*), Rusty Willow (*S. cinerea* subsp. *oleifolia*), Crack Willow (*S. fragilis*) and Osier (*S. viminalis*), along with Iris (*Iris pseudacorus*), Hemlock Water-dropwort (*Oenanthe crocata*), Wild Angelica (*Angelica sylvestris*), Thin-spiked Wood-sedge (*Carex strigosa*), Pendulous Sedge (*C. pendula*), Meadowsweet (*Filipendula ulmaria*), Common Valerian (*Valeriana officinalis*) and the Red Data Book species Nettle-leaved Bellflower (*Campanula trachelium*).

A good example of petrifying springs with tufa formations occurs at Dysart Wood along the Nore. This is a rare habitat in Ireland and one listed with priority status on Annex I of the E.U. Habitats Directive. These hard water springs are characterised by lime encrustations, often associated with small waterfalls. A rich bryophyte flora is typical of the habitat and two diagnostic species, *Palustriella commutata* and *Eucladium verticillatum*, have been recorded.

The best examples of old oak woodlands are seen in the ancient Park Hill woodland in the estate at Abbeyleix; at Kyleadohir, on the Delour, Forest Wood House, Kylecorragh and Brownstown Woods on the Nore; and at Cloghristic Wood, Drummond Wood and Borris Demesne on the Barrow, though other patches occur throughout the site. Abbeyleix Woods is a large tract of mixed deciduous woodland which is one of the only remaining true ancient woodlands in Ireland. Historical records show that Park Hill has been continuously wooded since the 16th century and has the most complete written record of any woodland in the country. It supports a variety of woodland habitats and an exceptional diversity of species including 22 native trees, 44 bryophytes and 92 lichens. It also contains eight indicator species of ancient woodlands. Park Hill is also the site of two rare plants, Nettle-leaved

Bellflower and the moss *Leucodon sciuroides*. The rare Myxomycete fungus, *Licea minima* has been recorded from woodland at Abbeyleix.

Oak woodland covers parts of the valley side south of Woodstock and is well developed at Brownsford where the Nore takes several sharp bends. The steep valley side is covered by oak (*Quercus* spp.), Holly (*Ilex aquifolium*), Hazel (*Corylus avellana*) and Downy Birch (*Betula pubescens*), with some Beech (*Fagus sylvatica*) and Ash (*Fraxinus excelsior*). All the trees are regenerating through a cover of Bramble (*Rubus fruticosus* agg.), Foxglove (*Digitalis purpurea*), Great Wood-rush (*Luzula sylvatica*) and Broad Buckler-fern (*Dryopteris dilatata*).

On the steeply sloping banks of the River Nore, about 5 km west of New Ross, in Co. Kilkenny, Kylecorragh Woods form a prominent feature in the landscape. This is an excellent example of relatively undisturbed, relict oak woodland with a very good tree canopy. The wood is quite damp and there is a rich and varied ground flora. At Brownstown, a small, mature oak dominated woodland occurs on a steep slope. There is younger woodland to the north and east of it. Regeneration throughout is evident. The understorey is similar to the woods at Brownsford. The ground flora of this woodland is developed on acidic, brown earth type soil and comprises a thick carpet of Bilberry (*Vaccinium myrtillus*), Heather (*Calluna vulgaris*), Hard Fern (*Blechnum spicant*), Common Cow-wheat (*Melampyrum pratense*) and Bracken (*Pteridium aquilinum*).

Borris Demesne contains a very good example of a semi-natural broadleaved woodland in very good condition. There is quite a high degree of natural regeneration of oak and Ash through the woodland. At the northern end of the estate oak species predominate. Drummond Wood, also on the Barrow, consists of three blocks of deciduous woods situated on steep slopes above the river. The deciduous trees are mostly oak species. The woods have a well-established understorey of Holly, and the herb layer is varied, with Bramble abundant. The whitebeam *Sorbus devoniensis* has also been recorded here.

Eutrophic tall herb vegetation occurs in association with the various areas of alluvial forest and elsewhere where the floodplain of the river is intact. Characteristic species of the habitat include Meadowsweet, Purple Loosestrife (*Lythrum salicaria*), Marsh Ragwort (*Senecio aquaticus*), Ground Ivy (*Glechoma hederacea*) and Hedge Bindweed (*Calystegia sepium*). Indian Balsam (*Impatiens glandulifera*), an introduced and invasive species, is abundant in places.

Floating river vegetation is well represented in the Barrow and in the many tributaries of the site. In the Barrow the species found include water-starworts (*Callitriche* spp.), Canadian Pondweed (*Elodea canadensis*), Bulbous Rush (*Juncus bulbosus*), water-milfoils (*Myriophyllum* spp.), the pondweed *Potamogeton x nitens*, Broad-leaved Pondweed (*P. natans*), Fennel Pondweed (*P. pectinatus*), Perfoliated Pondweed (*P. perfoliatus*) and crowfoots (*Ranunculus* spp.). The water quality of the Barrow has improved since the vegetation survey was carried out (EPA, 1996).

Dry heath at the site occurs in pockets along the steep valley sides of the rivers especially in the Barrow Valley and along the Barrow tributaries where they occur in the foothills of the Blackstairs Mountains. The dry heath vegetation along the slopes of the river bank consists of Bracken and Gorse (*Ulex europaeus*) with patches of acidic grassland vegetation. Additional typical species include Heath Bedstraw (*Galium saxatile*), Foxglove, Common Sorrel (*Rumex acetosa*) and Creeping Bent (*Agrostis stolonifera*). On the steep slopes above New Ross the Red Data Book species Greater Broomrape (*Orobanche rapum-genistae*) has been recorded. Where rocky outcrops are shown on the maps Bilberry and Great Wood-rush are present. At Ballyhack a small area of dry heath is interspersed with patches of lowland dry grassland. These support a number of clover species, including the legally protected Clustered Clover (*Trifolium glomeratum*) - a species known from only one other site in Ireland. This grassland community is especially well developed on the west side of the mud-capped walls by the road. On the east of the cliffs a group of rock-dwelling species occur, i.e. English Stonecrop (*Sedum anglicum*), Sheep's-bit (*Jasione montana*) and Wild Madder (*Rubia peregrina*). These rocks also support good lichen and moss assemblages with *Ramalina subfarinacea* and *Hedwigia ciliata*.

Dry heath at the site generally grades into wet woodland or wet swamp vegetation lower down the slopes on the river bank. Close to the Blackstairs Mountains, in the foothills associated with the Aughnabriskey, Aughavaud and Mountain Rivers there are small patches of wet heath dominated by Purple Moor-grass (*Molinia caerulea*) with Heather, Tormentil (*Potentilla erecta*), Carnation Sedge (*Carex panicea*) and Bell Heather (*Erica cinerea*).

Salt meadows occur at the southern section of the site in old meadows where the embankment has been breached, along the tidal stretches of in-flowing rivers below Stokestown House, in a narrow band on the channel side of Common Reed (*Phragmites australis*) beds and in narrow fragmented strips along the open shoreline. In the larger areas of salt meadow, notably at Carrickloney, Ballinlaw Ferry and Rochestown on the west bank; Fisherstown, Alderton and Great Island to Dunbrody on the east bank, the Atlantic and Mediterranean sub types are generally intermixed. At the upper edge of the salt meadow in the narrow ecotonal areas bordering the grasslands where there is significant percolation of salt water, the legally protected species Borrer's Saltmarsh-grass (*Puccinellia fasciculata*) and Meadow Barley (*Hordeum secalinum*) are found. The very rare and also legally protected Divided Sedge (*Carex divisa*) is also found. Sea Rush (*Juncus maritimus*) is also present. Other plants recorded and associated with salt meadows include Sea Aster (*Aster tripolium*), Thrift (*Armeria maritima*), Sea Couch (*Elymus pycnanthus*), Spear-leaved Orache (*Atriplex prostrata*), Lesser Sea-spurrey (*Spergularia marina*), Sea Arrowgrass (*Triglochin maritima*) and Sea Plantain (*Plantago maritima*).

Glassworts (*Salicornia* spp.) and other annuals colonising mud and sand are found in the creeks of the saltmarshes and at the seaward edges of them. The habitat also occurs in small amounts on some stretches of the shore free of stones.

The estuary and the other E.U. Habitats Directive Annex I habitats within it form a large component of the site. Extensive areas of intertidal flats, comprised of substrates ranging from fine, silty mud to coarse sand with pebbles/stones are present. Good quality intertidal sand and mudflats have developed on a linear shelf on the western side of Waterford Harbour, extending for over 6 km from north to south between Passage East and Creadaun Head, and in places are over 1 km wide. The sediments are mostly firm sands, though grade into muddy sands towards the upper shore. They have a typical macro-invertebrate fauna, characterised by polychaetes and bivalves. Common species include *Arenicola marina*, *Nephtys hombergii*, *Scoloplos armiger*, *Lanice conchilega* and *Cerastoderma edule*. An extensive area of honey-comb worm biogenic reef occurs adjacent to Duncannon, Co. Wexford on the eastern shore of the estuary. It is formed by the polychaete worm *Sabellaria alveolata*. This intertidal *Sabellaria alveolata* reef is formed as a sheet of interlocking tubes over a considerable area of exposed bedrock. This polychaete species constructs tubes, composed of aggregated sand grains, in tightly packed masses with a distinctive honeycomb-like appearance. These can be up to 25cm proud of the substrate and form hummocks, sheets or more massive formations. A range of species are reported from these reefs including: *Enteromorpha* sp.; *Ulva* sp.; *Fucus vesiculosus*; *Fucus serratus*; *Polysiphonia* sp.; *Chondrus crispus*; *Palmaria palmate*; *Coralinus officinalis*; *Nemertea* sp.; *Actinia equine*; *Patella vulgate*; *Littorina littorea*; *Littorina obtusata* and *Mytilus edulis*.

The western shore of the harbour is generally stony and backed by low cliffs of glacial drift. At Woodstown there is a sandy beach, now much influenced by recreation pressure and erosion. Behind it a lagoonal marsh has been impounded which runs westwards from Gaultiere Lodge along the course of a slow stream. An extensive reedbed occurs here. At the edges is a tall fen dominated by sedges (*Carex* spp.), Meadowsweet, willowherbs (*Epilobium* spp.) and rushes (*Juncus* spp.). Wet woodland also occurs.

The dunes which fringe the strand at Duncannon are dominated by Marram (*Ammophila arenaria*) towards the sea. Other species present include Wild Clary/Sage (*Salvia verbenaca*), a rare Red Data Book species. The rocks around Duncannon ford have a rich flora of seaweeds typical of a moderately exposed shore and the cliffs themselves support a number of coastal species on ledges, including Thrift, Rock Samphire (*Crithmum maritimum*) and Buck's-horn Plantain (*Plantago coronopus*).

Other habitats which occur throughout the site include wet grassland, marsh, reedswamp, improved grassland, arable land, quarries, coniferous plantations, deciduous woodland, scrub and ponds.

Seventeen Red Data Book plant species have been recorded within the site, most in the recent past. These are Killarney Fern (*Trichomanes speciosum*), Divided Sedge, Clustered Clover, Basil Thyme (*Acinos arvensis*), Red Hemp-nettle (*Galeopsis angustifolia*), Borrer's Saltmarsh-grass, Meadow Barley, Opposite-leaved Pondweed (*Groenlandia densa*), Meadow Saffron/Autumn Crocus (*Colchicum autumnale*), Wild Clary/Sage, Nettle-leaved Bellflower, Saw-wort (*Serratula tinctoria*), Bird Cherry

(*Prunus padus*), Blue Fleabane (*Erigeron acer*), Fly Orchid (*Ophrys insectifera*), Ivy Broomrape (*Orobanche hederæ*) and Greater Broomrape. Of these, the first nine are protected under the Flora (Protection) Order, 2015. Divided Sedge was thought to be extinct but has been found in a few locations in the site since 1990. In addition plants which do not have a very wide distribution in the country are found in the site including Thin-spiked Wood-sedge, Field Garlic (*Allium oleraceum*) and Summer Snowflake. Six rare lichens, indicators of ancient woodland, are found including *Lobaria laetevirens* and *L. pulmonaria*. The rare moss *Leucodon sciuroides* also occurs.

The site is very important for the presence of a number of E.U. Habitats Directive Annex II animal species including Freshwater Pearl Mussel (both *Margaritifera margaritifera* and *M. m. durrovensis*), White-clawed Crayfish, Salmon, Twaite Shad, three lamprey species – Sea Lamprey, Brook Lamprey and River Lamprey, the tiny whorl snail *Vertigo moulinsiana* and Otter. This is the only site in the world for the hard water form of the Freshwater Pearl Mussel, *M. m. durrovensis*, and one of only a handful of spawning grounds in the country for Twaite Shad. The freshwater stretches of the River Nore main channel is a designated salmonid river. The Barrow/Nore is mainly a grilse fishery though spring salmon fishing is good in the vicinity of Thomastown and Inistioge on the Nore. The upper stretches of the Barrow and Nore, particularly the Owenass River, are very important for spawning.

The site supports many other important animal species. Those which are listed in the Irish Red Data Book include Daubenton's Bat, Badger, Irish Hare and Common Frog. The rare Red Data Book fish species Smelt (*Osmerus eperlanus*) occurs in estuarine stretches of the site. In addition to the Freshwater Pearl Mussel, the site also supports two other freshwater mussel species, *Anodonta anatina* and *A. cygnea*.

Three rare invertebrates have been recorded in alluvial woodland at Murphy's of the River. These are: *Neoascia obliqua* (Order Diptera: Syrphidae), *Tetanocera freyi* (Order Diptera: Sciomyzidae) and *Dictya umbrarum* (Order Diptera: Sciomyzidae). The rare invertebrate, *Mitostoma chrysomelas* (Order Arachnida), occurs in the old oak woodland at Abbeyleix and only two other sites in the country. Two flies (Order Diptera) *Chrysogaster virescens* and *Hybomitra muhlfeldi* also occur at this woodland.

The site is of ornithological importance for a number of E.U. Birds Directive Annex I species, including Greenland White-fronted Goose, Whooper Swan, Bewick's Swan, Bar-tailed Godwit, Peregrine and Kingfisher. Nationally important numbers of Golden Plover and Bar-tailed Godwit are found during the winter. Wintering flocks of migratory birds are seen in Shanahoe Marsh and the Curragh and Goul Marsh, both in Co. Laois, and also along the Barrow Estuary in Waterford Harbour. There is also an extensive autumnal roosting site in the reedbeds of the Barrow Estuary used by Swallows before they leave the country. The old oak woodland at Abbeyleix has a typical bird fauna including Jay, Long-eared Owl and Raven. The reedbed at Woodstown supports populations of typical waterbirds including Mallard, Snipe, Sedge Warbler and Water Rail.

Land use at the site consists mainly of agricultural activities – mostly intensive in nature and principally grazing and silage production. Slurry is spread over much of the area. Arable crops are also grown. The spreading of slurry and fertiliser poses a threat to the water quality of the salmonid river and to the populations of E.U. Habitats Directive Annex II animal species within the site. Many of the woodlands along the rivers belong to old estates and support many non-native species. Little active woodland management occurs. Fishing is a main tourist attraction along stretches of the main rivers and their tributaries and there are a number of Angler Associations, some with a number of beats. Fishing stands and styles have been erected in places. Both commercial and leisure fishing takes place on the rivers. There is net fishing in the estuary and a mussel bed also. Other recreational activities such as boating, golfing and walking, particularly along the Barrow towpath, are also popular. There is a golf course on the banks of the Nore at Mount Juliet and GAA pitches on the banks at Inistioge and Thomastown. There are active and disused sand and gravel pits throughout the site. Several industrial developments, which discharge into the river, border the site. New Ross is an important shipping port. Shipping to and from Waterford and Belview ports also passes through the estuary.

The main threats to the site and current damaging activities include high inputs of nutrients into the river system from agricultural run-off and several sewage plants, over-grazing within the woodland areas, and invasion by non-native species, for example Cherry Laurel (*Prunus laurocerasus*) and Rhododendron (*Rhododendron ponticum*). The water quality of the site remains vulnerable. Good quality water is necessary to maintain the populations of the Annex II animal species listed above. Good quality is dependent on controlling fertilisation of the grasslands, particularly along the Nore. It also requires that sewage be properly treated before discharge. Drainage activities in the catchment can lead to flash floods which can damage the many Annex II species present. Capital and maintenance dredging within the lower reaches of the system pose a threat to migrating fish species such as lamprey and shad. Land reclamation also poses a threat to the salt meadows and the populations of legally protected species therein.

Overall, the site is of considerable conservation significance for the occurrence of good examples of habitats and of populations of plant and animal species that are listed on Annexes I and II of the E.U. Habitats Directive. Furthermore it is of high conservation value for the populations of bird species that use it. The occurrence of several Red Data Book plant species including three rare plants in the salt meadows and the population of the hard water form of the Freshwater Pearl Mussel, which is limited to a 10 km stretch of the Nore, add further interest to this site.

National Parks and Wildlife Service

Conservation Objectives

River Barrow and River Nore SAC 002162



*An Roinn
Ealaíon, Oidhreachta agus Gaeltachta*
*Department of
Arts, Heritage and the Gaeltacht*

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

* indicates a priority habitat under the Habitats Directive

002162 River Barrow and River Nore SAC

QI	Description
1016	Desmoulin's whorl snail <i>Vertigo moulinsiana</i>
1029	Freshwater pearl mussel <i>Margaritifera margaritifera</i>
1092	White-clawed crayfish <i>Austropotamobius pallipes</i>
1095	Sea lamprey <i>Petromyzon marinus</i>
1096	Brook lamprey <i>Lampetra planeri</i>
1099	River lamprey <i>Lampetra fluviatilis</i>
1103	Twaite shad <i>Alosa fallax</i>
1106	Atlantic salmon (<i>Salmo salar</i>) (only in fresh water)
1130	Estuaries
1140	Mudflats and sandflats not covered by seawater at low tide
1310	<i>Salicornia</i> and other annuals colonizing mud and sand
1330	Atlantic salt meadows (<i>Glauco-Puccinellietalia maritimae</i>)
1355	Otter <i>Lutra lutra</i>
1410	Mediterranean salt meadows (<i>Juncetalia maritimi</i>)
1421	Killarney fern <i>Trichomanes speciosum</i>
1990	Nore freshwater pearl mussel <i>Margaritifera durrovensis</i>
3260	Water courses of plain to montane levels with the <i>Ranunculion fluitantis</i> and <i>Callitricho-Batrachion</i> vegetation
4030	European dry heaths
6430	Hydrophilous tall herb fringe communities of plains and of the montane to alpine levels
7220	* Petrifying springs with tufa formation (<i>Cratoneurion</i>)
91A0	Old sessile oak woods with <i>Ilex</i> and <i>Blechnum</i> in the British Isles
91E0	* Alluvial forests with <i>Alnus glutinosa</i> and <i>Fraxinus excelsior</i> (<i>Alno-Padion</i> , <i>Alnion incanae</i> , <i>Salicion albae</i>)

Supporting documents, relevant reports & publications (listed by date)

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

-
- Title:** Desmoulin's whorl snail (*Vertigo moulinsiana* - 1016) Conservation Status Assessment Report
Year: 2011
Author: Moorkens, E. ; Killeen, I.
Series: Unpublished Report to NPWS
-
- Title:** River Barrow and River Nore SAC (002162): Conservation objectives supporting document - woodland habitats [Version 1]
Year: 2011
Author: NPWS
Series: Unpublished Report to NPWS
-
- Title:** River Barrow and River Nore SAC (002162): Conservation objectives supporting document - coastal habitats [Version 1]
Year: 2011
Author: NPWS
Series: Unpublished Report to NPWS
-
- Title:** River Barrow and River Nore SAC (002162): Conservation objectives supporting document - marine habitats [Version 1]
Year: 2011
Author: NPWS
Series: Unpublished Report to NPWS
-
- Title:** Second Draft Nore Freshwater Pearl Mussel Sub-basin Management Plan (2009-2015)
Year: 2010
Author: DEHLG
Series: Unpublished Report to NPWS
-
- Title:** Site investigations for *Sabellaria alveolata* (Honey-comb worm) biogenic reefs in Ireland
Year: 2010
Author: NPWS
Series: Unpublished Report to NPWS
-
- Title:** Irish Semi-natural Grasslands Survey. Annual report no. 3: Counties Donegal, Dublin, Kildare & Sligo
Year: 2010
Author: O'Neill, F.H.; Martin, J.R.; Devaney, F.M.; McNutt, K.E.; Perrin, P.M. ; Delaney, A.
Series: Unpublished Report to NPWS
-
- Title:** A provisional inventory of ancient and long-established woodland in Ireland
Year: 2010
Author: Perrin, P.M.; Daly, O.H.
Series: Irish Wildlife Manuals No. 46
-
- Title:** Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland [Version 1.0]
Year: 2010
Author: Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series: Irish Wildlife Manuals No. 48
-

Title:	A technical manual for monitoring white-clawed crayfish <i>Austropotamobius pallipes</i> in Irish lakes
Year:	2010
Author:	Reynolds, J.D.; O'Connor, W.; O'Keeffe, C.; Lynn, D.
Series:	Irish Wildlife Manuals No. 45
Title:	Report of the standing scientific committee to the DCENR. The status of Irish salmon stocks in 2010 and precautionary catch advice for 2011
Year:	2010
Author:	SSC
Series:	Unpublished Report to DCENR
Title:	The European Communities Environmental Objectives (Freshwater Pearl Mussel) Regulations 2009. [S.I. 296 of 2009]
Year:	2009
Author:	Government of Ireland
Series:	Irish Statute Book
Title:	The European Communities Environmental Objectives (Surface Water) Regulations 2009. [S.I. 272 of 2009]
Year:	2009
Author:	Government of Ireland
Series:	Irish Statute Book
Title:	Saltmarsh Monitoring Report 2007-2008
Year:	2009
Author:	McCorry, M.; Ryle, T.
Series:	Unpublished Report to NPWS
Title:	<i>Margaritifera durrovensis</i> Survey of Nore River. June – July 2009. NS 2 project
Year:	2009
Author:	Moorkens, E. A.
Series:	Unpublished Report to NPWS
Title:	Benthic Biotope classification of subtidal sedimentary habitats in the Lower River Suir candidate Special Area of Conservation and the River Nore and River Barrow candidate Special Area of Conservation
Year:	2008
Author:	ARMS
Series:	Unpublished Report to NPWS
Title:	A survey of mudflats and sandflats in Ireland. An intertidal soft sediment survey of Waterford Estuary
Year:	2008
Author:	ASU
Series:	Unpublished Report to NPWS
Title:	Assessment of the Risk of Barriers to Fish Migration in the Nore Catchment, Southern Regional Fisheries Board
Year:	2008
Author:	CFB; Compass Informatics
Series:	Unpublished Report to CFB

Title: Poor water quality constrains the distribution and movements of Twaite shad *Alosa fallax fallax* (Lacepede, 1803) in the watershed of river Scheldt

Year: 2008

Author: Maas, J.; Stevens, M. ; Breine, J.

Series: Hydrobiologia 602, 129 - 143

Title: All Ireland Species Action Plan - Killarney fern

Year: 2008

Author: NPWS ; EHS-NI

Series: Unpublished Report to NPWS & EHS-NI

Title: National Survey of Native Woodlands 2003-2008

Year: 2008

Author: Perrin, P.; Martin, J.; Barron, S.; O'Neill, F.; McNutt, K.; Delaney, A.

Series: Unpublished Report to NPWS

Title: Saltmarsh Monitoring Report 2006

Year: 2007

Author: McCorry, M.

Series: Unpublished Report to NPWS

Title: Supporting documentation for the Habitats Directive Conservation Status Assessment - backing documents, Article 17 forms and supporting maps

Year: 2007

Author: NPWS

Series: Unpublished Report to NPWS

Title: A Survey of Juvenile Lamprey Populations in the Corrib and Suir Catchments

Year: 2007

Author: O'Connor, W.

Series: Irish Wildlife Manuals No. 26

Title: Assessment of fish passage and the ecological impact of migration barriers on the River Nore catchment

Year: 2007

Author: Sullivan, A.

Series: Nore Suir Rivers Trust & OPW

Title: Otter Survey of Ireland 2004/2005

Year: 2006

Author: Bailey, M.; Rochford, J.

Series: Irish Wildlife Manuals No. 23

Title: The status of host fish populations and fish species richness in European freshwater pearl mussel (*Margaritifera margaritifera*) streams

Year: 2006

Author: Geist, J.; Porkka, M.; Kuehn, R.

Series: Aquatic Conservation: Marine and Freshwater Ecosystems 16, 251–266

Title: The distribution of Lamprey in the River Barrow SAC

Year: 2006

Author: King, J.J.

Series: Irish Wildlife Manuals No. 21

- Title:** Otters - ecology, behaviour and conservation
Year: 2006
Author: Kruuk, H.
Series: Oxford University Press
-
- Title:** The ecology and conservation of the gametophyte generation of the Killarney Fern (*Trichomanes speciosum* Willd.) in Ireland
Year: 2005
Author: Kingston, N. ; Hayes, C.
Series: Biology and Environment: Proceedings of the Royal Irish Academy 105B(2): 71-79
-
- Title:** Pilot Project for Monitoring Populations of the Freshwater Pearl Mussel. Baseline survey of the Nore River SAC, Counties Laois and Kilkenny
Year: 2004
Author: Moorkens, E. A.
Series: Unpublished Report to NPWS
-
- Title:** Monitoring the river, sea and brook lamprey, *Lampetra fluviatilis*, *L. planeri* and *Petromyzon marinus*
Year: 2003
Author: Harvey, J.; Cowx, I.
Series: Conserving Natura 2000 Rivers Monitoring Series No. 5, English Nature, Peterborough
-
- Title:** Ecology of Watercourses Characterised by *Ranunculion fluitantis* and *Callitriche-Batrachion* Vegetation
Year: 2003
Author: Hatton-Ellis, T.W.; Grieve, N.
Series: Conserving Natura 2000 Rivers Ecology Series No. 11. English Nature, Peterborough.
-
- Title:** Ecology of the Allis and Twaite shad
Year: 2003
Author: Maitland, P.S.; Hatton-Ellis, T.W.
Series: Conserving Natura 2000 Rivers Ecology Series No. 3. English Nature, Peterborough
-
- Title:** A survey of the white-clawed crayfish, *Austropotamobius pallipes* (Lereboullet) and of water quality in two catchments of Eastern Ireland
Year: 2002
Author: Demers, A.; Reynolds, J. D.
Series: Bulletin Français de la Pêche et de la Pisciculture, 367: 729-740
-
- Title:** Reversing the habitat fragmentation of British woodlands
Year: 2002
Author: Peterken, G.
Series: WWF-UK, London
-
- Title:** A survey of broadleaf woodlands in 3 SACs: Barrow-Nore, River Unshin & Lough Forbes
Year: 2000
Author: Browne, A.; Dunne, F.; Roche, N.
Series: Unpublished Report to NPWS
-
- Title:** Diet of Otters *Lutra lutra* on Inishmore, Aran Islands, west coast of Ireland
Year: 1999
Author: Kingston, S.; O'Connell, M.; Fairley, J.S.
Series: Biol & Environ Proc R Ir Acad B 99B:173-182

-
- Title:** Conservation Management of the White-clawed Crayfish, *Austropotamobius pallipes*
Year: 1998
Author: Reynolds, J.D.
Series: Irish Wildlife Manuals No. 1
-
- Title:** Studies on the biology and ecology of Margaritifera in Ireland
Year: 1996
Author: Moorkens, E.A.
Series: Unpublished PhD thesis, University of Dublin, Trinity College.
-
- Title:** Imminent extinction of the Nore freshwater pearl mussel *Margaritifera durrovensis* Phillips: a species unique to Ireland
Year: 1994
Author: Moorkens, E.A. ; Costello, M.J.
Series: Aquatic Conservation: Marine and Freshwater Ecosystems 4,363-365
-
- Title:** The spatial organization of otters (*Lutra lutra*) in Shetland
Year: 1991
Author: Kruuk, H.; Moorhouse, A.
Series: J. Zool, 224: 41-57
-
- Title:** The vegetation of Irish rivers
Year: 1987
Author: Heuff, H.
Series: Unpublished Report
-
- Title:** Otter survey of Ireland
Year: 1982
Author: Chapman, P.J.; Chapman, L.L.
Series: Unpublished Report to Vincent Wildlife Trust
-

Spatial data sources

Year:	2010
Title:	EPA transitional waterbody data
GIS operations:	Clipped to SAC boundary
Used for:	1130 (map 2)
Year:	Interpolated 2011
Title:	Intertidal and subtidal surveys 2008 & 2010
GIS operations:	Polygon feature classes from marine community types base data sub-divided based on interpolation of marine survey data
Used for:	Marine community types, 1140 (maps 3 & 4)
Year:	2005
Title:	OSi Discovery series vector data
GIS operations:	High water mark (HWM) and low water mark (LWM) polyline feature classes converted into polygon feature classes and combined; Saltmarsh and Sand Dune datasets erased out if applicable
Used for:	Marine community types base data (map 4)
Year:	Revision 2010
Title:	Saltmarsh Monitoring Project 2007-2008. Version 1
GIS operations:	QIs selected; clipped to SAC boundary; overlapping regions with Sand Dune data investigated and resolved with expert opinion used
Used for:	1310, 1330, 1410 (map 5)
Year:	Derived 2011
Title:	Internal NPWS files
GIS operations:	Dataset created from spatial reference contained in files
Used for:	7220 (map 6)
Year:	Revision 2010
Title:	National Survey of Native Woodlands 2003-2008. Version 1
GIS operations:	QIs selected; clipped to SAC boundary
Used for:	91A0, 91E0 (map 6)
Year:	2011
Title:	NPWS rare and threatened species database
GIS operations:	Dataset created from spatial references in database records
Used for:	1016, 1092, 1421, 1990 (map 7)
Year:	2005
Title:	OSi Discovery series vector data
GIS operations:	Creation of an 80m buffer on the marine side of the high water mark (HWM); creation of a 10m buffer on the terrestrial side of the HWM; combination of 80m and 10m HWM buffer datasets; creation of a 10m buffer on the landward side of the river banks data; creation of a 20m buffer applied to river centerline and stream data; combination of 10m river banks and 20m river and stream centerline buffer datasets; combined river and stream buffer dataset clipped to HWM; combination of HWM buffer dataset with river and stream buffer dataset; overlapping regions investigated and resolved; resulting dataset clipped to SAC boundary
Used for:	1355 (no map)

1016 Desmoulin's whorl snail *Vertigo moulinsiana*

To maintain the favourable conservation condition of Desmoulin's whorl snail in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution: occupied sites	Number	No decline. Two known sites: Borris Bridge, Co. Carlow S711503; Boston Bridge, Kilnaseer S338774, Co. Laois. See map 7	Data from NPWS rare and threatened species database
Population size: adults	Number per positive sample	At least 5 adults snails in at least 50% of samples	Attribute and target from Moorkens and Killeen (2011)
Population density	Percentage positive samples	Adult snails present in at least 60% of samples per site	Attribute and target from Moorkens and Killeen (2011)
Area of occupancy	Hectares	Minimum of 1ha of suitable habitat per site	Attribute and target from Moorkens and Killeen (2011)
Habitat quality: vegetation	Percentage of samples with suitable vegetation	90% of samples in habitat classes I and II as defined in Moorkens & Killeen (2011)	Attribute and target from Moorkens and Killeen (2011)
Habitat quality: soil moisture levels	Percentage of samples with appropriate soil moisture levels	90% of samples in moisture class 3-4 as defined in Moorkens & Killeen (2011)	Attribute and target from Moorkens and Killeen (2011)

1029 Freshwater pearl mussel *Margaritifera margaritifera*

The status of the freshwater pearl mussel (*Margaritifera margaritifera*) as a qualifying Annex II species for the River Barrow and River Nore SAC is currently under review. The outcome of this review will determine whether a site-specific conservation objective is set for this species. Please note that the Nore freshwater pearl mussel (*Margaritifera durrovensis*) remains a qualifying species for this SAC. This document contains a conservation objective for the latter species.

1092 White-clawed crayfish *Austropotamobius pallipes*

To maintain the favourable conservation condition of White-clawed crayfish in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Occurrence	No reduction from baseline. See map 7	The crayfish is present almost throughout this SAC. The records extend as far downstream as Thomastown on the Nore and Graiguenamanagh on the Barrow
Population structure: recruitment	Percentage occurrence of juveniles and females with eggs	Juveniles and/or females with eggs in at least 50% of positive samples	See Reynolds et al. (2010) for further details
Negative indicator species	Occurrence	No alien crayfish species	Alien crayfish species are identified as major direct threat to this species and as disease vector. See Reynolds (1998) for further details
Disease	Occurrence	No instances of disease	Disease is identified as major threat and has occurred in Ireland even in the absence of alien vectors. See Reynolds (1998) for further details
Water quality	EPA Q value	At least Q3-4 at all sites sampled by EPA	Target taken from Demers and Reynolds (2002). Q values based on triennial water quality surveys carried out by the Environmental Protection Agency (EPA)
Habitat quality: heterogeneity	Occurrence of positive habitat features	No decline in heterogeneity or habitat quality	Crayfish need high habitat heterogeneity. Larger crayfish must have stones to hide under, or an earthen bank in which to burrow. Hatchlings shelter in vegetation, gravel and among fine tree-roots. Smaller crayfish are typically found among weed and debris in shallow water. Larger juveniles in particular may also be found among cobbles and detritus such as leaf litter. These conditions must be available on the whole length of occupied habitat

1095 Sea lamprey *Petromyzon marinus*

To restore the favourable conservation condition of Sea lamprey in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution: extent of anadromy	% of river accessible	Greater than 75% of main stem length of rivers accessible from estuary	Artificial barriers can block or cause difficulties to lampreys' upstream migration, thereby limiting species to lower stretches and restricting access to spawning areas. See King (2006), Sullivan (2007) and CFB and Compass Informatics (2008) for further information on artificial barriers
Population structure of juveniles	Number of age/size groups	At least three age/size groups present	Attribute and target based on data from Harvey and Cowx (2003) and O'Connor, (2007). King (2007) provides survey information for the Barrow
Juvenile density in fine sediment	Juveniles/m ²	Juvenile density at least 1/m ²	Juveniles burrow in areas of fine sediment in still water. Attribute and target based on data from Harvey and Cowx (2003)
Extent and distribution of spawning habitat	m ² and occurrence	No decline in extent and distribution of spawning beds	Attribute and target based on spawning bed mapping by Inland Fisheries Ireland (IFI). Lampreys spawn in clean gravels. Artificial barriers are currently preventing lamprey from accessing suitable spawning habitat. See King (2006), Sullivan (2007) and CFB and Compass Informatics (2008) for further information
Availability of juvenile habitat	Number of positive sites in 3rd order channels (and greater), downstream of spawning areas	More than 50% of sample sites positive	Artificial barriers are currently preventing juvenile lampreys from accessing the full extent of suitable habitat. See King (2006), Sullivan (2007) and CFB and Compass Informatics (2008) for further information

1096 Brook lamprey *Lampetra planeri*

To restore the favourable conservation condition of Brook lamprey in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	% of river accessible	Access to all watercourses down to first order streams	Artificial barriers can block lampreys' upstream migration, thereby limiting species to lower stretches and restricting access to spawning areas. See King (2006), Sullivan (2007) and CFB and Compass Informatics (2008) for further information on artificial barriers
Population structure of juveniles	Number of age/size groups	At least three age/size groups of brook/river lamprey present	Attribute and target based on data from Harvey and Cowx (2003). King (2007) provides survey information for the Barrow. It is impossible to distinguish between brook and river lamprey juveniles in the field, hence they are considered together in this target
Juvenile density in fine sediment	Juveniles/m ²	Mean catchment juvenile density of brook/river lamprey at least 2/m ²	Juveniles burrow in areas of fine sediment in still water. Attribute and target based on data from Harvey and Cowx (2003) who state 10/m ² in optimal conditions and more than 2/m ² on a catchment basis
Extent and distribution of spawning habitat	m ² and occurrence	No decline in extent and distribution of spawning beds	Attribute and target based on spawning bed mapping by Inland Fisheries Ireland (IFI). Lampreys spawn in clean gravels. Artificial barriers are currently preventing lamprey from accessing suitable spawning habitat. See King (2006), Sullivan (2007) and CFB and Compass Informatics (2008) for further information
Availability of juvenile habitat	Number of positive sites in 2nd order channels (and greater), downstream of spawning areas	More than 50% of sample sites positive	Artificial barriers are currently preventing juvenile lampreys from accessing the full extent of suitable habitat. See King (2006), Sullivan (2007) and CFB and Compass Informatics (2008) for further information

1099 River lamprey *Lampetra fluviatilis*

To restore the favourable conservation condition of River lamprey in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution: extent of anadromy	% of river accessible	Greater than 75% of main stem and major tributaries down to second order accessible from estuary	Artificial barriers can block lampreys' upstream migration, thereby limiting species to lower stretches and restricting access to spawning areas. See King (2006), Sullivan (2007) and CFB and Compass Informatics (2008) for further information on artificial barriers
Population structure of juveniles	Number of age/size groups	At least three age/size groups of river/brook lamprey present	Attribute and target based on data from Harvey and Cowx (2003). King (2007) provides survey information for the Barrow. It is impossible to distinguish between brook and river lamprey juveniles in the field, hence they are considered together in this target
Juvenile density in fine sediment	Juveniles/m ²	Mean catchment juvenile density of brook/river lamprey at least 2/m ²	Juveniles burrow in areas of fine sediment in still water. Attribute and target based on data from Harvey and Cowx (2003) who state 10/m ² in optimal conditions and more than 2/m ² on a catchment basis
Extent and distribution of spawning habitat	m ² and occurrence	No decline in extent and distribution of spawning beds	Attribute and target based on spawning bed mapping by Inland Fisheries Ireland (IFI). Lampreys spawn in clean gravels. Artificial barriers are currently preventing lamprey from accessing suitable spawning habitat. See King (2006), Sullivan (2007) and CFB and Compass Informatics (2008) for further information
Availability of juvenile habitat	Number of positive sites in 2nd order channels (and greater), downstream of spawning areas	More than 50% of sample sites positive	Artificial barriers are currently preventing juvenile lampreys from accessing the full extent of suitable habitat. See King (2006), Sullivan (2007) and CFB and Compass Informatics (2008) for further information

1103 Twaite shad *Alosa fallax*

To restore the favourable conservation condition of Twaite shad in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution: extent of anadromy	% of river accessible	Greater than 75% of main stem length of rivers accessible from estuary	In some catchments, artificial barriers block twaite shads' upstream migration, thereby limiting species to lower stretches and restricting access to spawning areas
Population structure: age classes	Number of age classes	More than one age class present	Regular breeding has been confirmed in the River Barrow in recent years, but not in the Nore
Extent and distribution of spawning habitat	m ² and occurrence	No decline in extent and distribution of spawning habitats	
Water quality: oxygen levels	Milligrammes per litre	No lower than 5mg/l	Attribute and target based on Maas, Stevens and Briene (2008)
Spawning habitat quality: Filamentous algae; macrophytes; sediment	Occurrence	Maintain stable gravel substrate with very little fine material, free of filamentous algal (macroalgae) growth and macrophyte (rooted higher plants) growth	See Maitland and Hatton-Ellis (2003) for further information

Conservation objectives for: River Barrow and River Nore SAC [002162]

1106 Atlantic salmon (*Salmo salar*) (only in fresh water)

To restore the favourable conservation condition of Salmon in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution: extent of anadromy	% of river accessible	100% of river channels down to second order accessible from estuary	Artificial barriers block salmon's upstream migration, thereby limiting species to lower stretches and restricting access to spawning areas. See Sullivan (2007) and CFB and Compass Informatics (2008) for further information on artificial barriers
Adult spawning fish	Number	Conservation Limit (CL) for each system consistently exceeded	A conservation limit is defined by the North Atlantic Salmon Conservation Organisation (NASCO) as "the spawning stock level that produces long-term average maximum sustainable yield as derived from the adult to adult stock and recruitment relationship". The target is based on the Standing Scientific Committee of the National Salmon Commission's annual model output of CL attainment levels. See SSC (2010). Stock estimates are either derived from direct counts of adults (rod catch, fish counter) or indirectly by fry abundance counts. The Nore is currently exceeding its CL, while the Barrow is below its CL
Salmon fry abundance	Number of fry/5 minutes electrofishing	Maintain or exceed 0+ fry mean catchment-wide abundance threshold value. Currently set at 17 salmon fry/5 min sampling	Target is threshold value for rivers currently exceeding their conservation limit (CL)
Out-migrating smolt abundance	Number	No significant decline	Smolt abundance can be negatively affected by a number of impacts such as estuarine pollution, predation and sea lice (<i>Lepeophtheirus salmonis</i>)
Number and distribution of redds	Number and occurrence	No decline in number and distribution of spawning redds due to anthropogenic causes	Salmon spawn in clean gravels. Artificial barriers are currently preventing salmon from accessing suitable spawning habitat
Water quality	EPA Q value	At least Q4 at all sites sampled by EPA	Q values based on triennial water quality surveys carried out by the Environmental Protection Agency (EPA)

1130 Estuaries

To maintain the favourable conservation condition of Estuaries in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	The permanent habitat area is stable or increasing, subject to natural processes. See map 2	Habitat area was estimated using OSI data and the defined Transitional Water Body area under the Water Framework Directive as 3856ha. See marine supporting document for further details
Community distribution	Hectares	The following sediment communities should be maintained in a natural condition: Muddy estuarine community complex; Sand to muddy fine sand community complex; Fine sand with <i>Fabulina fabula</i> community. See map 4	The likely area of sediment communities was derived from a combination of intertidal and subtidal surveys undertaken in 2008 (ARMS, 2008; ASU, 2008). See marine supporting document for further details
Community extent	Hectares	Maintain the natural extent of the <i>Sabellaria alveolata</i> reef, subject to natural process. See map 4	The likely area of this community is derived from a survey undertaken in 2010 (NPWS, 2010). See marine supporting document for further details

1140 Mudflats and sandflats not covered by seawater at low tide

To maintain the favourable conservation condition of the Mudflats and sandflats not covered by seawater at low tide in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	The permanent habitat area is stable or increasing, subject to natural processes. See map 3	Habitat area was estimated using OSI data as 926ha. See marine supporting document for further details
Community distribution	Hectares	The following sediment communities should be maintained in a natural condition: Muddy estuarine community complex; Sand to muddy fine sand community complex. See map 4	The likely area of sediment communities was derived from a combination of intertidal and subtidal surveys undertaken in 2008 (ARMS, 2008; ASU, 2008). See marine supporting document for further details

1310 Salicornia and other annuals colonizing mud and sand

To maintain the favourable conservation condition of *Salicornia* and other annuals colonizing mud and sand in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For the one sub-site mapped: Ringville - 0.03ha. See map 5	Based on data from the Saltmarsh Monitoring Project (McCorry and Ryle, 2009). The Ringville sub-site was mapped and no additional areas of potential <i>Salicornia</i> mudflat were identified from an examination of aerial photographs, giving a total estimated area of 0.03ha. NB further unsurveyed areas maybe present within the site. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 5	See coastal habitats supporting document for further details
Physical structure: sediment supply	Presence/absence of physical barriers	Maintain or where necessary restore natural circulation of sediments and organic matter, without any physical obstructions	See coastal habitats supporting document for further details
Physical structure: flooding regime	Hectares flooded; frequency	Maintain natural tidal regime	See coastal habitats supporting document for further details
Physical structure: creeks and pans	Occurrence	Maintain/restore creek and pan structure, subject to natural processes, including erosion and succession	Based on McCorry and Ryle (2009). See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain range of saltmarsh habitat zonations including transitional zones, subject to natural processes including erosion and succession. See map 5	Based on McCorry and Ryle (2009). See coastal habitats supporting document for further details
Vegetation structure: vegetation height	Centimetres	Maintain structural variation within sward	Based on McCorry and Ryle (2009). See coastal habitats supporting document for further details
Vegetation structure: vegetation cover	Percentage cover at a representative sample of monitoring stops	Maintain more than 90% of area outside creeks vegetated.	Based on McCorry and Ryle (2009). See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative sample of monitoring stops	Maintain range of sub-communities with typical species listed in Saltmarsh Monitoring Project (McCorry & Ryle, 2009).	See coastal habitats supporting document for further details
Vegetation structure: negative indicator species: <i>Spartina anglica</i>	Hectares	No significant expansion of <i>Spartina</i> . No new sites for this species and an annual spread of less than 1% where it is already known to occur	Based on McCorry and Ryle (2009). See coastal habitats supporting document for further details

1330 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*)

To restore the favourable conservation condition of Atlantic salt meadows in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For sub-sites mapped: Dunbrody Abbey - 1.25ha, Killowen - 2.59ha, Rochestown - 17.50ha, Ringville - 6.70ha. See map 5	Based on data from the Saltmarsh Monitoring Project (McCorry and Ryle, 2009). Four sub-sites were mapped and additional areas of potential saltmarsh were identified from an examination of aerial photographs, giving a total estimated area of Atlantic salt meadow of 35.07ha. NB further unsurveyed areas maybe present within the site. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 5	See coastal habitats supporting document for further details
Physical structure: sediment supply	Presence/absence of physical barriers	Maintain/restore natural circulation of sediments and organic matter, without any physical obstructions	See coastal habitats supporting document for further details
Physical structure: flooding regime	Hectares flooded; frequency	Maintain natural tidal regime	See coastal habitats supporting document for further details
Physical structure: creeks and pans	Occurrence	Maintain/restore creek and pan structure, subject to natural processes, including erosion and succession	Based on McCorry and Ryle (2009). See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain range of saltmarsh habitat zonations including transitional zones, subject to natural processes including erosion and succession. See map 5	Based on McCorry and Ryle (2009). See coastal habitats supporting document for further details
Vegetation structure: vegetation height	Centimetres	Maintain structural variation within sward	Based on McCorry and Ryle (2009). See coastal habitats supporting document for further details
Vegetation structure: vegetation cover	Percentage cover at a representative sample of monitoring stops	Maintain more than 90% of area outside creeks vegetated	Based on McCorry and Ryle (2009). See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative sample of monitoring stops	Maintain range of sub-communities with typical species listed in Saltmarsh Monitoring Project (McCorry & Ryle, 2009)	See coastal habitats supporting document for further details
Vegetation structure: negative indicator species: <i>Spartina anglica</i>	Hectares	No significant expansion of <i>Spartina</i> . No new sites for this species and an annual spread of less than 1% where it is already known to occur	Based on McCorry and Ryle (2009). See coastal habitats supporting document for further details

1355 Otter *Lutra lutra*

To restore the favourable conservation condition of Otter in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Percentage positive survey sites	No significant decline	Measure based on standard otter survey technique. FCS target, based on 1980/81 survey findings, is 88% in SACs. Current range in south-east estimated at 73% (Bailey and Rochford, 2006)
Extent of terrestrial habitat	Hectares	No significant decline. Area mapped and calculated as 122.8ha above high water mark (HWM); 1136.0ha along river banks / around ponds	No field survey. Areas mapped to include 10m terrestrial buffer along shoreline (above HWM and along river banks) identified as critical for otters (NPWS, 2007)
Extent of marine habitat	Hectares	No significant decline. Area mapped and calculated as 857.7ha	No field survey. Area mapped based on evidence that otters tend to forage within 80m of the shoreline (HWM) (NPWS, 2007; Kruuk, 2006)
Extent of freshwater (river) habitat	Kilometres	No significant decline. Length mapped and calculated as 616.6km	No field survey. River length calculated on the basis that otters will utilise freshwater habitats from estuary to headwaters (Chapman and Chapman, 1982)
Extent of freshwater (lake) habitat	Hectares	No significant decline. Area mapped and calculated as 2.6ha	No field survey. Area mapped based on evidence that otters tend to forage within 80m of the shoreline (NPWS, 2007)
Couching sites and holts	Number	No significant decline	Otters need lying up areas throughout their territory where they are secure from disturbance (Kruuk, 2006; Kruuk and Moorhouse, 1991)
Fish biomass available	Kilograms	No significant decline	Broad diet that varies locally and seasonally, but dominated by fish, in particular salmonids, eels and sticklebacks in freshwater (Bailey and Rochford, 2006) and wrasse and rockling in coastal waters (Kingston et al., 1999)

1410 Mediterranean salt meadows (*Juncetalia maritimi*)

To restore the favourable conservation condition of Mediterranean salt meadows in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For sub-sites mapped: Dunbrody Abbey - 0.08ha, Rochestown - 0.04ha, Ringville - 6.70ha. See map 5	Based on data from the Saltmarsh Monitoring Project (McCorry and Ryle, 2009). Three sub-sites were mapped and no additional areas of potential saltmarsh were identified from an examination of aerial photographs, giving a total estimated area of Mediterranean salt meadow of 6.82ha. NB further unsurveyed areas maybe present within the site. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 5	See coastal habitats supporting document for further details
Physical structure: sediment supply	Presence/absence of physical barriers	Maintain or where necessary restore natural circulation of sediments and organic matter, without any physical obstructions	See coastal habitats supporting document for further details
Physical structure: flooding regime	Hectares flooded; frequency	Maintain natural tidal regime	See coastal habitats supporting document for further details
Physical structure: creeks and pans	Occurrence	Maintain/restore creek and pan structure, subject to natural processes, including erosion and succession	Based on McCorry and Ryle (2009). See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain range of saltmarsh habitat zonations including transitional zones, subject to natural processes including erosion and succession. See map 5	Based on McCorry and Ryle (2009). See coastal habitats supporting document for further details
Vegetation structure: vegetation height	Centimetres	Maintain structural variation within sward	Based on McCorry and Ryle (2009). See coastal habitats supporting document for further details
Vegetation structure: vegetation cover	Percentage cover at a representative sample of monitoring stops	Maintain more than 90% of area outside creeks vegetated.	Based on McCorry and Ryle (2009). See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative sample of monitoring stops	Maintain range of sub-communities with typical species listed in Saltmarsh Monitoring Project (McCorry & Ryle, 2009)	See coastal habitats supporting document for further details
Vegetation structure: negative indicator species: <i>Spartina anglica</i>	Hectares	No significant expansion of <i>Spartina</i> . No new sites for this species and an annual spread of less than 1% where it is already known to occur	Based on McCorry and Ryle (2009). See coastal habitats supporting document for further details

1421 Killarney fern *Trichomanes speciosum*

To maintain the favourable conservation condition of Killarney Fern in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Location	No decline. Three locations known, with three colonies of gametophyte and one sporophyte colony. See map 7	Data from NPWS rare and threatened species database
Population size	Number	Maintain at least three colonies of gametophyte, and at least one sporophyte colony of over 35 fronds	Data from NPWS rare and threatened species database
Population structure: juvenile fronds	Occurrence	At least one of the locations to have a population structure comprising sporophyte, unfurling fronds, 'juvenile' sporophyte and gametophyte generations	'Juvenile' sporophytes, which appear as small entire fronds, are known from this site. However, it is unknown whether they are due to apogamous growth or sexual reproduction. Based on Kingston and Hayes (2005) and Ni Dhuill (pers. Comm.)
Habitat extent	m ²	No loss of suitable habitat, such as shaded rock crevices, caves or gullies in or near to, known colonies. No loss of woodland canopy at or near to known locations	Based on Kingston and Hayes (2005) and Ni Dhuill (pers. Comm.)
Hydrological conditions: visible water	Occurrence	Maintain hydrological conditions at the locations so that all colonies are in dripping or damp seeping habitats, and water is visible at all locations	Based on Kingston and Hayes (2005) and Ni Dhuill (pers. Comm.)
Hydrological conditions: humidity	Number of dessicated fronds	No increase. Presence of dessicated sporophyte fronds or gametophyte mats indicates conditions are unsuitable	Based on Kingston and Hayes (2005) and Ni Dhuill (pers. Comm.)
Light levels: shading	Percentage	No changes due to anthropogenic impacts	Based on Kingston and Hayes (2005) and Ni Dhuill (pers. Comm.)
Invasive species	Occurrence	Absent or under control	NPWS and EHS-NI (2008) provides further details

1990 Nore freshwater pearl mussel *Margaritifera durrovensis*

To restore the favourable conservation condition of the Nore freshwater pearl mussel in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Kilometres	Maintain at 15.5km. See map 7	The population stretches from Poorman's Bridge (S407859) to Lismaine Bridge (S442660), with most of the population found between Poorman's Bridge and the Avonmore Creamery above Ballyragget (S 440 722) (Moorkens, 1996)
Population size: adult mussels	Number	Restore to 5,000 adult mussels	The extant wild population of Nore freshwater pearl mussel is estimated as 300 adult individuals (Moorkens, 2009)
Population structure: recruitment	Percentage per size class	Restore to at least 20% of population no more than 65mm in length; and at least 5% of population no more than 30mm in length	Mussels of no more than 65mm are considered 'young mussels' and may be found buried in the substratum and/or beneath adult mussels. Mussels of no more than 30mm are 'juvenile mussels' and are always buried in the substratum. This species is known not to have reproduced successfully in the River Nore since 1970 (Moorkens and Costello, 1994; Moorkens, 2004; Government of Ireland, 2009 [S.I. 272 of 2009])
Population structure: adult mortality	Percentage	No more than 5% decline from previous number of live adults counted; dead shells less than 1% of the adult population and scattered in distribution	5% is considered the cut-off between the combined errors associated with natural fluctuations and sampling methods and evidence of true population decline. 1% of dead shells is considered to be indicative of natural losses
Habitat extent	Kilometres	Restore suitable habitat in length of river corresponding to distribution target (15.5km; see map 7) and any additional stretches necessary for salmonid spawning	The species habitat is a stretch of large lowland river and is a combination of 1) the area of habitat adult and juvenile mussels can occupy and 2) the area of spawning and nursery habitats the host fish can occupy. Fish nursery habitat typically overlaps with mussel habitat. Fish spawning habitat is generally adjacent mussel habitat, but may lie upstream of the generalised mussel distribution. Only those salmonid spawning areas that could regularly contribute juvenile fish to the areas occupied by adult mussels should be considered. The availability of mussel habitat and fish spawning and nursery habitats are determined by flow and substratum conditions. The habitat for the species is currently unsuitable for the survival of adult mussels or the recruitment of juveniles

Conservation objectives for: River Barrow and River Nore SAC [002162]

1990 Nore freshwater pearl mussel *Margaritifera durrovensis*

To restore the favourable conservation condition of the Nore freshwater pearl mussel in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Water quality: Macroinvertebrates and phytobenthos (diatoms)	Ecological quality ratio (EQR)	Restore water quality-macroinvertebrates: EQR greater than 0.90; phytobenthos: EQR greater than 0.93	These EQRs correspond to high ecological status for these two Water Framework Directive biological quality elements. They represent high water quality with very low nutrient concentrations (oligotrophic conditions). The habitat of the Nore pearl mussel failed both standards during 2009 sampling for the Sub-basin Management Plan (DEHLG, 2010). See also The European Communities Environmental Objectives (Surface Water Objectives) Regulations 2009
Substratum quality: Filamentous algae (macroalgae), macrophytes (rooted higher plants)	Percentage	Restore substratum quality-filamentous algae: absent or trace (<5%); macrophytes: absent or trace (<5%)	High abundance of macroalgae was recorded during 2009 sampling for the Sub-basin Management Plan (DEHLG, 2010). Recruitment of juvenile mussels is being prevented by the poor quality of the river substrate
Substratum quality: sediment	Occurrence	Restore substratum quality-stable cobble and gravel substrate with very little fine material; no artificially elevated levels of fine sediment	The habitat for the species is currently unsuitable for the survival of adult mussels or the recruitment of juveniles owing to sedimentation of the substratum. Significant sedimentation has been recorded during all recent mussel monitoring surveys. Recruitment of juvenile mussels is being prevented by the poor quality of the river substrate
Substratum quality: oxygen availability	Redox potential	Restore to no more than 20% decline from water column to 5cm depth in substrate	Differences in redox potential between the water column and the substrate correlate with differences in oxygen levels. Juvenile mussels require full oxygenation while buried in gravel. In suitable habitat, there should be very little loss of redox potential between the water column and underlying gravels. The redox potential loss in 2009 was 58-64% at 5cm depth (DEHLG, 2010)
Hydrological regime: flow variability	Metres per second	Restore appropriate hydrological regimes	The availability of suitable Nore freshwater pearl mussel habitat is largely determined by flow (catchment geology being the other important factor). In order to restore the habitat for the species, flow variability over the annual cycle must be such that: 1) high flows can wash fine sediments from the substratum, 2) low flows do not exacerbate the deposition of fines and 3) low flows do not cause stress to mussels in terms of exposure, water temperatures, food availability or aspects of the reproductive cycle

1990 Nore freshwater pearl mussel *Margaritifera durrovensis*

To restore the favourable conservation condition of the Nore freshwater pearl mussel in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Host fish	Number	Maintain sufficient juvenile salmonids to host glochidial larvae	Salmonid fish are host to the larval form of freshwater pearl mussels and thus, they are essential to the completion of the life cycle. 0+ and 1+ fish are typically used, both because of the habitat overlaps and the development of immunity with age in the fish. Fish presence is considered sufficient, as higher densities and biomass of fish is indicative of enriched conditions in mussel rivers. Geist et al. (2006) found that higher densities of host fish coincided with eutrophication, poor substrate quality for pearl mussels and a lack of pearl mussel recruitment, while significantly lower densities and biomass of host fish were associated with high numbers of juvenile mussels. Fish movement patterns must be such that 0+ fish in the vicinity of the mussel habitat remain in the mussel habitat until their 1+ summer. As native brown trout appear to be favoured by the Nore freshwater pearl mussel, it is particularly important that these are not out-competed by stocked fish

3260 Water courses of plain to montane levels with the *Ranunculion fluitantis* and *Callitricho-Batrachion* vegetation

To maintain the favourable conservation condition of Water courses of plain to montane levels with the *Ranunculion fluitantis* and *Callitricho-Batrachion* vegetation in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat distribution	Occurrence	No decline, subject to natural processes	The full distribution of this habitat and its sub-types in this site is currently unknown. The basis of the selection of the SAC for the habitat is the presence of an excellent example of the vegetation community (nutrient-rich type) associated with extensive tufa deposits on the river bed in the Kings tributary of the Nore (Heuff, 1987). Other examples of this or other sub-types may be present within the SAC
Habitat area	Kilometres	Area stable or increasing, subject to natural processes	The full extent of this habitat in this site is currently unknown. See above
Hydrological regime: river flow	Metres per second	Maintain appropriate hydrological regimes	Due to regular disturbance (through variations in flow), river macrophytes rarely reach a climax condition but frequently occur as transient communities. A natural (relatively unmodified) flow regime is required for both plant communities and channel geomorphology to be in favourable condition, exhibiting typical dynamics for the river type (Hatton-Ellis and Grieve, 2003). For most of the sub-types of this habitat, high flows are required to maintain the substratum (see below) necessary for the characteristic species. Flow variation is particularly important, with high and flood flows being critical to the hydromorphology
Hydrological regime: groundwater discharge	Metres per second	The groundwater flow to the habitat should be permanent and sufficient to maintain tufa formation	This attribute refers to sub-types with tufa formations. Groundwater discharges to this habitat throughout the year
Substratum composition: particle size range	Millimetres	The substratum should be dominated by large particles and free from fine sediments	The tufaceous sub-types develop on relatively stable substrata such as bedrock, boulders and cobbles, where tufa can deposit and accumulate. Tufa deposition is believed to be biologically mediated, by algae and bryophytes. The substratum must remain free of fine sediments such as clay, silt and fine sand, which would adversely affect the growth of algae and mosses

3260 Water courses of plain to montane levels with the *Ranunculion fluitantis* and *Callitricho-Batrachion* vegetation

To maintain the favourable conservation condition of Water courses of plain to montane levels with the *Ranunculion fluitantis* and *Callitricho-Batrachion* vegetation in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Water chemistry: minerals	Milligrammes per litre	The groundwater and surface water should have sufficient concentrations of minerals to allow deposition and persistence of tufa deposits	The tufaceous sub-types require mineral- (typically calcium-) rich groundwaters to allow deposition of tufa. Surface water must also be sufficiently base-rich to prevent chemical erosion. Alkalinity and/or total hardness data may also be relevant
Water quality: suspended sediment	Milligrammes per litre	The concentration of suspended solids in the water column should be sufficiently low to prevent excessive deposition of fine sediments	See substratum composition above. Turbidity data may also be relevant
Water quality: nutrients	Milligrammes per litre	The concentration of nutrients in the water column should be sufficiently low to prevent changes in species composition or habitat condition	Phosphorus (MRP) is typically the limiting nutrient, however increased nitrogen (NO ₃ ⁻) negatively impacts upon the N-fixing blue-green algal communities that frequently contribute to tufa deposition. Nutrient enrichment of the habitat typically leads to increased filamentous-green-algal biomass, and consequent changes in other algae, bryophyte and macrophyte species composition and abundance. Water quality should reach a minimum of Water Framework Directive good status, in terms of nutrient standards, and macroinvertebrate and phytobenthos quality elements
Vegetation composition: typical species	Occurrence	Typical species of the relevant habitat sub-type should be present and in good condition	The sub-types of this habitat are poorly understood and their typical species have not yet been defined. Typical species and appropriate targets may emerge to be site-specific. The typical species of the tufaceous sub-type in the Kings tributary of the Nore are identified in Heuff (1987). The typical species may include higher plants, bryophytes, macroalgae and microalgae
Floodplain connectivity	Area	The area of active floodplain at and upstream of the habitat should be maintained	River connectivity with the floodplain is essential for the functioning of this habitat. The site of the tufaceous sub-type in the King's River is within an area of floodplain, with further large floodplains upstream. Floodplains regulate fine sediment deposition within the channel. See substratum composition above

4030 European dry heaths

To maintain the favourable conservation condition of European dry heaths in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat distribution	Occurrence	No decline from current habitat distribution, subject to natural processes	Spatial extent currently unmapped but indicated as occurring on the steep, free-draining, river valley sides especially the Barrow and tributaries in the foothills of the Blackstairs Mountains (based on NPWS NHA Survey - 1997/98 Site Notes; Natura 2000 Form Explanatory Notes - May 2006; The above NHA survey was prior to the extensions to the SAC that included river habitat and estuary at Ballyhack which may have incorporated additional dry heath habitat)
Habitat area	Hectares	Area stable or increasing, subject to natural processes. Habitat area is not known but estimated as less than 400ha of the area of the SAC, occurring in dispersed locations	Based on NPWS NHA Survey Site Notes (1997/98); Natura 2000 Form Explanatory Notes - May 2006
Physical structure: free-draining, acid, low nutrient soil; rock outcrops	Occurrence	No significant change in soil nutrient status, subject to natural processes. No increase or decrease in area of natural rock outcrop	Based on NPWS NHA Survey Site Notes - 1997/98; Natura 2000 Form Explanatory Notes - May 2006
Vegetation structure: sub-shrub indicator species	Percentage cover	Cover of characteristic sub-shrub indicator species at least 25%: gorse (<i>Ulex europaeus</i>) and where rocky outcrops occur bilberry (<i>Vaccinium myrtillus</i>) and woodrush (<i>Luzula sylvatica</i>). Some rock outcrops support English stonecrop (<i>Sedum anglicum</i>), sheep's bit (<i>Jasione montana</i>) and wild madder (<i>Rubia peregrina</i>) as well as important moss and lichen assemblages	Dry heath in this SAC occurs on free-draining nutrient poor soils and is often characterised by gorse and open acid grassland areas. A characteristic coastal dry heath of the southeast also occurs. Several rare plants occur including two species listed in the Red Data Book (Curtis and McGough, 1988). The species occurring on the site are listed in NPWS NHA Survey Site Notes - 1997/98. A brief overview of the principal characteristics of the dry heath habitat of this SAC is given in the Natura 2000 Explanatory Notes - May 2006
Vegetation structure: senescent gorse	Percentage cover	Cover of senescent gorse less than 50%	Based on NPWS NHA Survey Site Notes and Natura 2000 Form Explanatory Notes - May 2006 and on a modified version of the dry heath condition assessment methodology of Perrin et al. (2010)
Vegetation structure: browsing	Percentage cover	Long shoots of bilberry with signs of browsing collectively less than 33%	Based on NPWS NHA Survey Site Notes and Natura 2000 Form Explanatory Notes - May 2006 and on a modified version of the dry heath condition assessment methodology of Perrin et al. (2010)

4030 European dry heaths

To maintain the favourable conservation condition of European dry heaths in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Vegetation structure: native trees and shrubs	Percentage cover	Cover of scattered native trees and shrub less than 20%	Based on NPWS NHA Survey Site Notes - 1997/98; Natura 2000 Form Explanatory Notes - May 2006 and on a modified version of the dry heath habitat condition assessment methodology of Perrin et al. (2010). From the NHA survey notes the main threats appear to be reclamation or invasion by scrub woodland
Vegetation composition: positive indicator species	Number	Number of positive indicator species at least 2 e.g. gorse and associated dry heath/ acid grassland flora	Dry heath in this SAC occurs on free-draining nutrient poor soils and is characterised by gorse and acid grassland areas. It corresponds to Annex I sub-type "heaths rich in gorse (<i>Ulex</i>) of the Atlantic margins" (European Commission, 2007). Based on NPWS NHA Survey Site Notes -1997/98; Natura 2000 Form Explanatory Notes - May 2006 and a modified version of the dry heath habitat condition assessment methodology of Perrin et al. (2010)
Vegetation structure: positive indicator species	Percentage cover	Cover of positive indicator species at least 60%. This should include plant species characteristic of dry heath in this SAC including gorse, bilberry and associated acid grassland flora	Dry heath in this SAC is characterised by gorse and acid grassland areas and locally bilberry and woodrush. Based on NPWS NHA Survey Site Notes and Natura 2000 Form Explanatory Notes - May 2006 and a modified version of the dry heath habitat condition assessment methodology of Perrin et al. (2010)
Vegetation composition: bryophyte and non-crustose lichen species	Number	Number of bryophyte or non-crustose lichen species present at least 2	Based on NPWS NHA Survey Site Notes and Natura 2000 Form Explanatory Notes - May 2006 and on a modified version of the dry heath habitat condition assessment methodology of Perrin et al. 2010
Vegetation composition: bracken (<i>Pteridium aquilinum</i>)	Percentage cover	Cover of bracken less than 10% - however see 'Notes'	Based on NPWS NHA Survey Site Notes and Natura 2000 Form Explanatory Notes - May 2006 and on a modified version of the dry heath habitat condition assessment methodology of Perrin et al. (2010). Bracken appears to be quite dense in places and before any management action is considered its rate of spread needs to be established as well as its threat, if any, to other dry heath species and its potential value to important fauna (e.g. Twite)

4030 European dry heaths

To maintain the favourable conservation condition of European dry heaths in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Vegetation structure: weedy negative indicator species	Percentage cover	Cover of agricultural weed species (negative indicator species) less than 1%	Based on NPWS NHA Survey Site Notes and Natura 2000 Form Explanatory Notes - May 2006 and on a modified version of the dry heath habitat condition assessment methodology of Perrin et al. (2010)
Vegetation composition: non-native species	Percentage cover	Cover of non-native species less than 1%.	Based on NPWS NHA Survey Site Notes and Natura 2000 Form Explanatory Notes - May 2006 and on a modified version of the dry heath habitat condition assessment methodology of Perrin et al. (2010)
Vegetation composition: rare/scarce heath species	Location, area and number	No decline in distribution or population sizes of rare, threatened or scarce species, including Greater Broomrape (<i>Orobanche rapum-genistae</i>) and the legally protected clustered clover (<i>Trifolium glomeratum</i>)	Broomrape is dependent on gorse at this site as it is parasitic on gorse roots. It is recorded as occurring on steep slopes above New Ross. A small area of excellent dry coastal heath at Ballyhack is interspersed with patches rock and of dry lowland grassland and has a high species diversity. Notably there is an excellent range of Clover (<i>Trifolium</i>) species including the legally protected clustered clover, a species known only from one other site in Ireland. Also <i>T. ornithopodioides</i> , <i>T. striatum</i> and <i>Torilus nodosa</i> . Based on Natura 2000 Form Explanatory Notes May 2006, Irish Red Data Book (Curtis and Mc Gough, 1988) and on the NPWS database of rare and threatened vascular plants. Other areas of coastal heath may also occur
Vegetation structure: disturbed bare ground	Percentage cover	Cover of disturbed bare ground less than 10% (but if peat soil less than 5%)	Based on NPWS NHA Survey Site Notes and Natura 2000 Form Explanatory Notes - May 2006 and on a modified version of the dry heath habitat condition assessment methodology of Perrin et al. (2010)
Vegetation structure: burning	Occurrence	No signs of burning within sensitive areas	Perrin et al. (2010) defines sensitive areas

6430 Hydrophilous tall herb fringe communities of plains and of the montane to alpine levels

To maintain the favourable conservation condition of Hydrophilous tall herb fringe communities of plains and of the montane to alpine levels in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat distribution	Occurrence	No decline, subject to natural processes	Distribution of this habitat in this site is currently unknown. Considered to occur in association with some riverside woodlands, unmanaged river islands and in narrow bands along the floodplain of slow-flowing stretches of river (Natura 2000 Form Explanatory Notes)
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Extent of this habitat in this site is currently unknown. See above
Hydrological regime: Flooding depth/height of water table	Metres	Maintain appropriate hydrological regimes	This habitat requires winter inundation, which results in deposition of naturally nutrient-rich sediment
Vegetation structure:sward height	Centimetres	30-70% of sward is between 40 and 150cm in height	Bare ground, due to natural inundation processes, may often be present. Attribute and target based on the Irish Semi-natural Grassland Survey (O'Neill et al., 2010)
Vegetation composition: broadleaf herb: grass ratio	Percentage	Broadleaf herb component of vegetation between 40 and 90%	Attribute and target based on O'Neill et al. (2010)
Vegetation composition: typical species	Number	At least 5 positive indicator species present	List of positive indicator species identified by O'Neill et al. (2010)
Vegetation composition: negative indicator species	Occurrence	Negative indicator species, particularly non-native invasive species, absent or under control- NB Indian balsam (<i>Impatiens glandulifera</i>), monkeyflower (<i>Mimulus guttatus</i>), Japanese knotweed (<i>Fallopia japonica</i>) and giant hogweed (<i>Heracleum mantegazzianum</i>)	Species listed as being present in the site (Natura 2000 Form Explanatory Notes)

7220 * Petrifying springs with tufa formation (*Cratoneurion*)

To maintain the favourable conservation condition of Petrifying springs with tufa formation (*Cratoneurion*) in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Square metres	Area stable or increasing, subject to natural processes	Extent of this habitat in this site is currently unknown. An area ("Tens of square metres") has been described at one location (Natura 2000 Form Explanatory Notes; internal NPWS files), see below
Habitat distribution	Occurrence	No decline. See map 6 for recorded location	Full distribution of this habitat in this site is currently unknown. It has been described in woodlands at Dysart, between Thomastown and Inistioge (Natura 2000 Form Explanatory Notes; internal NPWS files). NB further areas are likely to occur within the site
Hydrological regime: height of water table; water flow	Metres; metres per second	Maintain appropriate hydrological regimes	Current hydrological regimes are unknown. Petrifying springs rely on permanent irrigation, usually from upwelling groundwater sources or seepage sources
Water quality	Water chemistry measures	Maintain oligotrophic and calcareous conditions	Water chemistry is currently unknown. Water supply to petrifying springs is characteristically oligotrophic and calcareous
Vegetation composition: typical species	Occurrence	Maintain typical species	The bryophytes <i>Cratoneurion commutatum</i> and <i>Eucladium verticillatum</i> are diagnostic of this habitat. Both are found at the location described above. Natura 2000 Form Explanatory Notes and internal NPWS files also list other typical species

91A0 Old sessile oak woods with *Ilex* and *Blechnum* in the British Isles

To restore the favourable conservation condition of Old oak woodland with *Ilex* and *Blechnum* in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, at least 85.08ha for sub-sites surveyed: see map 6	Minimum area, based on 13 sites surveyed by Perrin et al. (2008) - site codes 14, 20, 49, 73, 125, 508, 509, 510, 514, 515, 518, 519, 521, and other sources. NB further unsurveyed areas maybe present within the site
Habitat distribution	Occurrence	No decline. Surveyed locations shown on map 6	Distribution based on Perrin et al. (2008). NB further unsurveyed areas maybe present within the site
Woodland size	Hectares	Area stable of increasing. Where topographically possible, "large" woods at least 25ha in size and "small" woods at least 3ha in size	The sizes of at least some of the existing woodlands need to be increased in order to reduce habitat fragmentation and benefit those species requiring 'deep' woodland conditions (Peterken, 2002). Topographical and land ownership constraints may restrict expansion
Woodland structure: cover and height	Percentage and metres	Diverse structure with a relatively closed canopy containing mature trees; subcanopy layer with semi-mature trees and shrubs; and well-developed herb layer	Described in Perrin et al. (2008); Browne et al. (2000). See woodland habitats supporting document for further details
Woodland structure: community diversity and extent	Hectares	Maintain diversity and extent of community types	Described in Perrin et al. (2008); Browne et al. (2000). See woodland habitats supporting document for further details
Woodland structure: natural regeneration	Seedling:sapling:pole ratio	Seedlings, saplings and pole age-classes occur in adequate proportions to ensure survival of woodland canopy	Oak regenerates poorly. In suitable sites ash can regenerate in large numbers although few seedlings reach pole size
Woodland structure: dead wood	m ³ per hectare; number per hectare	At least 30m ³ /ha of fallen timber greater than 10cm diameter; 30 snags/ha; both categories should include stems greater than 40cm diameter	Dead wood is a valuable resource and an integral part of a healthy, functioning woodland ecosystem.
Woodland structure: veteran trees	Number per hectare	No decline	Mature and veteran trees are important habitats for bryophytes, lichens, saproxylic organisms and some bird species. Their retention is important to ensure continuity of habitats/niches and propagule sources

91A0 Old sessile oak woods with *Ilex* and *Blechnum* in the British Isles

To restore the favourable conservation condition of Old oak woodland with *Ilex* and *Blechnum* in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Woodland structure: indicators of local distinctiveness	Occurrence	No decline	Includes ancient or long-established woodlands, archaeological and geological features as well as red-listed and other rare or localised species. Perrin and Daly (2010) list sites 14, 20, 73, 125, 508, 509, 510, 514, 515, 518, 521 as potential ancient/long established woodlands
Vegetation composition: native tree cover	Percentage	No decline. Native tree cover not less than 95%	Species reported in Perrin et al. (2008); Browne et al. (2000)
Vegetation composition: typical species	Occurrence	A variety of typical native species present, depending on woodland type, including oak (<i>Quercus petraea</i>) and birch (<i>Betula pubescens</i>)	Species reported in Perrin et al. (2008); Browne et al. (2000)
Vegetation composition: negative indicator species	Occurrence	Negative indicator species, particularly non-native invasive species, absent or under control	The following are the most common invasive species in this woodland type: beech (<i>Fagus sylvatica</i>), rhododendron (<i>Rhododendron ponticum</i>), cherry laurel (<i>Prunus laurocerasus</i>)

Conservation objectives for: River Barrow and River Nore SAC [002162]

91E0 * Alluvial forests with *Alnus glutinosa* and *Fraxinus excelsior* (*Alno-Padion*, *Alnion incanae*, *Salicion albae*)

To restore the favourable conservation condition of Alluvial forests with *Alnus glutinosa* and *Fraxinus excelsior* (*Alno-Padion*, *Alnion incanae*, *Salicion albae*) in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:


Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, at least 181.54ha for sites surveyed: see map 6	Minimum area, based on 16 sites surveyed by Perrin et al. (2008) - site codes 10, 15, 17, 126, 127, 262, 282, 287, 511, 516, 517, 518, 520, 608, 1021; Coillte LIFE project and other sources. NB further unsurveyed areas maybe present within the SAC
Habitat distribution	Occurrence	No decline. Surveyed locations shown on map 6	Distribution based on Perrin et al. (2008). NB further unsurveyed areas maybe present within the site
Woodland size	Hectares	Area stable of increasing. Where topographically possible, "large" woods at least 25ha in size and "small" woods at least 3ha in size	The sizes of at least some of the existing woodlands need to be increased in order to reduce habitat fragmentation and benefit those species requiring 'deep' woodland conditions (Peterken, 2002). Topographical and land ownership constraints may restrict expansion
Woodland structure: cover and height	Percentage and metres	Diverse structure with a relatively closed canopy containing mature trees; subcanopy layer with semi-mature trees and shrubs; and well-developed herb layer	Described in Perrin et al. (2008); Browne et al. (2000). See woodland habitats supporting document for further details
Woodland structure: community diversity and extent	Hectares	Maintain diversity and extent of community types	Described in Perrin et al. (2008); Browne et al. (2000). See woodland habitats supporting document for further details
Woodland structure: natural regeneration	Seedling:sapling:pole ratio	Seedlings, saplings and pole age-classes occur in adequate proportions to ensure survival of woodland canopy	Alder and oak regenerate poorly. Ash often regenerates in large numbers although few seedlings reach pole size
Hydrological regime: Flooding depth/height of water table	Metres	Appropriate hydrological regime necessary for maintenance of alluvial vegetation	Periodic flooding is essential to maintain alluvial woodlands along river flood plains but not for woodland around springs/seepage areas
Woodland structure: dead wood	m ³ per hectare; number per hectare	At least 30m ³ /ha of fallen timber greater than 10cm diameter; 30 snags/ha; both categories should include stems greater than 40cm diameter (greater than 20cm diameter in the case of alder)	Dead wood is a valuable resource and an integral part of a healthy, functioning woodland ecosystem

Conservation objectives for: River Barrow and River Nore SAC [002162]

91E0 * Alluvial forests with *Alnus glutinosa* and *Fraxinus excelsior* (*Alno-Padion*, *Alnion incanae*, *Salicion albae*)

To restore the favourable conservation condition of Alluvial forests with *Alnus glutinosa* and *Fraxinus excelsior* (*Alno-Padion*, *Alnion incanae*, *Salicion albae*) in the River Barrow and River Nore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Woodland structure: veteran trees	Number per hectare	No decline	Mature and veteran trees are important habitats for bryophytes, lichens, saproxylic organisms and some bird species. Their retention is important to ensure continuity of habitats/niches and propagule sources
Woodland structure: indicators of local distinctiveness	Occurrence	No decline	Includes ancient or long-established woodlands, archaeological and geological features as well as red-listed and other rare or localised species. Perrin and Daly (2010) list sites 10, 15, 17, 127, 282, 516, 517, 518, 608 as potential ancient/long established woodlands
Vegetation composition: native tree cover	Percentage	No decline. Native tree cover not less than 95%	Species reported in Perrin et al. (2008); Browne et al. (2000)
Vegetation composition: typical species	Occurrence	A variety of typical native species present, depending on woodland type, including ash (<i>Fraxinus excelsior</i>) alder (<i>Alnus glutinosa</i>), willows (<i>Salix</i> spp) and locally, oak (<i>Quercus robur</i>)	Species reported in Perrin et al. (2008); Browne et al. (2000)
Vegetation composition: negative indicator species	Occurrence	Negative indicator species, particularly non-native invasive species, absent or under control	The following are the most common invasive species in this woodland type: sycamore (<i>Acer pseudoplatanus</i>), beech (<i>Fagus sylvatica</i>), rhododendron (<i>Rhododendron ponticum</i>), cherry laurel (<i>Prunus laurocerasus</i>), dogwood (<i>Cornus sericea</i>), Himalayan honeysuckle (<i>Leycesteria formosa</i>) and Himalayan balsam (<i>Impatiens grandiflora</i>)



Legend

SAC 002162

An Roinn Ealaíon, Oidhreachta agus Gaeltachta
 Department of Arts, Heritage and the Gaeltacht

MAP 1: RIVER BARROW AND RIVER NORE CONSERVATION OBJECTIVES SAC DESIGNATION


Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE: SAC 002162
 CO. CARLOW; version 1.03, CO. KILDARE; version 1.04,
 CO. KILKENNY; version 1.1, CO. LAOIS; version 1.07,
 CO. OFFALY; version 1.01, CO. TIPPERARY; version 1.01,
 CO. WATERFORD; version 1.01, CO. WEXFORD; version 1.01

0 5 10 15 km

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision. Reproduced from Ordnance Survey material by permission of the Government (Permit number EN 0059208).
 Níl sna teorainneacha ar na léarscálanna ach nod garshuíomhach ginearálta. Féadfar athbheithníthe a déanamh ar theorainneacha na gceantar comharthaíthe. Macasamhail d'ábhar na Suirbhéaracha Ordoínáis le chead ón Rialtas (Ceadúnas Uimh. EN 0059208).

Map Version 1
Date: April 2011



Legend

- SAC 002162
- 1130 Estuaries
- OSi Discovery Series County Boundary

An Roinn
Ealaíon, Oidhreacht agus Gaeltachta
Department of
Arts, Heritage and the Gaeltacht

MAP 2:
RIVER BARROW AND RIVER NORE
CONSERVATION OBJECTIVES
ESTUARIES

Map to be read in conjunction with the NPWS Conservation Objectives Document.


SITE CODE: SAC 002162
CO. CARLOW; version 1.03, CO. KILDARE; version 1.04,
CO. KILKENNY; version 1.1, CO. LAOIS; version 1.07,
CO. OFFALY; version 1.01, CO. TIPPERARY; version 1.01,
CO. WATERFORD; version 1.01, CO. WEXFORD; version 1.01

0 1 2 3 4 5 km

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision. Reproduced from Ordnance Survey material by permission of the Government (Permit number EN 0059208).
Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithithe a déanamh ar theorainneacha na gceantar comharthaithe. Macsamhail d'ábhar na Suirbhéarachta Ordonáis le chead ón Rialtas (Ceadúnas Uimh. EN 0059208)


N

Map Version 1
Date: April 2011



Legend

- SAC 002162
- 1140 Mudflats and sandflats not covered by sea water at low tide
- OSi Discovery Series County Boundary





Legend

- SAC 002162
- OSi Discovery Series County Boundary

Community Types

- Fine sand with *Fabulina fabula* community
- Muddy estuarine community complex
- Sand to Muddy fine sand community complex
- Sabellaria alveolata* reef






Legend

- SAC 002162
- OSi Discovery Series County Boundary

Saltmarsh Habitats

- 1310 *Salicornia* and other annuals colonising mud and sand
- 1330 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*)
- 1330 / 1410 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*) / Mediterranean salt meadows (*Juncetalia maritimi*)
- 1410 Mediterranean salt meadows (*Juncetalia maritimi*)
- Potential 1330 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*)




- Legend**
- SAC 002162
 - OSI Discovery Series County Boundary
 - ▲ 7220 *Petrifying springs with tufa formation (Cratoneurion)
- Woodland Habitats**
- 91A0 Old sessile oak woods with *Ilex* and *Blechnum* in the British Isles
 - 91E0 *Alluvial forests with *Alnus glutinosa* and *Fraxinus excelsior* (Alno-padion, Alnion incanae, Salicion albae)
 - 91A0 / 91E0 Old sessile oak woods with *Ilex* and *Blechnum* in the British Isles / *Alluvial forests with *Alnus glutinosa* and *Fraxinus excelsior* (Alno-padion, Alnion incanae, Salicion albae)
 - WD1 (Mixed) broadleaved woodland
 - WN2 / WD1 Oak-ash-hazel woodland / (Mixed) broadleaved woodland
 - WN2 / WN6 Oak-ash-hazel woodland / Wet willow-alder-ash woodland

**MAP 6:
RIVER BARROW AND RIVER NORE
CONSERVATION OBJECTIVES
OLD OAK WOODLANDS, ALLUVIAL
FORESTS & PETRIFYING SPRINGS**

Map to be read in conjunction with the NPWS Conservation Objectives Document.


SITE CODE: SAC 002162
CO. CARLOW; version 1.03, CO. KILDARE; version 1.04,
CO. KILKENNY; version 1.1, CO. LAOIS; version 1.07,
CO. OFFALY; version 1.01, CO. TIPPERARY; version 1.01,
CO. WATERFORD; version 1.01, CO. WEXFORD; version 1.01



The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision. Reproduced from Ordnance Survey material by permission of the Government (Permit number EN 0059208).
Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithnithe a déanamh ar theorainneacha na gceantar conharthaithe. Macasamhail d'ábhar na Suirbhéarachta Ordonáis le chead ón Rialtas (Ceadúnas Uimh. EN 0059208)



**Map Version 1
Date: April 2011**



Legend


- SAC 002162
- OSI Discovery Series County Boundary
- 1016 Desmoulin's Whorl Snail - *Vertigo moulinsiana*
- 1092 White-Clawed Crayfish - *Austropotamobius pallipes*
- 1421 Killarney Fern - *Trichomanes speciosum*
- 1990 Nore Freshwater Pearl Mussel - *Margaritifera durrovensis*



**MAP 7:
RIVER BARROW AND RIVER NORE
CONSERVATION OBJECTIVES
DESMOULIN'S WHORL SNAIL, WHITE-
CLAWED CRAYFISH, NORE FRESHWATER
PEARL MUSSEL & KILLARNEY FERN**

Map to be read in conjunction with the NPWS Conservation Objectives Document.

SITE CODE: SAC 002162
CO. CARLOW; version 1.03, CO. KILDARE; version 1.04,
CO. KILKENNY; version 1.1, CO. LAOIS; version 1.07,
CO. OFFALY; version 1.01, CO. TIPPERARY; version 1.01,
CO. WATERFORD; version 1.01, CO. WEXFORD; version 1.01



The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision. Reproduced from Ordnance Survey material by permission of the Government (Permit number EN 0059208).
Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithniú a déanamh ar theorainneacha na gceantar comharthaithe. Macasamhail d'ábhar na Suirbhéarachta Ordonáis le chead ón Rialtas (Ceadúnas Uimh. EN 0059208)



**Map Version 1
Date: April 2011**



An Roinn
Ealaíon, Oidhreacht agus Gaeltachta

Department of
Arts, Heritage and the Gaeltacht

**Produced by: National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: natureconservation@environ.ie**

Citation:

NPWS (2011) Conservation Objectives: River Barrow and River Nore SAC 002162. Version 1.0. National Parks and Wildlife Service, Department of Arts, Heritage and the Gaeltacht.

Series Editors: Rebecca Jeffrey & Naomi Kingston

ISSN 2009-4086

APPENDIX II – RESPONSE TO SCOPING CONSULTATION

Patrick O'Donnell
Earth Science Partnership
Consulting Engineers
Tonranny
Westport
Co. Mayo

19 November 2018

**EIA Screening & Scoping document for substitute consent & planning
in respect of a quarry at Roscat, Ardristan, Tullow, Co. Carlow.**

Dear Mr. O'Donnell,

The site of this facility is in close proximity to a tributary of the Burren River which is in turn a tributary of the River Barrow. The Barrow River is an important Spring Salmon & sea trout fishery. The Barrow system supports several species listed in Annex II of the Directive including Salmon, River Lamprey, Brook Lamprey, Sea Lamprey, Freshwater Pearl Mussel and Otter. Much of the main channel of the Barrow River is a candidate Special Area for Conservation (SAC) under the European Habitats Directive.

An examination of the old 6 and 25 inch maps for this site highlight that the south-western boundary of the site, borders a very wet area with a number of springs rising from there.

An examination of more recent aerial photos of the site highlight that there was significant production of concrete blocks and other concrete products on-site in the recent past.

The following are our concerns:

1. IFI request clarification if the applicant proposes to use the site for production of concrete blocks and other concrete products. Uncured concrete can kill fish and macro-invertebrates by altering the pH of the water. Given the proximity of the watercourse and the ground conditions on-site, we have concerns that there may be potential for contamination of the nearby watercourse from on-site operations. We would also point out that passage through silt ponds does not represent adequate treatment of waters which have come in contact with uncured concrete and have an pH.
2. One of the potential impacts of projects such as this, is the discharge of silt-laden waters to fisheries streams from sites at which earth moving and excavation works are ongoing. Silt can clog salmonid spawning beds, and juvenile salmonids are particularly sensitive to siltation of gill structures. Similarly plant and macro-invertebrate communities can literally be blanketed over, and this can lead to loss or degradation of valuable habitat.
3. Systems should be put in place to ensure that there shall be no discharge of suspended solids or other deleterious matter to watercourses during any phase of works at this site.

4. It is our understanding that no sand washing is carried out on-site and that there are no future plans to wash sand at this facility.
5. It is our understanding from the information submitted that there is no pumping from the quarry site to the surface waters.
6. It is our understanding that all sand extraction is carried out above groundwater level.
7. The design and sizing of the surface water drainage system must ensure that no suspended solids enter the neighbouring watercourse, even during periods of prolonged heavy rainfall.
8. All surface waters from the site and access roads should be channelled through adequately sized petrol/oil interceptors prior and be subject to attenuation prior to discharge to surface waters.
9. Given the proximity of surface waters draining to the Barrow the pollution threat from concrete and concrete/cement washings is significant. Good housekeeping is of the utmost importance while using concrete or cement, near watercourses.
10. Refuelling of machinery must be carried out in bunded areas.
11. Fuels, oils, greases and hydraulic fluids must be stored in bunded compounds.
12. All waste oil, empty oil containers and other hazardous wastes are disposed of in conjunction with the requirements of the Waste Management Act 1996.
13. The activities proposed for this site are likely to result in significant lorry traffic to and from the site, with potential for the generation of significant suspended solids pollution in the associated road run-off. It is imperative that the potential for suspended solids pollution from road run-off associated with vehicles entering and leaving this site is addressed.
14. We note that the applicant proposes to provide a wheel wash to prevent soils/clays being deposited on the public road. Our experience is that wheel-washes, if not designed/managed properly can represent a significant source of suspended solids pollution to surface waters. The maintenance of this wheel-wash should be addressed & wash water from any wheel wash must be directed to a suitable treatment facility.
15. Discharges to surface waters from wheel-washes and silt settlement ponds should be subject to S4 Discharge Licences.
16. It is important that the access/haul road, which is 1.1km in length is adequate to cope with the quantity and the type of traffic that this proposal will entail. Of concern to IFI is the fact that the 1.1km haul road between the proposed quarry and the N81 has not been included in the site layout plan.
17. The dirty waters generated from this 1.1km haul road, during times of high rainfall and also from waters used for dust suppression during the Summer months should be addressed.
18. IFI ask if the 1.1km haul road is a paved road or is a hard-core/clay road.
19. We request that all practical steps be taken to minimise dust being generated on this access road.

Yours faithfully



Donnachadh Byrne
Senior Fisheries Environmental Officer

Please note that any further correspondence regarding this matter should be addressed to Mr. Donnachadh Byrne, Senior Fisheries Environmental Officer, Inland Fisheries Ireland, 3044 Lake Drive, Citywest Business Campus, Dublin 24

