

## Appendix 7-4

# 2020 Breeding Bird Survey Report

## Breeding 2020 Bird Surveys Shronowen Wind Farm





ISSUE FORM	
Project number	19746
Document number	6005
Document revision	A
Document title	Shronowen Wind Farm - Bird Surveys - Breeding 2020
Document status	Final
Document prepared by	DO'B (MWP) 12-01-2021
Document checked by	MKe (MWP) 12-01-2021

## Table of Contents

<b>1</b>	<b>SUMMARY OF FINDINGS.....</b>	<b>2</b>
<b>2</b>	<b>INTRODUCTION .....</b>	<b>3</b>
<b>3</b>	<b>PURPOSE OF SURVEY .....</b>	<b>3</b>
<b>4</b>	<b>CONSTRAINTS.....</b>	<b>4</b>
<b>5</b>	<b>SURVEY DESIGN .....</b>	<b>4</b>
<b>6</b>	<b>SCOPING TO IDENTIFY TARGET SPECIES.....</b>	<b>5</b>
<b>6.1</b>	<b>Criteria for Selection of Target Species .....</b>	<b>5</b>
6.1.1	Legislative Protection and Conservation Status.....	5
6.1.2	Potential effects of wind farms on birds.....	6
6.1.3	Existing data, Records and Expert Knowledge .....	6
<b>7</b>	<b>SITE RECONNAISSANCE SURVEY.....</b>	<b>6</b>
<b>8</b>	<b>DESK STUDY.....</b>	<b>6</b>
<b>8.1</b>	<b>Description of the Survey Area .....</b>	<b>6</b>
<b>8.2</b>	<b>Bird Sensitivity to Wind Energy Development .....</b>	<b>10</b>
<b>8.3</b>	<b>Sites of International Importance in Proximity to the Survey Area.....</b>	<b>11</b>
8.3.1	Special Protection Areas (SPAs) - Birds Directive Species .....	11
8.3.2	Important Bird and Biodiversity Areas (IBAs) and Ramsar Sites .....	12
<b>8.4</b>	<b>Species known from the AREA.....</b>	<b>13</b>
8.4.1	In-house Expert Knowledge .....	13
<b>9</b>	<b>SELECTION OF SURVEY TYPES.....</b>	<b>14</b>
<b>9.1</b>	<b>Distribution and Abundance Surveys .....</b>	<b>14</b>
9.1.1	Moorland Breeding Birds .....	14
9.1.2	Raptors and Owls .....	14
9.1.3	Breeding Divers .....	14
9.1.4	Woodland Passerines.....	15
9.1.5	Nocturnal Species.....	15
9.1.6	Lowland and Farmland Birds.....	15
<b>10</b>	<b>SELECTION OF TARGET SPECIES.....</b>	<b>16</b>
<b>10.1</b>	<b>Primary Target Species.....</b>	<b>16</b>
<b>10.2</b>	<b>Secondary Target Species .....</b>	<b>17</b>
<b>11</b>	<b>VANTAGE POINT (VP) SURVEYS .....</b>	<b>24</b>
<b>11.1</b>	<b>Vantage Point (VP) Methodology .....</b>	<b>25</b>
11.1.1	Behaviour Codes .....	26
11.1.2	Habitat Codes <sup>23</sup> .....	26

---

<b>12</b>	<b>RESULTS: TARGET SPECIES ACTIVITY .....</b>	<b>27</b>
<b>12.1</b>	<b>Primary Target Species .....</b>	<b>27</b>
12.1.1	Hen harrier Observations.....	27
12.1.2	Kestrel Observations.....	29
12.1.3	Sparrowhawk Observations.....	30
<b>12.2</b>	<b>Secondary Target Species .....</b>	<b>33</b>
12.2.1	Cormorant Observations .....	33
12.2.2	Snipe Observations .....	34
<b>12.3</b>	<b>Other species observed .....</b>	<b>37</b>
12.3.1	Mallard Observations .....	37
12.3.2	Lesser black-backed gull Observations .....	38
12.3.3	Grey heron, Observations.....	39
<b>13</b>	<b>DISCUSSION .....</b>	<b>41</b>
<b>13.1</b>	<b>Primary Target Species.....</b>	<b>41</b>
<b>13.2</b>	<b>Secondary Target Species.....</b>	<b>42</b>
<b>13.3</b>	<b>Other species observed .....</b>	<b>42</b>
<b>14</b>	<b>REFERENCES.....</b>	<b>44</b>

## Table of Figures

Figure 1: Site Location in red. ....	3
Figure 2: Corine Landcover (2006) [from EPA Maps] .....	7
Figure 3: Bird Sensitivity to Wind Energy Development.....	10
Figure 4: Bird Sensitivity to Wind Energy Development.....	11
Figure 5: VP Locations.....	25
Figure 6: Hen harrier flight paths.....	28
Figure 7: Kestrel flight paths .....	29
Figure 8: Sparrowhawk flight paths .....	30
Figure 9: Cormorant flight paths.....	33
Figure 10: Snipe areas of activity .....	34
Figure 11 Mallard flight paths.....	37
Figure 12 Lesser black-backed gull flight paths .....	38

## Table of Tables

Table 1: Determining the sensitivity and importance of bird species (adapted from Percival, 2003) ...	5
Table 3: Target Species Ratings and Rationale for the Ratings Assigned .....	18
Table 4: Total Observation Time.....	27
Table 5 : Total Observation Time.....	29
Table 6: Total Observation Time.....	30
Table 7: Summary characteristics of hen harrier flights observed .....	32
Table 8: Summary characteristics of kestrel flights observed .....	32
Table 9: Summary characteristics of sparrowhawk flights observed .....	32
Table 10: Total Observation Time.....	33
Table 11: Total Observation Time.....	34
Table 12: Summary characteristics of cormorant flights observed .....	36
Table 13: Summary characteristics of snipe flights observed.....	36
Table 14: Total Observation Time.....	37
Table 15: Total Observation Time.....	38
Table 16: Total Observation Time.....	39
Table 17: Summary characteristics of Mallard flights observed.....	40
Table 18: Summary characteristics of Lesser black-backed gull flights observed .....	40
Table 19: Summary characteristics of Grey heron flights observed .....	40

## List of Appendices

<b>Appendix 1</b>	NPWS Recommended Methodology for Assessment of Impacts of Proposed Windfarms
<b>Appendix 2</b>	Survey Field Sheets
<b>Appendix 3</b>	Vantage Point Survey Summary
<b>Appendix 4</b>	Target/Secondary Species Observations
<b>Appendix 5</b>	Flight Paths and Activity Areas

- Appendix 6** Non-Target Species of Conservation Concern recorded during VP Surveys
- Appendix 7** List of All Species Recorded



NOTE: The following conventions have been followed with regard to species.

- 1. First instance of any species name in the text: Common name followed by full form Scientific Name**

Daisy (*Bellis perennis*)

- 2. 2<sup>nd</sup> instance: Common name followed by abbreviated Scientific Name**

Daisy (*B. perennis*)

- 3. Within tables: 1 or 2 above depending on circumstance.**

- 4. In Headings and within body of text: Unless first instance – Common name only**

Daisy

## 1 SUMMARY OF FINDINGS

Only three of the 13 Primary Target Species<sup>1</sup> and two of the 15 Secondary Target Species were recorded during the survey period and the numbers of observations of individual Target Species, and the activity of bird species generally, was extremely low. These species are as follows:

- Primary Target Species:
  - Hen harrier (*Circus cyaneus*): 1 observation;
  - Kestrel (*Falco tinnunculus*): 3 observations;
  - Sparrowhawk (*Accipiter nisus*): 1 observation;
  
- Secondary Target Species
  - Cormorant (*Phalacrocorax carbo*): 1 observation; and
  - Snipe (*Gallinago gallinago*): 1 recording.

In addition, non-target species namely, mallard (*Anas platyrhynchos*), lesser black-backed gull (LBBG) (*Larus fuscus*) and grey heron (*Ardea cinerea*) were also recorded.

While the full results of the survey are described in comprehensive detail in **Section 12**, a brief summary is presented here for information and for ease of review.

Hen harrier was recorded on one occasion which comprised a brief observation of an adult male which didn't extend beyond 30 seconds. This male was observed hunting low over the bog at <5m height to the northeast inside the site boundary. This low flight, hugging the ground while hunting is typical as they conceal themselves from predators. Kestrel was recorded on three occasions (one of which was an incidental sighting), all observations were quite short, the longest of which lasted 17seconds. All observations of kestrel were made to the east of the site inside the site boundary from VP1. Kestrel flight heights ranged from 0m-20m. Sparrowhawk was observed flying on 1 occasion. This agile hunter was only observed for a brief 15seconds to the south of the site from VP2. Sparrowhawk flight heights ranged from 0m-20m.

There was one recording of cormorant and snipe during the breeding 2020 survey period. A cormorant was observed in flight during from VP1 to the east of the site and this occurred inside the site boundary. This cormorant was observed flying over bog habitat in the east of the site. There was one recording of snipe during this survey period. Snipe drumming was heard from two areas inside the site boundary to the west of the site.

Mallard were observed on one occasion. This observation was made from VP1 to the east of the site. Mallard flight heights fall within 0m-20m as it flew and was observed on the ground over bog habitat. Lesser black-backed gulls were observed on one occasion. This observation was made from VP1 to the east of the site. The flight heights fall within 0m-30m as it flew over bog habitat. There was one observation of grey heron from VP3 to the east of the site. The flight heights fall within <10m and the heron was flying over bog habitat and on the ground.

---

<sup>1</sup> See **Section 10**

## 2 INTRODUCTION

Malachy Walsh and Partners, Engineering and Environmental Consultants, were commissioned by Emerging Markets Power (NI) Ltd., to conduct bird surveys, during the summer 2020<sup>2</sup>, at the location of a proposed wind farm development at Shronowen Bog near Ballylongford, County Kerry, (Irish Grid Co-ordinates: R 00498 40715). The survey area, outlined in red, in **Figure 1**, below, includes the proposed development site and areas adjacent. This report presents the results of the breeding 2020 survey. Previous reports (report ref. 19746-6002-A) has been completed for the winter 2018-2019 survey, (report ref. 19746-6003-A) has been completed for the breeding 2019 survey and (report ref. 19746-6004-A) has been completed for the winter 2019-2020.

This report comprises a description of those surveys and the results.




Figure 1: Site Location in red.

## 3 PURPOSE OF SURVEY

The survey was designed to determine the mix of species present and their behaviours and distribution within the survey area during the survey period. As reliable comparisons can then be made between these data and any subsequent survey data and, collectively, these will form a baseline upon which any future monitoring/multiyear surveys may be compared and, in the event of a consent application, will inform any impact assessments. The survey was conducted in compliance with the primary

<sup>2</sup> Summer survey period: April to September.

guidance used by the competent authorities in Ireland when assessing planning applications for a wind farm in circumstances where the impacts on avian ecology are germane, namely SNH (2017).

In summary the survey design will identify the species assemblage and the spatial and temporal distribution of activity. The range of methods used and survey effort involved are site and species specific and are informed by a desk study, site reconnaissance, by extensive survey experience in the surrounding area and by knowledge of the bird assemblage present in the north Kerry area.

#### 4 CONSTRAINTS

Surveyors did not have permission to access any lands outside the client's control. However, this did not impose a significant constraint on sampling as these lands comprise, almost exclusively, agricultural grassland habitats and it was expected, in light of the fact that several of the vantage points are located close to these agricultural habitats, that the typical species associated with these areas would be detected during the vantage point surveys.

#### 5 SURVEY DESIGN

Compliance with SNH (2017) requires that two main broad survey types are included in the survey design.

- **Distribution and Abundance Surveys.** These are surveys to record numbers and distribution of breeding, wintering and migrant birds using the site. They will allow the evaluation of a site's importance and provide information to help quantify predicted impacts from disturbance and displacement.
- **Vantage Point (VP) Surveys.** These surveys, which, in the case of the Shronowen site, will be required, comprise a series of watches from a fixed location to quantify the flight activity of birds at a proposed development site, which provides data to estimate the collision risk.

The decision as to which of the survey methodologies are required is based on the outcome of a scoping exercise which determines which species are considered likely to use the habitats in the study area.

The survey includes a number of methodologies, described in **Sections 9.1** and **11**, below, that have been selected, from the list of survey types identified in SNH (2017), for their capacity to detect and record the activities of the species expected to be present in the survey area during the survey period. The methodologies selected ensured that a structured approach to survey work was implemented throughout. While all aspects of the activities of the observed Target Species were recorded, the primary aim of the surveys is to understand bird use of the survey area; a secondary purpose is to provide data for Collision Risk Modelling (CRM). A detailed description of how information on flight behaviours was recorded will be provided, under the appropriate headings, in **Section 11**.

The survey design and execution is informed by extensive in house experience across a broad range of comparable surveys conducted in similar areas with specific reference to those carried out in the north Kerry and west Limerick.

## 6 SCOPING TO IDENTIFY TARGET SPECIES

Compliance with SNH (2017) requires that prior to the commencement of surveys a scoping exercise is carried out to determine a broad overview of which species are likely to be at the site, their likely sensitivity to impacts from wind farms and the proximity of relevant designated sites. This allows the selection of Target Species (see **Section 9**) and these species will form the basis of the survey programme.

### 6.1 CRITERIA FOR SELECTION OF TARGET SPECIES

#### 6.1.1 Legislative Protection and Conservation Status

When compiling the list(s) of Target Species, consideration of legislative protection and conservation status are of primary importance, in this regard, there are three important species lists from which Target Species may be drawn:

- Listed in Annex 1 of the EC Birds Directive;
- Protected under the Wildlife Acts, 1976 to 2012; and
- Red-listed species as per Colhoun & Cummins (2013)<sup>3</sup>.

Within the scope of the criteria outlined above, SNH (2017) recommends that the Target Species should be limited to:

- Those species which are afforded a higher level of legislative protection; and
- Those species which, as a result of their behaviours, are more likely to be subject to impact from wind farms.

A precautionary approach was adopted and the selection followed the guidance set out for determining the sensitivity and importance of bird species as outlined in Percival (2003). Percival's methodology was considered alongside the other literature relating to the effects of wind farms on birds as reviewed in Whitfield and Madders (2006) and Drewitt and Langston (2006). These sensitivities were evaluated using the criteria set out in **Table 1**. When compiling the list cognisance was also taken of the constraints imposed on the distributions on the species due to their known habitat requirements and distributions.<sup>4</sup> Those species selected as Primary Target Species are listed in **Section 10.1** and those selected as Secondary Target Species are listed in **Section 10.2**.

**Table 1: Determining the sensitivity and importance of bird species (adapted from Percival, 2003)**

Sensitivity	Determining Factor
VERY HIGH	Where the site is an SPA
	Species present in nationally important numbers (>1% Irish population)
HIGH	Ecologically sensitive species (e.g. divers, common scoter, golden eagle, hen harrier, chough and roseate tern)
	EU Bird Directive Annex I species
	Red-listed Species of Conservation Concern
MEDIUM	Amber-listed Species of Conservation Concern

<sup>3</sup> Birds on the Red List birds are those of highest conservation concern, Amber List birds are of medium conservation concern and the Green List birds are not considered threatened.

<sup>4</sup> As outlined at <https://www.birdwatchireland.ie>

	Species present in locally important numbers (>1% of county population)
LOW	Amber-listed Species

### 6.1.2 Potential effects of wind farms on birds

Detailed knowledge of bird distribution and flight activity is necessary in order to predict the potential effects of a wind farm on birds. However, the scope and scale of the survey data taken and the suite of species on which data is collected should be informed by the analysis that wind farms present three main potential risks to birds (Drewitt & Langston 2006, 2008; Band *et al.* 2007, cited in SNH, 2017). These are:

- Direct habitat loss through construction of wind farm infrastructure;
- Displacement (sometimes called indirect habitat loss) if birds avoid the wind farm and its surrounding area due to turbine construction and operation. Displacement may also include barrier effects in which birds are deterred from using normal routes to feeding or roosting grounds; and
- Death through collision or interaction with turbine blades and other infrastructure.

Due to the unique ecology of each species each will have different sensitivities to each of these three impact sources.

### 6.1.3 Existing data, Records and Expert Knowledge

Cognisance must also be taken of existing data and records, expert knowledge of the species assemblage present in the wider north Kerry/west Limerick area, and the influence on bird distribution of the habitat mix within and adjacent to the survey area whose presence within the survey area is reasonably foreseeable in light of the habitats present, both within the survey area and in the surrounding landscape.

## 7 SITE RECONNAISSANCE SURVEY

As per SNH (2017) requirements that, prior to the commencement of surveys, a scoping exercise is carried out reconnaissance of the site and its surrounds was carried out by MWP staff ecologists. These visits enabled an evaluation to be made of the habitat characteristics of the site and the identification of VP locations considered suitable to provide maximum site coverage. As stipulated by the client, all surveys were undertaken within lands within which landowner's permission had been arranged or on public roads. Access was not permitted to private lands outside the client's control.

## 8 DESK STUDY

### 8.1 DESCRIPTION OF THE SURVEY AREA

The site largely comprises cut-over bog (*sensu* Fossitt, 2000), which in its original form was a blanket bog, but which is now substantially cut-over and significantly altered by turf cutting. It is situated within a landscape dominated by agricultural grassland habitats and with some commercial conifer plantations against which the bog itself abuts (see **Figure 2** for Corine Landcover)<sup>5</sup>. The topography of the site is essentially flat, albeit, with the slight peat dome that is a characteristic of the lowland bog


<sup>5</sup> Areas of bog are shown in purple, forestry in green and pastureland is shown in yellow.

type. The site is intersected by a network of access tracks of robust construction that, while too rough for cars, are, for the most part, in good condition.

Turbary rights pertain to the entire site and much of the original peat mass has been removed. While a large central area remains relatively uncut, a crisscross network of drains intersects the site and significant proportion of the bog now comprises a mix of exhausted banks or banks that are currently being, or historically have been, worked. A significant effect of the peat extraction is the extent to which the water table across the site has been lowered permanently. Because the water table plays an important role in aerobic and anaerobic processes in a bog, the lowering of the water table within the peat boundary, between the upper aerobic acrotelm (living) layer and the underlying, water-logged and compacted, catotelm (dead) layer, has fundamentally altered the peat forming capacity of Shronowen Bog.

While the dominant current practice is removal of peat by excavator to a hopper from which the peat is then extruded (see **Drone Flown Image 1**) there is clear evidence of historic sausage cutting in the eastern part of the site (see **Drone Flown Image 2**). **Aerial Image 1** illustrates the extent to which, over time, the peat mass has been removed progressively and incrementally from the edge of the bog (represented in blue) to the interior area of the peat mass.

Corine Landcover



November 23, 2018




Figure 2: Corine Landcover (2006) [from [EPA Maps](#)]



**Aerial Image 1: Typical view showing distinct signature of turf banks progressing from edge to centre at northern section of Shronowen Bog. (Red circle: approximate location of Drone Image 1; Yellow circle approximate location of Drone Image 2).**



**Drone Flown Image 1: Extruded turf with excavated bank adjacent (2019)**





**Drone Flown Image 2: Evidence of historic sausage cutting (parallel 'scars' aligned left to right)**

The vegetation communities that the bog supports are constrained by the nutrient poor conditions that pertain and the cover currently comprises a relatively uniform and homogenous cover of Purple Moor-grass (*Molinia caerulea*). While heather is present, surveys indicate that it is not a significant component in the overall plant mix. A few isolated treelines are present; these consist primarily of birch (*Betula* spp.) and all are of a relatively low stature with an average canopy height in the region of 5 m. Areas of willow scrub (*Salix* spp.) are also present; however, these are primarily distributed within the transitional marginal habitats that fringe the bog, in the interface areas between the agricultural and commercial forestry habitats and the bog itself. Willow shrub lines also fringe the sides of the tracks in many places. A variety of grasses and ruderal species have colonised the margins along the sides of the tracks where disturbance has disrupted the dominance of the indigenous vegetation that dominates the remainder of the site. A significant proportion of the site comprises bare unvegetated ground which is present in areas where sustained peat extraction has been occurring recently.

While the site is intersected by a network of man-made drains, the only natural water body within the site is an unnamed tributary<sup>6</sup> of the Ballylongford River which drains from a point of origin in the north of the site. Apart from some localised ponding of water in some of the lower lying peat banks no established ponds or other bodies of standing water were noted during the site surveys and none are visible in the range of aerial imagery reviewed<sup>7</sup>. While stands of Bulrush (*Typha latifolia*) are present in some trackside drains in the western part of the site, the individual stands are generally small and localised and the distribution within the site is somewhat uneven and diffuse.

In summary the site is, both topographically and ecologically, relatively homogeneous, a characteristic that inhibits species diversity not only in terms of the floristic communities and insect species but also

<sup>6</sup> River Waterbody Code: IE\_SH\_24B030700 <https://gis.epa.ie/EPAMaps/>


<sup>7</sup> OSI aerial imagery (1995 to 2012); Google imagery (2017); Bing (undated)

in the variety of bird species, particularly passerines, likely to be present. It is unlikely to provide significant foraging, roosting or breeding habitats for many bird species.

**8.2 BIRD SENSITIVITY TO WIND ENERGY DEVELOPMENT**

The National Biodiversity Data Centre’s (NBDC) online mapper<sup>8</sup> includes a layer which provides information on sensitivity to wind energy development. This layer is derived from a collation of existing distributional data, which indicates, by assessing the characteristics of a selected number of the most-sensitive bird species, whether protected birds are likely to be sensitive to wind energy developments in the areas mapped. The mapping layer is derived from McGuinness *et al.* (2015) and while it does not include all vulnerable species - due to data and other issues - and does not replace SEA, AA or EIA requirements nor the need to tailor survey and research to specific sites, it provides a useful metric to rank sites, at the initial scoping stage, in terms of their potential sensitivity to wind energy development. The layer has four sensitivity ratings, namely Low, Medium, High and Highest. These ratings are mapped at 2km grid square resolution for which ‘All Birds Sensitivity Scores’ (ABSS) are provided.

The survey area and the geographical area extending away from it is categorised as Low Sensitivity (see **Figure 3** and **Figure 4**, below) and the ABSS is 14.8.



**Figure 3: Bird Sensitivity to Wind Energy Development (from <http://maps.biodiversityireland.ie/#/Map>)**

<sup>8</sup> <https://maps.biodiversityireland.ie/Map>

## Bird Sensitivity to Wind Energy2




Figure 4: Bird Sensitivity to Wind Energy Development (from <http://maps.biodiversityireland.ie/#/Map>)

### 8.3 SITES OF INTERNATIONAL IMPORTANCE IN PROXIMITY TO THE SURVEY AREA

#### 8.3.1 Special Protection Areas (SPAs) - Birds Directive Species

The survey area is situated approximately 3 km due south of the site boundary of the River Shannon and River Fergus Estuaries SPA (004077) which is selected for the conservation of the non-breeding, wintering populations<sup>9</sup> of 21 Special Conservation Interest (SCI) species and for the SCI Wetlands [A999] habitats that are a resource for the regularly-occurring migratory water birds that utilise the SPA. The proposal site is also approximately 10 km to the west of the Stack's to Mullaghareirk Mountains, West Limerick Hills and Mount Eagle SPA (004161) which is selected for the conservation of a resident, breeding, population of one SCI species, namely hen harrier (*Circus cyaneus*) [A082]<sup>10</sup>.

The SCI species for which the River Shannon and River Fergus Estuaries SPA (004077) is selected are:

- Cormorant (*Phalacrocorax carbo*) [A017]
- Whooper Swan (*Cygnus cygnus*) [A038]
- Light-bellied Brent Goose (*Branta bernicla hrota*) [A046]
- Shelduck (*Tadorna tadorna*) [A048]
- Wigeon (*Anas penelope*) [A050]
- Teal (*Anas crecca*) [A052]
- Pintail (*Anas acuta*) [A054]
- Shoveler (*Anas clypeata*) [A056]

<sup>9</sup> <https://www.npws.ie/sites/default/files/protected-sites/natura2000/NF004077.pdf>

<sup>10</sup> <https://www.npws.ie/protected-sites/spa/004161>

- Scaup (*Aythya marila*) [A062]
- Ringed Plover (*Charadrius hiaticula*) [A137]
- Golden Plover (*Pluvialis apricaria*) [A140]
- Grey Plover (*Pluvialis squatarola*) [A141]
- Lapwing (*Vanellus vanellus*) [A142]
- Knot (*Calidris canutus*) [A143]
- Dunlin (*Calidris alpina*) [A149]
- Black-tailed Godwit (*Limosa limosa*) [A156]
- Bar-tailed Godwit (*Limosa lapponica*) [A157]
- Curlew (*Numenius arquata*) [A160]
- Redshank (*Tringa totanus*) [A162]
- Greenshank (*Tringa nebularia*) [A164]
- Black-headed Gull (*Chroicocephalus ridibundus*) [A179]

This list includes species from a number of groups including, *inter alia*, swans, geese, waders and gulls. While the foraging or breeding behaviours of most of these populations are not strongly associated with the habitats available in the survey area (NPWS, 2012), it is possible that some of the species do overfly the site when commuting between roosting and foraging grounds.

### 8.3.2 Important Bird and Biodiversity Areas (IBAs) and Ramsar Sites

#### 8.3.2.1 Important Bird and Biodiversity Areas (IBAs)

The Important Bird and Biodiversity Areas (IBA) Programme is a BirdLife International initiative aimed at identifying and protecting a network of critical sites for the conservation of the world's birds. A total of 140 Important Bird Areas (IBAs) have been identified in Ireland, covering an area of about 4,309km<sup>2</sup>, equivalent to 6% of the land area. These sites are important for breeding seabirds and for wintering wildfowl.

There are two IBA site within 15 km of the survey area, namely the Shannon and Fergus Estuaries (IE08) and The Stacks to Mullaghareirk Mountains, West Limerick Hills and Mount Eagle (IBA Criteria C6 (2009)). Shannon and Fergus Estuaries (IE08) is encompassed within the significantly larger River Shannon and River Fergus Estuaries SPA (004077), is one of the most important sites in Ireland for wintering and migrating waterfowl and it supports 10 species in numbers of international importance all which are also protected under the SPA designation. These species are<sup>11</sup>:

- Whooper swan (*C. cygnus*)
- Brent goose (*Branta bernicla*)<sup>12</sup>
- Scaup (*A. marila*)
- Golden plover (*P. apricaria*)
- Knot (*C. canutus*)
- Dunlin (*C. alpina*)
- Black-tailed godwit (*L. limosa*)
- Bar-tailed godwit (*L. lapponica*)

<sup>11</sup> <http://datazone.birdlife.org/site/factsheet/shannon-and-fergus-estuaries-iba-ireland/details>

<sup>12</sup> Light-bellied brent goose, a species for which the SPA site (004077) is selected, is a sub species of brent goose

- Curlew (*N. arquata*)
- Redshank (*T. totanus*)

A further 13 species occur in numbers of national importance, including, *inter alia*,

- Greylag goose (*Anser anser*)
- Shelduck (*T. tadorna*)
- Wigeon (*A. penelope*)
- Teal (*A. crecca*)
- Pintail (*A. acuta*)
- Shoveler (*A. clypeata*)
- Lapwing (*V. vanellus*)
- Greenshank (*T. nebularia*)<sup>13</sup>

Of these species only greylag goose is not an SCI species for which the River Shannon and River Fergus Estuaries SPA (004077) is selected.

The Stacks to Mullaghareirk Mountains, West Limerick Hills and Mount Eagle (IBA Criteria C6 (2009)) is encompassed within The Stacks to Mullaghareirk Mountains, West Limerick Hills and Mount Eagle SPA (004161), both sites are important for breeding hen harrier (*Circus cyaneus*)<sup>14</sup>.

#### 8.3.2.2 Ramsar Sites

The Ramsar Convention on Wetlands of International Importance especially as Waterfowl Habitat is an international treaty for the conservation and sustainable use of wetlands. The Ramsar Convention was ratified by Ireland in 1984 and came into force for Ireland on 15 March 1985. Ireland presently has 45 sites designated as Wetlands of International Importance, with a surface area of 66,994 hectares.

No Ramsar site is located within 15km of the survey area.

## 8.4 SPECIES KNOWN FROM THE AREA

### 8.4.1 In-house Expert Knowledge

On the basis of extensive formal and informal in house expertise the following species are known to be present in the wider geographical area extending away from the survey area:

- Barn owl (*Tyto alba*)
- Kestrel (*F. tinnunculus*)
- Merlin (*Falco columbarius*)
- Mute swan (*Cygnus olor*)
- Sparrowhawk (*A. nisus*)
- Short-eared owl (*Asio flammeus*)

<sup>13</sup> No further information on the other species is provided on the website.

<sup>14</sup><http://datazone.birdlife.org/site/factsheet/stacks-to-mullaghareirk-mountains-west-limerick-and-mount-eagle-iba-ireland/details>

A hinterland survey undertaken to inform the winter 2018-19 survey detected a waterbird site used by whooper swan comprising agricultural grassland fields about 0.5-1km northwest of the site where a flock of between 11 and 15 individuals were observed on the ground and foraging during the months of six separate dates in February and March 2019.

## 9 SELECTION OF SURVEY TYPES

As outlined, previously, in **Section 5** compliance with SNH (2017) requires that two main broad survey types are included in the survey design.

- Distribution and Abundance Surveys; and
- Vantage Point (VP) Surveys.

Within these broad types SNH (2017) lists a number of different methodologies and these are outlined hereunder. In each case a site specific assessment is carried out and recommendations are made as to which of the survey types should be carried out

### 9.1 DISTRIBUTION AND ABUNDANCE SURVEYS

#### 9.1.1 Moorland Breeding Birds

The site is of limited suitability for breeding waders, skuas, gulls, or red grouse (grouse would have been heard in late winter calling if present) and thus a dedicated survey was not carried out.

#### 9.1.2 Raptors and Owls

Of the four species of owl known in Ireland, namely barn owl (*Tyto alba*), snowy owl (*Nyctea scandiaca*), long-eared owl (*Asio otus*) and short-eared owl (*Asio flammeus*) only barn owl and long-eared owl are purely nocturnal. Surveys for nocturnal species are assessed in **Section 11**, below.

With regard to snowy owl (*Nyctea scandiaca*) it is noted that because this species is a rare winter visitor, mainly to western counties such as Mayo<sup>15</sup>, it is not expected to be present. With regard to short-eared owl, it is a scarce winter visitor throughout Ireland and rare breeding species, mainly in the south and east, should it be present in the survey area it is expected that this species and other raptors would be detected by the VP surveys described in **Section 11**, below.

#### 9.1.3 Breeding Divers

This survey type was not required. Only one species from this group is known to breed in Ireland, namely red-throated diver (*Gavia stellata*). Very few pairs do breed in Ireland and those that have bred have been restricted to Co. Donegal<sup>16</sup>.

With regard to the likelihood that the other species from this group will frequent the site, the populations of these species are associated with shallow sandy bays and feed on open water plunging to catch fish or other food. Due to the specialised nature of their feeding techniques they are not expected to present at the site due to its terrestrial location and habitat mix.

<sup>15</sup> <https://www.birdwatchireland.ie/IrelandsBirds/Owls/SnowyOwl/tabid/1125/Default.aspx>

<sup>16</sup> <https://www.birdwatchireland.ie/Default.aspx?tabid=125>

#### 9.1.4 Woodland Passerines

While the site boundary does overlap with a number of commercial conifer plantations the buildable area does not overlap with any of them. In light of this and bearing in mind that surveys of woodland passerines, especially in commercial conifer forest, are generally not required (SNH, 2017) and because there is very little evidence that passerines are significantly affected by wind farms (DGE, 2014) it was concluded that this survey type was not required. In addition, because the VPs (see **Section 11**, below) are located adjacent to locations that are good examples of the typical, albeit limited, variation in habitats present within the survey area, it was expected that the typical species associated with these habitats and the broader more typical habitats would be detected during the VP surveys.

#### 9.1.5 Nocturnal Species

##### 9.1.5.1 Owls

Of the species of owl resident in Ireland only barn owl and long-eared owl are purely nocturnal. As a result any flights would not be observable and systematic flight path mapping would not be possible, therefore, neither was selected as Target Species. However, extensive in-house experience of the species mix present in the wider geographical area indicates that the survey area could be within the foraging territory of barn owl and, although equivalent knowledge on the presence of long-eared owl is not available, it is considered, on the basis of the precautionary principle, that surveys for both species should be undertaken.

The surveys were conducted, as per SNH (2017) and BirdWatch Ireland<sup>17</sup>, by listening for calling birds around dusk from February onwards during winter VP surveys. SNH (2017) further recommends that late evening surveys for calling juveniles in May-July can also be useful in detecting successful pairs; adults may also be active during this time. Should calling birds be detected, in the event that specific breeding sites are identified, surveys can be complemented by searches for signs of occupation, such as moulted feathers and pellets. If present, these evidences of occupancy in the environs of the site can be recorded.

##### 9.1.5.2 Other nocturnal species

Nightjar (*Caprimulgus europaeus*): as this species is a rare summer-visitor to uplands in southern Ireland<sup>18</sup> it was not expected to be present. Surveys were not required.

#### 9.1.6 Lowland and Farmland Birds

Surveys of farmland, moorland or woodland passerines are generally not required (SNH, 2017) and there is very little evidence that passerines are significantly affected by wind farms (DGE, 2014). However, in order to fully characterise the use of the survey area by birds, all species encountered were recorded; however, recording of these species was subsidiary to recording of Target Species and comprised recording of simple counts of species observed only. Because the VPs (see **Section 11**, below) are located adjacent to locations that are good examples of the typical, albeit limited, variation in habitats present within the survey area, it was expected that the typical species associated with these habitats and the broader more typical habitats would be detected during the VP surveys.

<sup>17</sup> <https://birdwatchireland.ie/birds/long-eared-owl/>

<sup>18</sup> <https://birdwatchireland.ie/birds/nightjar/>

## 10 SELECTION OF TARGET SPECIES

Target Species, for which comprehensive data were recorded, were limited to those species likely to be affected by wind farms. The habitat mix within and adjacent to the proposed development site, described in **Section 8.1**, allowed a preliminary assessment to be made, in 2018, prior to commencement of surveys at the site, of the bird populations likely to be present in the study area. This assessment was cognisant of the known habitat preferences of the species evaluated and the restrictions on their distributions that result from these preferences. This assessment when viewed in combination with the information on the proximity of relevant designated sites, outlined in **Section 8.3**, and those species known to be present in the wider area, identified in **Section 8.4**, allowed the selection of primary and, potentially, Secondary Target Species as per SNH (2017). In selecting species for inclusion in the Target Species lists a precautionary approach was adopted and the selection also followed the guidance set out for determining the sensitivity and importance of bird species as outlined in Percival (2003), Whitfield & Madders (2006) and Drewitt & Langston (2006). This evaluation is summarised in **Table 2**.

Because there is very little evidence that passerines are significantly affected by wind farms (DGE, 2014; SNH, 2017) and unless rare/restricted passerines are present surveys are not required (SNH, 2017) transects or point counts such as those outlined in Anon (2012) or Bibby *et al.* (2000) were not carried out. However, in order to fully characterise the species mix present in the survey area all species encountered, including passerines, were recorded. However, recording of these species is subsidiary to recording of Target Species and will comprise recording of simple counts of species observed. This element of the survey design is to provide the additional data on bird usage of the site that will be required for subsequent assessments of the impacts on the broad avian biodiversity of the survey area in the event that an application for planning permission is submitted. An example of the survey sheet is included in **Appendix 2**.

Those species selected as Primary Target Species are listed in **Section 10.1** and those selected as Secondary Target Species are listed in **Section 10.2**. The evaluation is summarised in **Table 2**.

### 10.1 PRIMARY TARGET SPECIES

The Primary Target Species are:

- Hen harrier (*C. cyaneus*)
- Merlin (*F. columbarius*)
- Kestrel (*F. tinnunculus*)
- Sparrowhawk (*A. nisus*)
- Short-eared owl (*A. flammeus*)
- Whooper Swan (*C. cygnus*)
- Mute Swan (*C. olor*)
- Light-bellied Brent Goose (*B. bernicla hrota*)
- Greylag goose (*A. anser*)
- Golden Plover (*P. apricaria*)
- Lapwing (*V. vanellus*)
- Curlew (*N. arquata*)
- Black-headed Gull (*C. ridibundus*)



## 10.2 SECONDARY TARGET SPECIES

The Secondary Target Species are:

- Cormorant (*P. carbo*)
- Shelduck (*T. tadorna*)
- Wigeon (*A. penelope*)
- Teal (*A. crecca*)
- Pintail (*A. acuta*)
- Shoveler (*A. clypeata*)
- Scaup (*A. marila*)
- Ringed Plover (*C. hiaticula*)
- Grey Plover (*P. squatarola*)
- Knot (*C. canutus*)
- Dunlin (*C. alpina*)
- Black-tailed Godwit (*L. limosa*)
- Bar-tailed Godwit (*L. lapponica*)
- Redshank (*T. totanus*)
- Greenshank (*T. nebularia*)
- Snipe (*G. gallinago*)<sup>19</sup>

While not included as target species surveys for the nocturnal barn owl and long-eared owl will be conducted as outlined in **Section 9.1.5**, above. In the event that either species is observed in daylight then any flight paths observed will be recorded as per **Section 11.1**, below.

---

<sup>19</sup> Dedicated dawn counting of drumming snipe in early spring will be carried out for this species as per **Section 6.1.1**, above.

**Table 2: Target Species Ratings and Rationale for the Ratings Assigned**

Raptors & Owls	Target Species Rating	Rationale
Hen harrier ( <i>C. cyaneus</i> )	Primary	Amber listed. EU Bird Directive Annex I species. Potential foraging and breeding habitat in survey area. Populations are vulnerable to habitat modifications that result from land use change (Wilson <i>et al.</i> , 2015). Raptors are among the species known to be most vulnerable to collision mortality at wind farms (Thaxter <i>et al.</i> , 2017). The construction and operation of wind turbines can impact on hen Harriers (displacement during construction and/or operation; collision with turbines). Known presence in wider geographical area year round <sup>20</sup> .
Merlin ( <i>F. columbarius</i> )	Primary	Amber listed. EU Bird Directive Annex I species. Potential foraging habitat in survey area but unlikely to breed in survey area or in area extending away from survey area. Raptors are among the species known to be most vulnerable to collision mortality at wind farms (Thaxter <i>et al.</i> , 2017). Known presence in wider geographical area during winter <sup>20</sup> .
Kestrel ( <i>F. tinnunculus</i> )	Primary	Amber listed. Potential foraging habitat in survey area. Potential breeding habitat in area extending away from survey area. Raptors are among the species known to be most vulnerable to collision mortality at wind farms (Thaxter <i>et al.</i> , 2017). Known presence in wider geographical area year round <sup>20</sup> .

<sup>20</sup> Known presence based on MWP in-house knowledge and experience.

Sparrowhawk ( <i>A. nisus</i> )	Primary	Amber listed. EU Bird Directive Annex I species. Potential foraging habitat in survey area. Potential breeding habitat in area extending away from survey area. Raptors are among the species known to be most vulnerable to collision mortality at wind farms (Thaxter <i>et al.</i> , 2017). Known presence in wider geographical area year round <sup>20</sup> .
Barn owl ( <i>T. alba</i> )	Not selected	Nocturnal species therefore flight lines not visible. While raptors are among the species known to be most vulnerable to collision mortality at wind farms (Thaxter <i>et al.</i> , 2017), barn owls are rarely affected by wind turbines <sup>21</sup> .
Long-eared owl ( <i>Asio otus</i> )	Not selected	Nocturnal species therefore flight lines not visible. Potential foraging habitat in survey area. Potential breeding habitat in area extending away from survey area.
Short-eared owl ( <i>Asio flammeus</i> )	Primary	Feeds mainly on small mammals in open habitats. Potential foraging habitat in survey area. Potential breeding habitat in area extending away from survey area. Known presence in wider geographical area <sup>20</sup> .
Swans and Geese	Target Species Rating	Rationale
Whooper Swan ( <i>C. cygnus</i> )	Primary	EU Bird Directive Annex I species. Nationally important population. Proximity of SPA selected for protection of this species. Grassland areas adjacent to the estuary are used by grazing Whooper Swans (Robinson <i>et al.</i> , 2004). The species is known to forage on grassland sites (Worden <i>et al.</i> , 2009) during the day. Possibility that the species overflies or transects through the survey area when commuting to foraging grounds further inland. Known poor flight manoeuvrability.

<sup>21</sup> <https://www.barnowltrust.org.uk/hazards-solutions/barn-owls-wind-turbines/>

Mute Swan ( <i>C. olor</i> )	Primary	Known presence in wider geographical area <sup>20</sup> . Possibility, albeit slight, that the species' flight lines intersect through the survey area when commuting between foraging grounds. Precautionary principle. Known poor flight manoeuvrability.
Light-bellied Brent Goose ( <i>B. bernicla hrota</i> )	Primary	EU Bird Directive Annex I species. Internationally important population <sup>22</sup> . Proximity of SPA selected for protection of this species. Possibility, albeit slight, that the species' flight lines intersect through the survey area. Known poor flight manoeuvrability.
Greylag goose ( <i>A. anser</i> )	Primary	Proximity of IBA selected for protection of this species. Possibility, albeit slight, that the species' flight lines intersect with the survey area. Known poor flight manoeuvrability. Precautionary principle.
<b>Cormorants</b>	<b>Target Species Rating</b>	<b>Rationale</b>
Cormorant ( <i>P. carbo</i> )	Secondary	EU Bird Directive Annex I species. Nationally important migratory population. Nationally important resident breeding population. Proximity of SPA selected for protection of this species. Possibility that the species' flight lines intersect with the survey area.
<b>Ducks</b>	<b>Target Species Rating</b>	<b>Rationale</b>
<b>Amber listed:</b> Shelduck ( <i>T. tadorna</i> ) Scaup ( <i>A. marila</i> ) Teal ( <i>A. crecca</i> )	Secondary	Notwithstanding the proximity of SPA selected for protection of these species and the national importance of the populations for which the SPA is selected, all are exclusively associated with open water habitats not present within the survey area or in the area extending away from it. Very limited likelihood that the species' flight lines intersect with the survey area.

<sup>22</sup> <https://www.npws.ie/sites/default/files/protected-sites/synopsis/SY004077.pdf>

<b>Red listed:</b> Pintail ( <i>A. acuta</i> ) Shoveler ( <i>A. clypeata</i> ) Wigeon ( <i>A. penelope</i> )		
Waders	Target Species Rating	Rationale
Golden Plover ( <i>P. apricaria</i> )	Primary	Red listed. EU Bird Directive Annex I species. Nationally important population. Proximity of SPA selected for protection of species. Possibility that the species overflies or transects through the survey area. Potential foraging habitat in survey area but unlikely to breed in survey area or in area extending away from survey area. Known presence in wider geographical area in winter <sup>20</sup> .
Curlew ( <i>N. arquata</i> )	Primary	Red listed; EU Bird Directive Annex I species. Nationally important population. Proximity of SPA selected for protection of species. Possibility that the species overflies or transects through the survey area. Potential foraging habitat in area extending away from survey area survey area but unlikely to breed in survey area or in area extending away from survey area. Known presence in wider geographical area <sup>20</sup> .

Lapwing ( <i>V. vanellus</i> )	Primary	<p>Red listed.</p> <p>EU Bird Directive Annex I species.</p> <p>Nationally important population.</p> <p>Proximity of SPA selected for protection of species.</p> <p>Possibility that the species overflies or transects through the survey area to foraging grounds where the variety of soil and surface-living invertebrates this species predares are available.</p> <p>Potential foraging habitat in area extending away from survey area survey area but unlikely to breed in survey area or in area extending away from survey area.</p>
<p><b>Green listed:</b></p> <p>Ringed Plover (<i>C. hiaticula</i>)</p> <p>Greenshank (<i>T. nebularia</i>)</p> <p><b>Amber listed:</b></p> <p>Grey Plover (<i>P. squatarola</i>)]</p> <p>Knot (<i>C. canutus</i>)</p> <p>Black-tailed Godwit (<i>L. limosa</i>)</p> <p>Bar-tailed Godwit (<i>L. lapponica</i>)</p> <p><b>Red listed:</b></p> <p>Dunlin (<i>C. alpina</i>)</p> <p>Redshank (<i>T. totanus</i>)</p>	Secondary	<p>Notwithstanding the proximity of SPA selected for protection of these species and the international and national importance of the populations for which the SPA is selected, all are essentially obligate feeders on marine and estuarine benthic invertebrates. Very limited likelihood that the species' flight lines intersect with the survey area.</p>
Gulls	Target Species Rating	Rationale

Black-headed Gull ( <i>C. ridibundus</i> )	Primary	Red listed. EU Bird Directive Annex I species. Proximity of SPA selected for protection of species. Nationally important population. Possibility that the species overflies or transects through the survey area to alternative foraging grounds inland from the estuary.
--	---------	---

## 11 VANTAGE POINT (VP) SURVEYS

VP surveys are designed to quantify the level of flight activity and its distribution over a survey area (SNH, 2017). The survey type comprises a series of watches from fixed locations that are repeated on a scheduled basis that are focused on recording flight behaviours that intersect with the turbine rotor envelope. The aim of the survey design is to set out a standard methodology for recording both the quantitative and qualitative aspects of these behaviours in order to produce sufficient information to assess the potential effects of the development on Target Species particularly with regard to collision risk. It also allows a determination to be made as to whether regular flight lines for any species intersect with the survey area.

VP surveys allow the collection of accurate data on Target Species that will enable estimates to be made of:


- The time spent flying over the survey area;
- The relative use of different parts of the survey area; and
- The proportion of flying time spent within the upper and lower height limits as determined by the rotor diameter and the hub height.

On the basis of extensive local knowledge and experience of the distribution of hen harrier in the north Kerry area and due to the proximity of an SPA designated for the protection of this species, VP surveys were required (SNH, 2017). To this end surveys from three VP locations were conducted during the survey period. The VPs, shown in **Figure 5** were selected to ensure that the fields of view covered all of the flight activity within the survey area (buildable area & 500m buffer) and are located such that no point within the survey area is greater than 2 km from a VP. When selecting the VP locations the visibility of the rotor swept area is critical; visibility at ground level is not. However, due to the almost uninterrupted fields of view afforded by the relatively flat topography of the site visibility to ground level is possible over much of the site. As per SNH (2017) 36 hours per VP were completed during the survey period.

Because bird species have varied seasonal, and within day, activity patterns the timing of survey sessions were adjusted to occur at times when birds are likely to be most active. Because bird flight behaviours change in response to wind conditions, particularly with regard to flight heights, weather will also be a factor in the scheduling of surveys.

The VP methodology outlined in **Section 11.1** also followed the NPWS Recommended Methodology for Assessment of Impacts of Proposed Windfarms included in **Appendix 1**. While the primary focus of the VP surveys were the Target Species listed in **Section 10** all species encountered were recorded on a presence/absence basis on separate field sheets (see **Appendix 2**).





**Figure 5: VP Locations**

### 11.1 VANTAGE POINT (VP) METHODOLOGY

The methodology is of particular use in providing details of the number of species and the extent to which birds use the site. It also provides supplementary information on flight activity and behaviour. The longer the overall survey period of VP surveys, the more accurate and precise the sample of flight behaviour.

The VPs are located at positions that provided clear views of turbine hub heights and blade swept area over the survey area. The surveyors based themselves at each VP for a fixed period of 6 hours each month of the survey period. VP sessions were conducted as a series of watches each of not more than 3 hours continuous duration at a time. There were breaks of at least 30 minutes between watches to minimise observer fatigue and a short 'settling in' period of approximately 10 minutes at each VP, before watches started, to allow the surveyor to organise and annotate field sheets, mapping, etc. and to ensure any disturbance from moving around the site had passed.

VP watches were conducted under conditions of good ground visibility (>2km) on days when the cloud base was high enough to allow observation of the full survey area and observations were to be suspended during periods of poor visibility and/or heavy rain. In order to ensure that any activity by soaring birds was sampled, surveys were undertaken in a range of wind conditions and on showery days providing showers were not too heavy or prolonged. For each sighting of a Primary Target Species in flight the following was recorded:

- The time that the bird was located and the duration of the observation;
- Sex and age of the bird(s), if possible;
- Behaviour observed such as foraging, commuting or displaying;

- Estimation of flight height;
- Habitats used during flight observation period; and
- Weather conditions at time of sighting.

From the point when an individual was detected it was followed until it ceased flying or was lost from view. The time of initial detection and the flight duration was recorded and the flight path followed was plotted, in the field, onto OSI 1:50 000 mapping. The bird's flight height was estimated at the time of detection and then at evenly spaced intervals thereafter. In order to avoid observer error narrow height bands were not used and flight heights were classified into height bands that can be used in post survey analysis to characterise and describe the flights.

Observations of Target Species took priority over completion of activity summaries. The survey sheet (See **Appendix 2**) is designed to facilitate data entry and allows for the addition of brief notes summarising the flight behaviours. These can subsequently be used to provide qualitative descriptions of the behaviour. Entry of this information was facilitated by use of the codes outlined in **Sections 11.1.1** and **11.1.2**.

Static birds, such as those that are perched were to be recorded on the sheets and the location marked on a map. For clarity, and for ease of post survey analysis, individual flight paths were recorded on separate maps and observation sheets.

#### **11.1.1 Behaviour Codes<sup>23</sup>**

The following codes were used in the survey sheets to indicate the behaviours observed for each sighting:

- (H) Hunting
- (F) Flying
- (S) Soaring
- (C) Circling
- (P) Perched
- (G) On Ground
- (M) Mobbing
- (D) Display
- (FP) Male
- (O) Other

#### **11.1.2 Habitat Codes<sup>23</sup>**

The following codes were used in the survey sheets to indicate the habitats transected by each flight path:

- IG - Improved grazing
- S - Scrub
- B - Bog
- RG - Rough grazing
- G - Grass moorland

---

<sup>23</sup> Derived from Irish Hen Harrier Survey 2015 Survey & recording guidelines for contributors

- 1F - First rotation forest
- 2F - Second rotation forest
- T - Thicket (or pole) stage forest
- CF - Clear fell
- H - Heather moorland
- O - Other (please specify)

## 12 RESULTS: TARGET SPECIES ACTIVITY

Three Primary Target Species and two Secondary Target Species were recorded during the survey period. These are, as follows:

- Primary Target Species:
  - Hen harrier (*C. cyaneus*)
  - Kestrel (*F. tinnunculus*)
  - Sparrowhawk (*A. nisus*)
- Secondary Target Species
  - Cormorant (*P. carbo*)
  - Snipe (*G. gallinago*)

In addition, non-target species namely, mallard (*Anas platyrhynchos*), LBBG (*L. fuscus*) and grey heron (*Ardea cinerea*) were also recorded.

### 12.1 PRIMARY TARGET SPECIES


#### 12.1.1 Hen harrier Observations

One observation of this species was recorded, and this occurred in May. This observation was recorded from VP3 of an adult male. This male hen harrier was observed hunting over bog within the site boundary (see **Figure 1**, above) from VP3. Flight heights were within the 0m-20 m range. This flight path is illustrated in **Figure 6** this is also included in A4 format in **Appendix 5**.

The total time of observations is shown in **Table 3**, below and the characteristics of the flights recorded are summarised in **Table 6**, below. Descriptions of the behaviors recorded are included in **Section 12.1.1.1**, below. A discussion of the survey results is included in **Section 13**, below.

**Table 3: Total Observation Time**

VP	Time in seconds
3	22
Total	22



**Figure 6: Hen harrier flight paths**

#### 12.1.1.1 VP3 (May 21<sup>st</sup>) Flight Path 1

At 08:55 a very clean looking adult male was seen from VP3. It was hunting low over the bog at <5m height to the northwest inside the site boundary. This bird was observed for 22seconds and flew off in a south easterly direction.


### 12.1.2 Kestrel Observations

In total there were three observations (one of which was an incidental sighting) of kestrels inside the site boundary during August and September. All three observations occurred at VP1. The kestrels were observed perched, flying and hunting at various heights ranging from 0m -20 m. The only habitat over flown was bog. These flight paths are illustrated in **Figure 7** this is also included in A4 format in **Appendix 5**.

The total time of observations is shown in **Table 4**, below. The flight characteristics are summarised in **Table 7**, below and the observations are described in **Section 12.1.2.1** to **Section 12.1.2.3**, inclusive, below. A discussion of the survey results is included in **Section 13**, below.

**Table 4 : Total Observation Time**

VP	Time in seconds
1	39
Total	39



**Figure 7: Kestrel flight paths**

#### 12.1.2.1 VP1 (August 8<sup>th</sup>) Flight Path 1

At 10:10 there was an incidental recording of an adult male kestrel as the surveyor drove onto the site. This kestrel was flushed southwest of VP1 over bog from a tree stump as a surveyor drove onto the site and it flew off in a westerly direction inside the site boundary at heights between 0m-20m.

12.1.2.2 VP1 (September 12<sup>th</sup>) Flight Path 2

At 10:30 a kestrel was observed northwest of VP1. This bird was flying at heights between 0m-20m and was observed for 17seconds. It flew north and west back and forth flying quickly over bog habitat inside the site boundary. It was lost to sight flying in a northerly direction.

12.1.2.3 VP1 (September 12<sup>th</sup>) Flight Path 3

At 13:30 a kestrel was observed northwest of VP1. This bird was observed briefly as it was hunting inside the site boundary, it dropped behind a ridge and out of sight flying in a westerly direction. This observation lasted 12seconds and the bird was flying between 0m-20m height over bog.

12.1.3 Sparrowhawk Observations

There was one observation of sparrowhawk during the survey period and this was observed inside the site boundary. This adult was observed at VP2 location and the species was recorded in May only. Flight heights were within the 0m-20 m range. The individual recorded was observed flying over bog habitat. The flight path is illustrated in **Figure 8** this is also included in A4 format in **Appendix 5**.

The total time of the observation is shown in **Table 5**. The flight characteristics are summarised in **Table 8** and the observation is described in **Section 12.2.1.1**. A discussion of the survey results is included in **Section 13**, below.

Table 5: Total Observation Time

VP	Time in seconds
2	15
Total	15




Figure 8: Sparrowhawk flight paths

#### *12.1.3.1 VP2 (May 10<sup>th</sup>) Flight Path 1*

At 07:48 an adult sparrowhawk was observed southeast of VP2. This adult was flying low over the bog inside the site boundary at <3m height and was observed for 15seconds before it flew off in a south easterly direction.

**Table 6: Summary characteristics of hen harrier flights observed**

Flight Path	Figure No.	Date	VP	Time of Observation	Gender/ age	Duration of observation (seconds)	Behaviour	Height Flown (m)	Habitat(s) over flown
<b>Breeding 2020</b>									
Orange	6	21/05/20	3	08:55	Male/ Adult	22	Hunting	0-20m	Bog

**Table 7: Summary characteristics of kestrel flights observed**

Flight Path	Drawing No.	Date	VP	Time of Observation	Gender/age	Duration of observation (in seconds)	Behaviour	Height Flown (m)	Habitat(s) over flown
<b>Breeding 2020</b>									
Blue	7	10/08/20 Incidental	1	10:10	Male/Adult	10	Perched, Flying	0- 20m	Bog
Brown	7	12/09/20	1	10:30	Unknown/ Unknown	17	Flying	0- 20m	Bog
Green	7	12/09/20	1	13:31	Unknown/ Unknown	12	Hunting	0- 20m	Bog

**Table 8: Summary characteristics of sparrowhawk flights observed**

Flight Path	Drawing No.	Date	VP	Time of Observation	Gender/age	Duration of observation (in seconds)	Behaviour	Height Flown (m)	Habitat(s) over flown
<b>Breeding 2020</b>									
Brown	8	15/05/20	2	07:48	Unknown/ Adult	15	Flying	0- 20m	Bog



## 12.2 SECONDARY TARGET SPECIES


### 12.2.1 Cormorant Observations

There was one observation of cormorant in flight during August and this occurred inside the site boundary. This cormorant was observed flying over bog habitat in the east of the site. Flight heights were within the 20m-50m range. The flight path is illustrated in **Figure 9** this is also included in A4 format in **Appendix 5**. The individual flight path is numbered and can be identified by cross reference to the Flight Path number found in Column 1, **Table 11**, below.

The total time of the observation is shown in **Table 9**. The flight characteristics are summarised in **Table 11** and the observation is described in **Section 12.2.1.1**. A discussion of the survey results is included in **Section 13**, below.

**Table 9: Total Observation Time**

VP Number	Time in seconds
VP1	10
Total	10



**Figure 9: Cormorant flight paths**

### 12.2.1.1 VP1 (August 15<sup>th</sup>) Flight Path 1

At 12:26 a cormorant was observed southwest of VP1. This bird was flying northwards though the site, inside the site boundary at 30m height and was observed for 10seconds as it flew over bog.


### 12.2.2 Snipe Observations

There was one recording of snipe during this survey period. In May drumming was heard from two areas to the west of the site, one occurred inside the site boundary. The areas where drumming was recorded from are illustrated in **Figure 10** this is also included in A4 format in **Appendix 5**. Individual flight paths are numbered and can be identified by cross reference to the Flight Path numbers found in Column 1, **Table 12**.

The total time of observations is shown in **Table 10**, below. The flight characteristics are summarised in **Table 12** and the observations are described in **Section 12.2.2.1**. A discussion of the survey results is included in **Section 13**, below.

**Table 10: Total Observation Time**

VP Number	Total (seconds)
VP1	10
Total	10



**Figure 10: Snipe areas of activity**

#### *12.2.2.1 VP3 (May 21st) Areas of activity.*

At 08:55 and 09:07, two snipe were heard drumming from VP3. A bird was heard each time for approximately 5seconds. The first drumming was heard from west of VP3 inside the site boundary as the hen harrier flew through the same area. The second was heard to the northeast of VP3.

**Table 11: Summary characteristics of cormorant flights observed**

Flight Path	Figure No.	Date	VP	Time of Observation	Gender/age	Duration of observation (in seconds)	Behaviour	Height Flown (m)	Habitat(s) over flown
Winter 2018 - 2019									
Purple	9	10/08/20	1	12:26	Unknown	10	Flying	0-20m	Bog

**Table 12: Summary characteristics of snipe flights observed**

Flight Path	Drawing No.	Date	VP	Time of Observation	Gender/ age	Duration of observation (seconds)	Behaviour	Height Flown (m)	Habitat(s) overflown
Breeding - 2020									
Cream	10	21/05/20	3	08:55	Unknown/	5	Drumming	NA	NA
Green				09:07	Unknown	5			

### 12.3 OTHER SPECIES OBSERVED


#### 12.3.1 Mallard Observations

In total there was one observation of mallard made during the breeding survey period. This observation was made from VP1 location inside the site boundary. Mallard appeared in April only, flight heights fall within 0m-20m and the mallard was flying and on the ground over bog habitat. The flight path is illustrated in **Figure 11** this is also included in A4 format in **Appendix 5**.

The total time of observations is shown in **Table 13**. The flight characteristics are summarised in **Table 16** and the observations are described in **Section 12.3.1.1**, below. A discussion of the survey results is included in **Section 13**.

**Table 13: Total Observation Time**

VP	Total in seconds
VP1	7
Total	7



**Figure 11 Mallard flight paths**

##### 12.3.1.1 VP1 (April 14<sup>th</sup>) Flight Path 1.

At 19:25 an adult female mallard was observed east of VP1. This bird was flying over cutover bog and then landed, this observation lasted 7seconds.


### 12.3.2 Lesser black-backed gull Observations

In total there was one observation of LBBG made during the breeding survey period. This observation was made from VP1 location. LBBG appeared in May only, flight heights fall within 0m-30m and the LBBG was flying over bog habitat. The flight path is illustrated in **Figure 12**, this is also included in A4 format in **Appendix 5**.

The total time of observations is shown in **Table 14**. The flight characteristics are summarised in **Table 17** and the observations are described in **Section 12.3.2.1**, inclusive. A discussion of the survey results is included in **Section 13**.

**Table 14: Total Observation Time**

VP	Total in seconds
VP1	33
Total	33



**Figure 12 Lesser black-backed gull flight paths**

#### 12.3.2.1 VP1 (May 5<sup>th</sup>). Flight Path 1.

At 11:25 an adult LBBG was observed west of VP1. This bird flew through the site and southwards over bog at c. 30m height and was observed for a total of 33 seconds.

### 12.3.3 Grey heron, Observations

In total there was one observation of grey heron made during the breeding survey period. This observation was made from VP3 location. Grey heron appeared in May only, flight heights were <10m and the heron was flying over bog habitat and on the ground.

The total time of observations is shown in **Table 15**. The flight characteristics are summarised in **Table 18** and the observations are described in **Section 12.3.3.1**, inclusive. A discussion of the survey results is included in **Section 13**.

**Table 15: Total Observation Time**

VP	Total in seconds
VP3	7
Total	7

#### 12.3.3.1 VP3 (May 14<sup>th</sup>). Flight Path 1.

At 13:58 an adult grey heron was observed VP3. This bird was flying low, <10m height in the bog and then landed within the bog.

**Table 16: Summary characteristics of Mallard flights observed**

Flight Path	Figure No.	Date	VP	Time of Observation	Gender/age	Duration of observation (in seconds)	Behaviour	Height Flown (m)	Habitat(s) overflown
Breeding - 2020									
Blue	11	14/04/20	1	19:25	Female/ Adult	7	Flying, on ground	0-20m	Bog

**Table 17: Summary characteristics of Lesser black-backed gull flights observed**

Flight Path	Figure No.	Date	VP	Time of Observation	Gender/age	Duration of observation (in seconds)	Behaviour	Height Flown (m)	Habitat(s) overflown
Breeding - 2020									
Green	12	11/05/20	1	11:25	Unknown/Adult	33	Flying	20-50m	Bog

**Table 18: Summary characteristics of Grey heron flights observed**

Flight Path	Figure No.	Date	VP	Time of Observation	Gender/age	Duration of observation (in seconds)	Behaviour	Height Flown (m)	Habitat(s) overflown
Breeding - 2020									
NA	/	16/05/20	3	13:58	Unknown/ Adult	7	Flying, on ground	0-20	Bog



## 13 DISCUSSION

Only three of the 13 Primary Target Species<sup>24</sup> and two of the 15 Secondary Target Species were recorded during the survey period and the numbers of observations of individual Target Species, and the activity of bird species generally, was extremely low.

These species are, as follows:

- Primary Target Species:
  - Hen harrier (*C. cyaneus*)
  - Kestrel (*F. tinnunculus*)
  - Sparrowhawk (*A. nisus*)
  
- Secondary Target Species
  - Cormorant (*P. carbo*)
  - Snipe (*G. gallinago*)

In addition, non target species namely, mallard (*A. platyrhynchos*), LBBG (*L. fuscus*) and grey heron (*A. cinerea*) were also recorded.

These species differ from those species recorded during the 2019 breeding survey period. Additional Secondary Target Species recorded in 2020 was cormorant (*P. carbo*). During 2019 breeding survey period buzzard (*Buteo buteo*), little egret (*Egret garzetta*), and unidentified gull's were also recorded

### 13.1 PRIMARY TARGET SPECIES

Hen harrier was recorded on one occasion which comprised a brief observation of an adult male which didn't extend beyond 30 seconds. This very clean male was observed hunting low over the bog at <5m height to the northeast inside the site boundary. During the 2019 breeding survey four observations were made which included two adult males, one adult female and the remaining bird was categorised as a juvenile female. During last year's breeding season (2019) hen harrier were observed from all VP's and for a significantly greater amount of time (412 seconds) compared to the 2020 breeding season (22 seconds) from VP3.

Kestrel was recorded on three occasions (one of which was an incidental sighting), all observations were quite short, the longest of which lasted 17seconds. These were all recorded inside the site boundary during August and September. All three observations occurred at VP1. The kestrels were observed perched, flying and hunting at various heights ranging from 0m-20m. The only habitat over flown was bog. There were significantly more observations of kestrel made in the previous breeding season when a total of 14 observations were made. The majority of the activity again was observed from VP1 from June to September. The kestrels were observed flying at various heights ranging from 0m-50m. These were seen within the bog habitat mainly but also in scrub, improved grassland, 1<sup>st</sup> rotation forestry, grassland moorland and bog track. The activities observed over these habitats include flying and hunting mainly, perched, soaring, being mobbed and circling. During last year's

---

<sup>24</sup> See Section 10

breeding season (2019) kestrel were observed from all VP's and for a significantly greater amount of time (1,137 seconds) compared to the 2020 breeding season (39 seconds) from VP1.

Sparrowhawk was observed flying on one occasion. This agile hunter was only observed for a brief 15seconds from VP2 location and the species was recorded in May only. Flight heights were within the 0m-20m range. The individual recorded was observed flying over bog habitat inside the site boundary. Sparrowhawk was not recorded in the previous breeding season, on the basis of the survey data, it is considered that sparrowhawk were not present to a significant extent during the survey period comprising 2019 and 2020 breeding season.

No observations of whooper swan were made during this 2020 breeding survey. During 2019 whooper swan were not observed during the breeding survey. On the 10<sup>th</sup> and 11<sup>th</sup> of April the site where whooper swan had previously been observed was surveyed. On these dates no whooper swans were observed and cattle were seen grazing in this improved grassland.

### 13.2 SECONDARY TARGET SPECIES

There was one observation of cormorant in flight during August 2020 and this occurred inside the site boundary. This cormorant was observed flying over bog habitat in the east of the site. It is considered, on the basis of the survey data, that cormorant were not present to a significant extent during the survey period comprising 2019 and 2020 breeding season.

Similar to 2019 snipe was recorded on few occasions. There was one recording of snipe during this survey period. In May drumming was heard from two areas to the west of the site one occurring inside the site boundary. There were two sightings of adult snipe during the previous breeding survey period (2019). The flight paths observed were all on the eastern side of the site from VP2 and these snipe were flying over bog and scrub at heights between 0m-20m. These observations were made in April and September. During last year's breeding season (2019) snipe were recorded from VP2 only and for a greater amount of time (30seconds) compared to the 2020 breeding season (10 seconds) from VP1. It is considered, on the basis of the survey data, that snipe were not present to a significant extent during the survey period comprising 2019 and 2020 breeding season.

### 13.3 OTHER SPECIES OBSERVED

Mallard were observed on one occasion during the 2020 breeding survey period. This observation was made from VP1 location inside the site boundary. Mallard appeared in April only, flight heights fall within 0m-20m and the mallard was flying and on the ground over bog habitat. Mallard were observed on more occasions (four) during the 2019 breeding survey period. Mallard appeared in April and May only and flight heights fall within 0m-50m and they were observed flying over bog habitat. During last year's breeding season (2019) mallard were recorded from VP2 and for a greater amount of time (214seconds) compared to the 2020 breeding season (7seconds) from VP1. It is considered, on the basis of the survey data, that mallard were not present to a significant extent during the survey period comprising 2019 and 2020 breeding season.

LBBG were observed on one occasion in the breeding survey period 2020. This observation was made from VP1 location in May. The flight heights fall within 0m-30m and the LBBG was flying over bog habitat inside the site boundary. In breeding survey period 2019 there were two observations of LBBG made. These observations were made from VP1 and VP3 location. LBBG appeared in June only and flight heights fall within 20m-100m. They were observed flying over bog, 1<sup>st</sup> rotation forestry and

grassland moorland. During last year's breeding season (2019) LBBG were recorded from VP1 and VP3 and for a greater amount of time (230seconds) compared to the 2020 breeding season (33 seconds) from VP1. It is considered, on the basis of the survey data, that LBBG were not present to a significant extent during the survey period comprising 2019 and 2020 breeding season.

In total there was one observation of grey heron made during the breeding survey period 2020. This observation was made from VP3 location in May. The flight heights fall within <10m and the heron was flying over bog habitat and on the ground. There were no observations of grey heron in the breeding survey period 2019. It is considered, on the basis of the survey data, that they were not present to a significant extent during the survey period comprising 2019 and 2020 breeding season.

Buzzard, little egret and an unidentified gull were the other species observed during the breeding survey period 2019, however they were not identified during this breeding survey period 2020.

## 14 REFERENCES

- Anon, (2012) *CBS Manual: Guidelines for Countryside Bird Survey participants*. CBS is a joint project of BirdWatch Ireland and the National Parks and Wildlife Service of the Department of Arts, Heritage and the Gaeltacht
- Bibby, C.J., Burgess, N.D. & Hill, D.A. & Mustoe, S. (2000) *Bird Census Techniques* (Second edition). Academic Press, London.
- Brown, A.F. & Shepherd, K. B. (1993) A method for censusing upland breeding waders, *Bird Study*, 40:3, 189-195.
- Colhoun, K. & Cummins, S. (2013), Birds of Conservation Concern in Ireland 2014–2019, *Irish Birds* 9: 523-544
- Directorate General Environment (DGE) (2014) Wind turbines have minor impact on small-bird populations *Science for Environment Policy* 395. A publication of the European Commission DGE News Alert Service, edited by SCU, The University of the West of England, Bristol.
- Drewitt, A.L. & Langston, R. H.W. (2006). Assessing the impacts on wind farms and birds. *Ibis* 148, 29-42. British Ornithologists Union.
- Fossitt, J. A. (2000) *A Guide to Habitats in Ireland*. Kilkenny: The Heritage Council.
- Green, R.E. (1985) Estimating the abundance of breeding Snipe. *Bird Study*, 32, 141-9.
- Mc Guinness, S., Muldoon, C., Tierney, N., Cummins, S., Murray, A., Egan, S., & Crowe, O. (2015). *Bird Sensitivity Mapping for Wind Energy Developments and Associated Infrastructure in the Republic of Ireland Guidance Document*. BirdWatch Ireland, Kilcoole, Wicklow.
- National Parks & Wildlife Service (NPWS) (2012). *River Shannon & River Fergus Estuaries Special Protection Area (Site Code 4077) Conservation Objectives Supporting Document VERSION 1*. (Available at [https://www.npws.ie/sites/default/files/publications/pdf/004077\\_River%20Shannon%20and%20River%20Fergus%20Estuaries%20SPA%20Supporting%20Doc\\_V1.pdf](https://www.npws.ie/sites/default/files/publications/pdf/004077_River%20Shannon%20and%20River%20Fergus%20Estuaries%20SPA%20Supporting%20Doc_V1.pdf))
- Percival, S.M. (2003). *Birds and Wind farms in Ireland: A Review of Potential Issues and Impact Assessment*. Sustainable Energy Ireland.
- Robinson, JA, Colhoun, K, McElwaine, V & Rees, EC. (2004). *Whooper Swan Cygnus cygnus (Iceland population) in Britain and Ireland 1960/61 – 1999/2000*. Waterbird Review Series, The Wildfowl & Wetlands Trust/Joint Nature Conservation Committee, Slimbridge.
- Scottish Natural Heritage (SNH), (2017). *Guidance: Recommended bird survey methods to inform impact assessment of onshore wind farms*. Scottish Natural Heritage
- Scottish Natural Heritage (SNH) (2016). *Assessing Connectivity with Special Protection Areas (SPAs) Guidance*. Scottish Natural Heritage, Inverness.

Thaxter, C.B, Buchanan, G.M., Carr, J., Butchart, S.H.M., Newbold, T., Green, R.E., Tobias, J.A., Foden, W.B., O'Brien, S., and Pearce-Higgins, J.W. (2017). Bird and bat species' global vulnerability to collision mortality at wind farms revealed through a trait-based assessment. *Proc. R. Soc. B* 284

Whitfield, D.P. & Madders, M. (2006). Upland Raptors and the Assessment of Wind farm Impacts. *Ibis* **148**, 43-56. British Ornithologists Union.

Wilson, M., Fernández-Bellon, D., Irwin, S. & O'Halloran, J. (2015). *The interactions between Hen Harriers and wind turbines WINDHARRIER FINAL PROJECT REPORT*. School of Biological, Earth & Environmental Sciences, University College Cork, Cork.

Worden, J., Crowe, O., Einarsson, O., Gardarsson, A., McElwaine, G. & Rees, E.C. 2009. Population size and breeding success of the Icelandic Whooper Swan *Cygnus cygnus*: results of the January 2005 international census. *Wildfowl* **59**: 17-40.

# **Appendix 1**

## NPWS Recommended Methodology for Assessment of Impacts of Proposed Windfarms

## **RECOMMENDED METHODOLOGY FOR ASSESSMENT OF IMPACTS OF PROPOSED WINDFARMS ON BREEDING HEN HARRIER WITHIN THE KNOWN RANGE OF THE SPECIES IN IRELAND**

Of the two main threats to Hen Harriers from windfarms (collision and displacement), the possibility of indirect habitat loss, or displacement, if birds avoid a windfarm area is seen as the most immediate issue. Research to improve assessments of collision risk is ongoing in other countries; the proportion of the breeding population at risk from windfarms that have planning permission at present is small. Other proposed windfarms, within areas of importance for Hen Harrier, should be subject to Environmental Impact Assessment.

### **RELEVANT SPECIES**

Although these recommendations focus on the Hen Harrier as the species of concern, breeding Short-eared Owl may possibly occur at some sites, in which case an assessment of site importance should be made using the same methodology, at times of day appropriate to the species.

### **ASSESSMENT OF SITE IMPORTANCE**

Nine upland areas have been identified by Dúchas as being of national importance for Hen Harrier. All areas of heath/bog habitats within the indicative boundaries of these areas lie within 5km of known nest sites located during the 1998-2000 survey, *i.e.* within the normal foraging range of the male of each pair. Any proposed development, which may have impacts on such habitats, should be subject to a detailed survey, to determine Hen Harrier usage for hunting (foraging).

Important aspects to be considered in an assessment are:

The numbers and breeding success of Hen Harriers that may forage in the area, ideally within 5km of the proposed development site,

The time spent by Hen Harriers in all parts of the site,

The cumulative impact of other windfarms in the area that have been granted planning permission,

Spatial variation in an area's importance to foraging Hen Harriers when:

either occupancy or breeding success are below normal,

fire, overgrazing or turbary temporarily reduce the vegetation cover and hence its value to foraging birds,

nest locations change from year to year.

## **METHODS**

### Survey of breeding occupancy:

An appropriate survey in good weather conditions, with at least two visits in April of breeding pairs within 5km of the site from outer turbines and a second series of visits in July to determine breeding success, would be necessary to interpret results from foraging observations. In years with a run of poor weather during April and May, an intermediate series of observations may be required in June to confirm occupancy by breeding pairs or locate late arriving pairs. Useful information is given in Gilbert *et al.* (1998).

Methodology should be detailed giving dates of survey, map of area searched, and habitat types searched. Results should not include detailed nest locations in public documents (e.g. EIS), but should include minimum distance from the development site.

Data on the number and distance from the site of breeding pairs recorded in the 1998-2000 survey (Norriss *et al.* 2002), and in subsequent years where available, can be provided by Dúchas (contact [dnorriss@duchas.ie](mailto:dnorriss@duchas.ie)).

### **Survey of proposed development site**

#### Description of survey area:

The assessment area should include a strip at least 500m beyond the outermost turbines.

A habitat map of the study area should be produced based on the habitat categories listed in Appendix 1. A more detailed habitat map (for example using the classification in Fossitt (2000)) may be appropriate in some cases.

#### Use of the site:

Madders' (2002) methodology, using timed watches from fixed vantage points (VPs), suits well and can be adapted to local circumstances; those aspects of his procedures relevant to Hen Harriers are summarised below. The objective is to estimate the amount of time birds spend foraging per unit area of the site.

Two 3hour watches per VP per month are recommended for the duration of the breeding season (April – July). A gap of at least one hour between watches is advised.

Restrict observations to 0700-2000 hours and suspend observations during periods of poor



visibility and rain.

Select the minimum number of VPs consistent with complete coverage of the site. VPs should be outside the site where feasible, or located so as to avoid disturbance within the site, but within 1km of the ground being observed. Choose inconspicuous locations, well away from nests, to minimise impact on the birds.

Foraging Harriers usually fly within 10m of the ground and characteristically change direction and height abruptly when searching for prey. Record duration of observation and activity of any Harriers observed according to habitat category.

Map the area of each habitat visible from each VP, either in the field, from photographs or using a GIS. If there is area overlap from different VPs, observation areas should be summed when calculating overall observation rates/unit area. Because fields of view can change substantially with even minor changes in VP location, exact relocation using a GPS and perhaps an inconspicuous marker on the ground is recommended if more than one observer is involved.

The Report should include a summary of the sections of the site used by foraging Hen Harriers, broken down by broad habitat category.

If successful breeding is demonstrated in or close to a site, then VP observations should be continued into August to identify areas used by recently fledged juveniles prior to dispersal.

## **References**

- Fossitt, J. A. (2000) A Guide to Habitats in Ireland. Heritage Council. Kilkenny.
- Gilbert, G., Gibbons, D.W. and Evans, J. (1998) Bird Monitoring Methods – a manual of techniques for key UK species. RSPB, Sandy.
- Madders, M. (2002) Method statement for Vantage Point watches. In: Survey methods to assess windfarm impacts on upland bird communities. Scottish Natural Heritage.
- Norriss, D.W., Marsh, J., McMahon, D. and Oliver, G.A. (2002) A national survey of breeding Hen Harriers *Circus cyaneus* in Ireland 1998-2000. Irish Birds 7: 1-10.

## **APPENDIX 1**

Recommended classification of habitat types for use in assessments of wind farm sites for Hen Harrier

Habitat code

## Description

NF

NF 2

New forestry plantation, trees 20-30 cm high

NF 3

New forestry plantation, trees c 1m in height

NF 4

New forestry plantation, trees >2m in height, patchy thickets

2nd F

2nd F 1 & 2

2nd rotation forestry plantation, trees 20-30 cm high

2nd F 3

New forestry plantation, trees c 1m in height

2nd F 4

New forestry plantation, trees >2m in height, patchy thickets

F

Post thicket plantation

G

Grazing

RG

Rough Grazing & rushy pasture

H/B

Heath / Bog

DE



Deciduous woodland & scrub

GO, Gorse

## **Appendix 2**

### Survey Field Sheets


See over for additional notes

Location: Shroneowen WF	Project No: 19746	VP No:	Observer	Date:		Visibility:			
				Time:	Start: Finish:				
 0 Sky completely clear 1 2 3 4 Sky half cloudy	 5 6 7 8 Sky completely cloudy	Weather		Wind Speed & Direction:		Temp:			
Barn Owl		Goldfinch		Long-eared Owl		Sand Martin		Whooper Swan	
Blackbird		Grasshopper Warbler		Long-tailed Tit		Sedge Warbler		Wigeon	
Blackcap		Grt Black-backed Gull		Magpie		Shelduck		Willow Warbler	
Black-headed Gull		Great Tit		Mallard		Siskin		Woodcock	
Blue Tit		Greenfinch		Meadow Pipit		Skylark		Woodpigeon	
Brambling		Grey Heron		Merlin		Snipe		Wren	
Bullfinch		Grey Partridge		Mistle Thrush		Song Thrush		Yellowhammer	
Buzzard		Grey Wagtail		Moorhen		Sparrowhawk		<b>Additional Species</b>	
Chaffinch		Greylag Goose		Mute Swan		Sptd Flycatcher			
Chiffchaff		Hen Harrier		Peregrine		Starling			
Coal Tit		Herring Gull		Pheasant		Stock Dove			
Collared Dove		Hooded Crow		Pied Wagtail		Stonechat			
Coot		House Martin		Raven		Swallow			
Crossbill		House Sparrow		Red Grouse		Swift			
Cuckoo		Jackdaw		Redpoll		Teal			
Curlew		Jay		Redshank		Tree Sparrow			
Dunlin		Kestrel		Redwing		Treecreeper			
Dunnock		Lapwing		Reed Bunting		Water Rail			
Fieldfare		Lsr-blk-bk Gull		Ringed Plover		Wheatear			
Goldcrest		Linnet		Robin		White-fronted Goose			
Golden Plover		Little Grebe		Rook		Whitethroat			

**TARGET SPECIES FIELD SHEET**

Project No: <b>19746</b>		VP:	Date:	Survey Sheet No:	Surveyor:	Species:			
Location: <b>Shroneowen</b>									
VP Start:			Wind Speed (B 'fort)	Wind Direction:	Visibility:				
VP Finish:									
Weather Conditions:									
Disturbance:									
Time first observed:	<b>Activity Codes:</b> (H) Hunting, (F) Flying, (S) Soaring, (C) Circling, (P) Perched, (G) On Ground, (M) Mobbing, (D) Display. <b>Habitat Codes:</b> (IG) Improved Grassland, (RG) Rough Grassland, (G) Grassland Moorland, (S) Scrub, (B) Bog, (1F) 1 <sup>st</sup> Rotation Forest, (2F) 2 <sup>nd</sup> Rotation Forest, (T) Thicket/Pole Stage Forest, (CF) Clear Fell, (H) Heather Moorland, (L) Lake, (P) Pond, (TSW) Temporary Standing Water, (O) Other (specify):								
Sex:									
Age:									
0m – 20m (Seconds)	Activity/Habitat	20-50m	Activity/Habitat	50-100m	Activity/Habitat	100-150m	Activity/Habitat	>150m	Activity/Habitat

Notes:



**Legend**

- Vantage Point Location
- Parking Location
- Boundary

Map Reproduced From Ordnance Survey Ireland  
 By Permission Of The Government.  
 Licence Number EN 0015720.

0 0.475 0.95 km

**Project Title:**  
 Shronowen Wind Farm

**Client:** Shronowen Wind Farm Limited

**Drawing Title:**  
 Vantage Point Locations

**Drawn:** JK      **Checked:** DoB

**Date:** 19/11/2020      **Scale (A4):** 1:25,000

**Malachy Walsh and Partners**  
 Engineering and Environmental Consultants

## **Appendix 3**

### Vantage Point Survey Summary

## Vantage Point Survey Summary

### Location. Shronowen

#### April 2020 VP 1-3

VP	Date	Observer	Start Time	Finish Time	Length of VP watch (hours)	Weather
1	14/04/2020	CMc	18.35	21.35	3	Cloud cover 2/8, no rain, wind f1, wind direction 4.5, visibility 2.5km
1	28/04/2020	CMc	16.30	19.30	3	Cloud cover 6/8, no rain, wind f2, wind direction 9, visibility 2.5km
2	26/04/2020	CMc	07.00	13.00	6	Cloud cover 8/8, temp 12oC, no rain, wind f2, wind direction 1.5, visibility 2.5km.
3	19/04/2020	CMc	13.00	16.00	3	Cloud cover 8/8, temp 12oC, no rain, wind f2, wind direction 4.5, visibility 2.5km.
3	19/04/2020	CMc	17.30	20.30	3	Cloud cover 8/8, temp 11oC, no rain, wind f2, wind direction 4.5, visibility 2.5km.

#### May 2020 VP 1-3

VP	Date	Observer	Start Time	Finish Time	Length of VP watch (hours)	Weather
1	06/05/2020	CMc	08.30	11.30	3	Cloud cover 6/8, temp 12oc, no rain, wind f3, wind direction 3, visibility 2.5km
1	31/05/2020	CMc	19.50	22.50	3	Cloud cover 2/8, temp 22oC, no rain, wind f3, wind direction 3, visibility 2.5km
2	15/05/2020	CMc	07.00	10.00	3	Cloud cover 2/8, temp 5oC, no rain, wind f1, wind direction 9, visibility 2.5km.
2	30/05/2020	CMc	11.00	14.00	3	Cloud cover 2/8, temp 24oC, no rain, wind f3, wind direction 4.5, visibility 2.5km.
3	21/05/2020	CMc	07.00	10.00	3	Cloud cover 2/8, temp 13oC, no rain, wind f3, wind direction 7.5, visibility 2.5km.
3	21/05/2020	CMc	15.30	18.30	3	Cloud cover 8/8, temp 15oC, no rain, wind f3, wind direction 4.5, visibility 2.5km.





## Vantage Point Survey Summary

### June 2019 VP 1-3

VP	Date	Observer	Start Time	Finish Time	Length of VP watch (hours)	Weather
1	11/05/2020	CMc	07.00	10.00	3	Cloud cover 8/8, temp 13oC, no rain, wind f3, wind direction 1.5, visibility 2.5km.
1	11/05/2020	CMc	10.44	13.44	3	Cloud cover 8/8, temp 14oC, no rain, wind f3, wind direction 1.5, visibility 2.5km.
2	11/05/2020	CMc	15.15	18.15	3	Cloud cover 6/8, temp 18oC, no rain, wind f3, wind direction 1.5, visibility 2.5km.
2	12/05/2020	CMc	07.00	10.00	3	Cloud cover 8/8, temp 12oC, no rain, wind f2, wind direction 12, visibility 2.5km..
3	15/05/2020	CMc	08.30	11.30	3	Cloud cover 2/8, temp 21oC, no rain, wind f1, wind direction 3, visibility 2.5km.
3	16/05/2020	CMc	13.00	16.00	3	Cloud cover 7/8, temp 16oC, no rain, wind f3, wind direction 12, visibility 2.5km.

### July 2020 VP 1-3

VP	Date	Observer	Start Time	Finish Time	Length of VP watch (hours)	Weather
1	04/07/2020	CMc	08.00	11.00	3	Cloud cover 8/8, temp 15oC, drizzle, wind f2, wind direction 9, visibility 1.5km.
1	04/07/2020	CMc	12.00	15.00	3	Cloud cover 8/8, temp 16oC, no rain, wind f1, wind direction 2.5, visibility 2.5km.
2	11/07/2020	CMc	07.30	10.30	3	Cloud cover 3/8, temp 11oC, no rain, wind f1, wind direction 9, visibility 2.5km.
2	12/07/2020	CMc	11.45	14.45	3	Cloud cover 8/8, temp 16oC, no rain, wind f2, wind direction 6, visibility 2.5km..
3	07/07/2020	CMc	12.30	15.30	3	Cloud cover 8/8, drizzle, wind f3, wind direction 9, visibility 1.5km.
3	08/07/2020	CMc	16.25	19.25	3	Cloud cover 8/8, temp 16oC, slight continuous rain, wind f3, wind direction 7.5, visibility 1.5km.

## Vantage Point Survey Summary

### August 2020 VP 1-3

VP	Date	Observer	Start Time	Finish Time	Length of VP watch (hours)	Weather
1	10/08/2020	CMc	10.20	13.20	3	Temp 18oC, no rain, wind f1, wind direction 3, visibility 2.5km.
1	15/08/2020	CMc	15.00	18.00	3	Cloud cover 7/8, temp 23oC, no rain, wind f2, wind direction 3, visibility 2.5km.
2	11/08/2020	CMc	09.30	12.30	3	Cloud cover 7/8, temp 18oC, no rain, wind f3, wind direction 3, visibility 2.5km.
2	24/08/2020	CMc	14.30	17.30	3	Cloud cover 8/8, temp 20oC, no rain, wind f1, wind direction 3, visibility 2.5km..
3	18/08/2020	CMc	09.30	12.30	3	Cloud cover 6/8, temp 18oC, no rain, wind f2, wind direction 7.5, visibility 2.5km.
3	22/08/2020	CMc	12.30	15.30	3	Cloud cover 8/8, temp 15oC, heavy intermittent rain, wind f4, wind direction 9, visibility 1.5km.

### September 2020 VP 1-3

VP	Date	Observer	Start Time	Finish Time	Length of VP watch (hours)	Weather
1	12/09/2020	CMc	09.00	12.00	3	Cloud cover 6/8, temp 14oC, no rain, wind f1, wind direction 7.5, visibility 2.5km.
1	12/09/2020	CMc	13.00	16.00	3	Cloud cover 5/8, temp 16oC, no rain, wind f3, wind direction 7.5, visibility 2.5km.
2	14/09/2020	CMc	14.20	17.20	3	Cloud cover 8/8, temp 19oC, no rain, wind f2, wind direction 3, visibility 2.5km.
2	29/09/2020	CMc	12.30	15.30	3	Cloud cover 7/8, temp 14oC, no rain, wind f2, wind direction 6, visibility 2.5km.
3	15/09/2020	CMc	14.45	17.45	3	Cloud cover 7/8, temp 24oC, slight intermittent rain, wind f1, wind direction 6, visibility 2.5km.
3	25/09/2020	CMc	08.30	11.30	3	Cloud cover 4/8, temp 11oC, moderate intermittent rain, wind f2, wind direction 12, visibility 2.5km.

## Appendix 4

### Target/Secondary Species Observations

Shronowen Breeding 2020 Target Species

Hen harrier														
Date	VP	Sex	Age	Map Flight Path Colour	Habitat	No. Of Birds	Time of Flight/ Obs.	Activity	Flight Height (m)	Time (sec) in Height Category				
										Non-flight	0-50m	50 – 100m	>100m	>200m
21/05/20	3	Male	Adult	Orange	Bog	1	08.55	Hunting	0-20m		22			

Kestrel														
Date	VP	Sex	Age	Map Flight Path Colour	Habitat	No. Of Birds	Time of Flight/ Obs.	Activity	Flight Height (m)	Time (sec) in Height Category				
										Non-flight	0-50m	50 – 100m	>100m	>200m
10/08/20 Incidental	1	Male	Adult	Blue	Bog	1	10.10	Perched, Flying	0- 20m		10			
12/09/20	1	Unknown	Unknown	Brown	Bog	1	10.30	Flying	0- 20m		17			
12/09/20	1	Unknown	Unknown	Green	Bog	1	13.31	Hunting	0- 20m		12			

Sparrowhawk														
Date	VP	Sex	Age	Map Flight Path Colour	Habitat	No. Of Birds	Time of Flight/ Obs.	Activity	Flight Height (m)	Time (sec) in Height Category				
										Non-flight	0-50m	50 – 100m	>100m	>200m
15/05/20	2	Unknown	Adult	Brown	Bog	1	07.48	Flying	0- 20m		15			

Cormorant														
Date	VP	Sex	Age	Map Flight Path Colour	Habitat	No. Of Birds	Time of Flight/ Obs.	Activity	Flight Height (m)	Time (sec) in Height Category				
										Non-flight	0-50m	50 – 100m	>100m	>200m
10/08/20	1	Unknown	Unknown	Purple	Bog	1	12.26	Flying	20-50m		10			

Shronowen Breeding 2020 Target Species

Snipe														
Date	VP	Sex	Age	Map Flight Path Colour	Habitat	No. Of Birds	Time of Flight/ Obs.	Activity	Flight Height (m)	Time (sec) in Height Category				
										Non-flight	0-50m	50 – 100m	>100m	>200m
21/05/20	3	Unknown	Unknown	Cream	NA	2	08.55	Drumming	NA	5				
				Green			09.07			5				


Mallard														
Date	VP	Sex	Age	Map Flight Path Colour	Habitat	No. Of Birds	Time of Flight/ Obs.	Activity	Flight Height (m)	Time (sec) in Height Category				
										Non-flight	0-50m	50 – 100m	>100m	>200m
14/04/20	1	Female	Adult	Blue	Bog	1	19.25	Flying, on ground	0-20m		7			

Lesser Black Backed Gull														
Date	VP	Sex	Age	Map Flight Path Colour	Habitat	No. Of Birds	Time of Flight/ Obs.	Activity	Flight Height (m)	Time (sec) in Height Category				
										Non-flight	0-50m	50 – 100m	>100m	>200m
11/05/20	1	Unknown	Adult	Green	Bog	1	11.25	Flying	20-50m		33			




Grey heron														
Date	VP	Sex	Age	Map Flight Path Colour	Habitat	No. Of Birds	Time of Flight/ Obs.	Activity	Flight Height (m)	Time (sec) in Height Category				
										Non-flight	0-50m	50 – 100m	>100m	>200m
16/05/20	3	Unknown	Adult	NA	Bog	1	13.58	Flying, on ground	0-20m		7			

## **Appendix 5**

### Flight Paths and Activity Areas




**Legend**

-  Vantage Point Location
-  21/05/2020
-  Boundary

Map Reproduced From Ordnance Survey Ireland  
 By Permission Of The Government.  
 Licence Number EN 0015720.

0 0.375 0.75 km



**Project Title:**  
 Shronowen Bird Survey Breeding 2020

**Client:** EMP


**Drawing Title:**  
 Hen Harrier Flightpath

<b>Drawn:</b> JK	<b>Checked:</b> FMcK
------------------	----------------------






<b>Date:</b> 19/11/2020	<b>Scale (A4):</b> 1:20,000
-------------------------	-----------------------------



**Malachy Walsh and Partners**  
 Engineering and Environmental Consultants




**Legend**

-  Vantage Point Location
-  10/08/2020
-  12/09/2020 (1st watch)
-  12/09/2020 (2nd watch)
-  Boundary

Map Reproduced From Ordnance Survey Ireland  
 By Permission Of The Government.  
 Licence Number EN 0015720.

0 0.375 0.75 km



**Project Title:**  
 Shronowen Bird Survey Breeding 2020

**Client:** EMP

**Drawing Title:**  
 Kestrel Flightpaths


<b>Drawn:</b> JK	<b>Checked:</b> FMcK
------------------	----------------------

<b>Date:</b> 19/11/2020	<b>Scale (A4):</b> 1:20,000
-------------------------	-----------------------------






**Malachy Walsh and Partners**  
 Engineering and Environmental Consultants






**Legend**

-  Vantage Point Location
-  15/05/2020
-  Boundary

Map Reproduced From Ordnance Survey Ireland  
 By Permission Of The Government.  
 Licence Number EN 0015720.

0 0.375 0.75 km



**Project Title:**  
 Shronowen Bird Survey Breeding 2020

**Client:** EMP


**Drawing Title:**  
 Sparrowhawk Flightpath

<b>Drawn:</b> JK	<b>Checked:</b> FMcK
------------------	----------------------




<b>Date:</b> 19/11/2020	<b>Scale (A4):</b> 1:20,000
-------------------------	-----------------------------



**Malachy Walsh and Partners**  
 Engineering and Environmental Consultants




**Legend**

-  Vantage Point Location
-  10/08/2020
-  Boundary

Map Reproduced From Ordnance Survey Ireland  
 By Permission Of The Government.  
 Licence Number EN 0015720.

0 0.375 0.75 km



**Project Title:**  
 Shronowen Bird Survey Breeding 2020

**Client:** EMP


**Drawing Title:**  
 Comorant Flightpath

<b>Drawn:</b> JK	<b>Checked:</b> FMcK
------------------	----------------------





<b>Date:</b> 19/11/2020	<b>Scale (A4):</b> 1:20,000
-------------------------	-----------------------------



**Malachy Walsh and Partners**  
 Engineering and Environmental Consultants




**Legend**

-  Vantage Point Location
-  21/05/2020 (08:55)
-  21/05/2020 (09:07)
-  Boundary

Map Reproduced From Ordnance Survey Ireland  
 By Permission Of The Government.  
 Licence Number EN 0015720.

0 0.375 0.75 km



**Project Title:**  
 Shronowen Bird Survey Breeding 2020

**Client:** EMP


**Drawing Title:**  
 Snipe Areas of Activity

<b>Drawn:</b> JK	<b>Checked:</b> FMcK
------------------	----------------------

<b>Date:</b> 19/11/2020	<b>Scale (A4):</b> 1:20,000
-------------------------	-----------------------------



**Malachy Walsh and Partners**  
 Engineering and Environmental Consultants



**Legend**

- ◆ Vantage Point Location
- ▶ 14/04/2020
- Boundary

Map Reproduced From Ordnance Survey Ireland  
 By Permission Of The Government.  
 Licence Number EN 0015720.

0 0.375 0.75 km

**Project Title:**  
 Shronowen Bird Survey Breeding 2020


**Client:** EMP

**Drawing Title:**  
 Mallard Flightpath




Drawn: JK	Checked: FMcK
-----------	---------------

Date: 19/11/2020	Scale (A4): 1:20,000
------------------	----------------------

**Malachy Walsh and Partners**  
 Engineering and Environmental Consultants




**Legend**

-  Vantage Point Location
-  11/05/2020
-  Boundary

Map Reproduced From Ordnance Survey Ireland  
 By Permission Of The Government.  
 Licence Number EN 0015720.

0 0.375 0.75 km



**Project Title:**  
 Shronowen Bird Survey Breeding 2020

**Client:** EMP

**Drawing Title:**  
 Lesser Black-Backed Gull Flightpath

<b>Drawn:</b> JK	<b>Checked:</b> FMcK
------------------	----------------------

<b>Date:</b> 19/11/2020	<b>Scale (A4):</b> 1:20,000
-------------------------	-----------------------------



**Malachy Walsh and Partners**  
 Engineering and Environmental Consultants

## **Appendix 6**

### Non-Target Species of Conservation Concern recorded during VP Surveys

The following summary outlines peak numbers of all non-target species of conservation concern recorded during the breeding 2020 VP surveys.

Swallow (*Saxicola torquatus*) was the most frequently recorded amber listed species. It was recorded in all months of the breeding survey, during April – September.

Meadow pipit (*Anthus pratensis*) was the only non-target red-listed species which was recorded. Meadow pipits were recorded in all months of the breeding survey, during April – September. Amber-listed species which were frequently recorded include robin (*Erithacus rubecula*) recorded on during all months of the surveys. The other amber-listed species recorded were stonechat (*Saxicola torquatus*) recorded in all months accept September, skylark (*Alauda arvensis*) recorded only in June and July, mistle thrush (*Turdus viscivorus*) recorded only in May and wheatear (*Oenanthe*) recorded only in September.

15 green-listed species were recorded during the summer vantage point surveys. The majority of these species are common and widespread and occur in a wide variety of habitat-types, many of which are found within the survey area. Most of these species are present throughout the year while some are summer visitors to Ireland.

The following table outlines monthly peak counts for all non-target species of conservation concern recorded during vantage point surveys at Shronowen breeding 2020.

Common Name	Scientific Name	April	May	June	July	Aug	Sept
Meadow pipit	<i>Anthus pratensis</i>	3	5	4	4	3	6
Mistle thrush	<i>Turdus viscivorus</i>		1				
Robin	<i>Erithacus rubecula</i>	1	2	1	1	1	1
Skylark	<i>Alauda arvensis</i>			2	2		
Stonechat	<i>Saxicola torquatus</i>	2	2	2	2	1	
Swallow	<i>Hirundo rustica</i>	2	5	4	2	3	40
Wheatear	<i>Oenanthe</i>						1

## **Appendix 7**

### List of All Species Recorded



The following table outlines peak counts for all species recorded during the breeding 2020 surveys at Shronowen. A total of 28 species were recorded (Annex I species\* are highlighted in bold).

Common Name	Scientific Name	April	May	June	July	Aug	Sept
Blackbird	<i>Turdus merula</i>	2	4	2	1	2	3
Chiffchaff	<i>Phylloscopus collybita</i>		1				
Cormorant	<i>Phalacrocorax carbo</i>					1	
Cuckoo	<i>Cuculus canorus</i>	1	3	2	1		
Goldfinch	<i>Carduelis carduelis</i>	1				20	
Great tit	<i>Parus major</i>		1				
<b>Hen Harrier*</b>	<i>Circus cyaneus</i>		1				
Hooded crow	<i>Corvus cornix</i>	2	1	2	2	2	3
Jay	<i>Garrulus glandarius</i>						3
Kestrel	<i>Falco tinnunculus</i>					1	2
Lesser Black-backed gull	<i>Larus fuscus</i>			1			
Magpie	<i>Pica pica</i>			1	1	1	2
Meadow pipit	<i>Anthus pratensis</i>	3	5	4	4	3	6
Mistle thrush	<i>Turdus viscivorus</i>		1				
Pheasant	<i>Phasianus colchicus</i>		1	1	1		
Raven	<i>Corvus corax</i>	4	2		3	1	4
Reed bunting	<i>Emberzia schoenichus</i>		2		1	1	
Robin	<i>Erithacus rubecula</i>	1	2	1	1	1	1
Rook	<i>Corvus frugilegus</i>	1					
Sedge warbler	<i>Acrocephalus schoenobaenus</i>	1					
Skylark	<i>Alauda arvensis</i>			2	2		
Snipe	<i>Gallinago gallinago</i>		2	2			
Sparrowhawk	<i>Accipiter nisus</i>		1				
Stonechat	<i>Saxicola torquatus</i>	2	2	2	2	1	
Swallow	<i>Hirundo rustica</i>	2	5	4	2	3	40
Wheatear	<i>Oenanthe</i>						1
Woodpigeon	<i>Columba palumbus</i>		2				
Wren	<i>Troglodytes troglodytes</i>	1	2	1	1	1	1