

Sure Partners Limited

ARKLOW BANK WIND PARK  
PHASE 2  
**ONSHORE GRID  
INFRASTRUCTURE**

**Natura Impact Statement**  
APPENDICES

**Appendix A. Site Synopsis**



# Appendix A

## Site Synopsis



## Buckroneys-Brittass Dunes and Fen SAC Site Synopsis Site Code: 000729

Buckroneys-Brittass Dunes and Fen is a complex of coastal habitats located about 10 km south of Wicklow town. It comprises two main sand dune systems, Brittass Bay and Buckroneys Dunes, connected on the coast by the rocky headland of Mizen Head. The dunes have cut off the outflow of a small river at Mizen Head and a fen, Buckroneys Fen, has developed. A further small sand dune system occurs south of Pennycomequick Bridge.

The site is a Special Area of Conservation (SAC) selected for the following habitats and/or species listed on Annex I / II of the E.U. Habitats Directive (\* = priority; numbers in brackets are Natura 2000 codes):

[1210] Annual Vegetation of Drift Lines [1220] Perennial Vegetation of Stony Banks [1410] Mediterranean Salt Meadows  
[2110] Embryonic Shifting Dunes  
[2120] Marram Dunes (White Dunes) [2130] Fixed Dunes (Grey Dunes)\*  
[2150] Decalcified Dune Heath\*  
[2170] Dunes with Creeping Willow  
[2190] Humid Dune Slacks  
[7230] Alkaline Fens

Along much of the higher parts of the beach at this site, typical annual strandline vegetation occurs. Species such as Sea Rocket (*Cakile maritima*), Prickly Saltwort (*Salsola kali*) and Spear-leaved Orache (*Atriplex prostrata*) are frequent in this zone, with the scarcer Yellow Horned-poppy (*Glaucium flavum*) present in places.

A shingle ridge occurs along the Buckroneys dune system. The amount of exposed shingle is low, but it is likely that shingle underlies much of the sandy areas also. The vegetation on the shingle is similar in composition to that which occurs as part of the drift line and embryonic dune habitats. Sea Sandwort (*Honkenya peploides*) is characteristic, and other species include Sand Couch (*Elymus farctus*), Sand Sedge (*Carex arenaria*), Sea Rocket and Yellow Horned-Poppy.

An area of saline vegetation which conforms to 'Mediterranean salt meadows' occurs in the Buckroneys dune system south of the inlet stream to the fen, and possibly in small areas elsewhere within the site. It is typically dominated by rushes (*Juncus* spp.), and of note is the presence of Sharp Rush (*J. acutus*). Sea Club-rush (*Scirpus maritimus*) also occurs. The area is inundated by the tide only occasionally via the narrow inlet leading to Buckroneys Fen.

Embryonic dune development occurs at the southern part of Brittass and more widely at Buckroneys and Pennycomequick. Typical species are couch grasses (*Elymus* sp.), Sand Sedge and Sea Sandwort. The main dune ridges are dominated by Marram (*Ammophila arenaria*), with herbaceous species such as Sea Spurge (*Euphorbia paralias*), Sea-holly (*Eryngium maritimum*) and Common Restharrow (*Ononis repens*) occurring throughout. The main dune ridges are well developed, reaching heights of 10 m at Brittass. The northern end of the Brittass system has fine examples of parabolic dunes.

Stable fixed dunes are well developed at Brittas and Buckronev. Marram is less frequent in these areas and is replaced by Red Fescue (*Festuca rubra*) as the most common grass species. A rich flora occurs, especially in the more open areas. Common species include Pyramidal Orchid (*Anacamptis pyramidalis*), Common Milkwort (*Polygala vulgaris*), Wild Pansy (*Viola tricolor* subsp. *curtisii*), Carlina Thistle (*Carlina vulgaris*), Biting Stonecrop (*Sedum acre*), Wild Thyme (*Thymus praecox*) and Common Bird's-foot-trefoil (*Lotus corniculatus*). The mature areas of fixed dune also contain Burnet Rose (*Rosa pimpinellifolia*), Bracken (*Pteridium aquilinum*), Wood Sage (*Teucrium scordonia*) and Common Sorrel (*Rumex acetosa*). Mosses such as *Tortula ruralis* subsp. *ruraliformis*, *Rhytidiadelphus triquetris*, and *Homalothecium lutescens* are frequent, along with lichens (*Cladonia* spp., *Peltigera canina*).

This is one of the few Irish east coast sites to possess good examples of wet dune slacks and dunes with Creeping Willow (*Salix repens*). These areas of the dunes have a rich and varied flora, including species such as Creeping Willow, Water Mint (*Mentha aquatica*), Silverweed (*Potentilla anserina*), Meadowsweet (*Filipendula ulmaria*) and Meadow Thistle (*Cirsium dissectum*). The slacks are notably rich in rushes and sedges. Of particular interest is the presence of Sharp Rush (*Juncus acutus*), a scarce species in eastern Ireland and one that is indicative of a saline influence.

The site is also notable for the presence, at the back of the dunes, of areas of decalcified dune heath, a rare habitat type, and one which is listed with priority status in the E.U. Habitats Directive. Heath species present include Heather (*Calluna vulgaris*), Bell Heather (*Erica cinerea*) and Gorse (*Ulex europaeus*).

Buckronev Fen lies west of Mizen Head. It is backed to the west by a dense swamp of Common Reed (*Phragmites australis*). The fen is dominated by Tussock Sedge (*Carex paniculata*), with Water Mint, Purple Loosestrife (*Lythrum salicaria*), Marsh Pennywort (*Hydrocotyle vulgaris*), Greater Bird's-foot-trefoil (*Lotus uliginosus*), Water Horsetail (*Equisetum fluviatile*), small sedges (*Carex* spp.) and other flowering plants. An extensive stand of Blunt-flowered Rush (*Juncus subnodulosus*) is of note. Throughout this area the rare Marsh Fern (*Thelypteris palustris*) is frequent. There are also extensive areas of Rusty Willow (*Salix cinerea* subsp. *oleifolia*) scrub.

This site contains two rare plant species protected under the Flora (Protection) Order, 1999: Wild Asparagus (*Asparagus officinalis* subsp. *prostratus*), in its most northerly Irish station, and Meadow Saxifrage (*Saxifraga granulata*). Other rare species which occur within the site include Green-flowered Helleborine (*Epipactis phyllanthes*), Bird's-foot (*Ornithopus perpusillus*) and Spring Vetch (*Vicia lathyroides*). All of these are Red Data Book species. The rare sedge hybrid *Carex riparia* x *C. vesicaria* (*Carex* x *csomadensis*) is only known from Mizen Head.

The invertebrate fauna of Buckronev fen has been investigated and some notable species have been recorded, including the beetle *Eurynebria complanata* and the following flies: *Machimus cowini*, *Anasimyia lunulata*, *Parhelophilus consimilis* and *Lejogaster splendida*.

Little Tern, a species listed on Annex I of the E.U. Birds Directive, has bred or attempted to breed at Buckronev strand in recent years. In 1992 between 7 and 10 pairs were present and in 1993 up to 8 pairs. Teal are regular in winter (119), as are Curlew (46), Lapwing (515) and Snipe (87). All figures are average peaks for 1994/95 - 1995/96.

The dune systems and beaches are subject to high amenity usage from day-trippers and several areas around the site have been developed as caravan parks, car parks and golf courses. The marginal areas of the fen have been reclaimed, especially at the south end, though these areas still flood in winter and attract waterfowl.

This site is important as an extensive sand dune/fen system with well developed plant communities. Several coastal habitats listed on the E.U. Habitats Directive, including two priority habitats - fixed dune and decalcified dune heath - are present. The area contains two legally protected plants, as well as a number of other rare or scarce plant species. The site provides habitat for some rare species of invertebrate and for the vulnerable Little Tern. A rich flora and fauna has persisted on this site despite extensive amenity use and adjacent farming. However, future land use practices will need to be managed to ensure the continued survival of this unique mosaic of coastal habitats.