

**METROLINK**

Integrated Transport. Integrated Life.

**A15.3**

**Habitat Survey  
Results –  
Species Lists**



**Contents**

1. Habitat Survey Results - Species Lists ..... 1

## 1. Habitat Survey Results - Species Lists

Habitat code <sup>1</sup>	Annex I habitat	Key species
BC1	N/A	Bread wheat <i>Triticum aestivum</i> , oat <i>Avena sativa</i> , potato <i>Solanum tuberosum</i> , root beet <i>Beta vulgaris</i> subsp. <i>vulgaris</i> , six-rowed barley <i>Hordeum vulgare</i> , annual meadow-grass <i>Poa annua</i> , barren brome <i>Anisantha sterilis</i> , false oat-grass <i>Arrhenatherum elatius</i> , wild-oat <i>Avena fatua</i> , American willowherb <i>Epilobium ciliatum</i> , common fumitory <i>Fumaria officinalis</i> , cut-leaved crane's-bill <i>Geranium dissectum</i> , fern-grass <i>Catapodium rigidum</i> , field forget-me-not <i>Myosotis arvensis</i> , germander speedwell <i>Veronica chamaedrys</i> , hedge woundwort <i>Stachys sylvatica</i> , long-headed poppy <i>Papaver dubium</i> , oil-seed rape <i>Brassica napus</i> subsp. <i>oleifera</i> , purple-loosestrife <i>Lythrum salicaria</i> , redshank <i>Persicaria maculosa</i> , shepherd's-purse <i>Capsella bursa-pastoris</i> , yarrow <i>Achillea millefolium</i> and two-rowed barley <i>Hordeum distichon</i> .
BC2	N/A	Wild oat <i>Avena fatua</i> , American willowherb <i>Epilobium ciliatum</i> , tall ramping-fumitory <i>Fumaria bastardii</i> , common fumitory <i>Fumaria officinalis</i> , henbit deadnettle <i>Lamium amplexicaule</i> , red deadnettle <i>Lamium purpureum</i> , common poppy <i>Papaver rhoeas</i> , redshank <i>Persicaria maculosa</i> , curled dock <i>Rumex crispus</i> , groundsel <i>Senecio vulgaris</i> , common chickweed <i>Stellaria media</i> , grey field-speedwell <i>Veronica polita</i> and bush vetch <i>Vicia sepium</i> .
BC4	N/A	Field horsetail <i>Equisetum arvense</i> , fumitory species <i>Fumaria</i> sp., Yorkshire-fog <i>Holcus lanatus</i> , black medick <i>Medicago lupulina</i> , common ragwort <i>Jacobaea vulgaris</i> , groundsel <i>Senecio vulgaris</i> and common nettle <i>Urtica dioica</i> .
BL1	N/A	Rustyback <i>Asplenium ceterach</i> , red valerian <i>Centranthus ruber</i> , ivy-leaved toadflax <i>Cymbalaria muralis</i> , Canadian fleabane <i>Erigeron canadensis</i> and prickly sow-thistle <i>Sonchus asper</i> .
BL2	N/A	Brambles <i>Rubus fruticosus</i> agg., cleavers <i>Galium aparine</i> , false oat-grass <i>Arrhenatherum elatius</i> , ash <i>Fraxinus excelsior</i> and hawthorn <i>Crataegus monogyna</i> .
BL3	N/A	Daisy <i>Bellis perennis</i> , shepherd's-purse <i>Capsella bursa-pastoris</i> , perennial rye-grass <i>Lolium perenne</i> , greater plantain <i>Plantago major</i> , common ragwort <i>Jacobaea vulgaris</i> , prickly sow-thistle <i>Sonchus arvensis</i> , smooth sow-thistle <i>Sonchus oleraceus</i> , common chickweed <i>Stellaria media</i> , dandelion <i>Taraxacum officinale</i> agg. and white clover <i>Trifolium repens</i> .
CW2	1130	Channelled wrack <i>Pelvetia canaliculata</i> .
ED1	N/A	No flora species recorded
ED2	N/A	No flora species recorded
ED3	N/A	Cock's-foot <i>Dactylis glomerata</i> , common couch <i>Elymus repens</i> , creeping bent <i>Agrostis stolonifera</i> , false oat-grass <i>Arrhenatherum elatius</i> , fern-grass <i>Catapodium rigidum</i> , oat species <i>Avena</i> sp., perennial rye-grass <i>Lolium perenne</i> , red fescue <i>Festuca rubra</i> , Yorkshire-fog <i>Holcus lanatus</i> , compact rush <i>Juncus conglomeratus</i> , hard rush <i>J. inflexus</i> , toad rush <i>J. bufonius</i> , water bent <i>Polypogon viridis</i> , glaucous sedge <i>Carex flacca</i> , American willowherb <i>Epilobium ciliatum</i> , Bilbao's fleabane <i>Erigeron floribundus</i> , black medick <i>Medicago lupulina</i> , burnet-saxifrage species <i>Pimpinella</i> sp., butterfly-bush <i>Buddleja davidii</i> , Canadian fleabane <i>Erigeron canadensis</i> , colt's-foot <i>Tussilago farfara</i> , common field-speedwell <i>Veronica persica</i> , common poppy <i>Papaver rhoeas</i> , cut-leaved crane's-bill <i>Geranium dissectum</i> , dandelion <i>Taraxacum officinale</i> agg., dog's mercury <i>Mercurialis perennis</i> , field bindweed <i>Convolvulus arvensis</i> , groundsel <i>Senecio vulgaris</i> , long-headed poppy <i>Papaver dubium</i> , oil-seed rape <i>Brassica napus</i> , petty spurge <i>Euphorbia peplus</i> , pineappleweed <i>Matricaria discoidea</i> , red valerian <i>Centranthus ruber</i> , ribbed melilot <i>Melilotus officinalis</i> , scarlet pimpernel <i>Lysimachia arvensis</i> ,

<sup>1</sup> Fossitt, J.A. (2000) *A guide to habitats in Ireland*. Heritage Council, Kilkenny, Ireland.

Habitat code <sup>1</sup>	Annex I habitat	Key species
		<p>scentless mayweed <i>Tripleurospermum inodorum</i>, smooth sowthistle <i>Sonchus oleraceus</i>, sun spurge <i>Euphorbia helioscopia</i>, thale cress <i>Arabidopsis thaliana</i>, broad-leaved dock <i>Rumex obtusifolius</i>, common mouse-ear <i>Cerastium fontanum</i>, common nettle <i>Urtica dioica</i>, common ragwort <i>Jacobaea vulgaris</i>, creeping buttercup <i>Ranunculus repens</i>, curled dock <i>Rumex crispus</i>, daisy <i>Bellis perennis</i>, greater plantain <i>Plantago major</i>, red clover <i>Trifolium pratense</i>, ribwort plantain <i>Plantago lanceolata</i>, spear thistle <i>Cirsium vulgare</i>, white clover <i>Trifolium repens</i>, cat's-ear <i>Hypochaeris radicata</i>, common bird's-foot-trefoil <i>Lotus corniculatus</i>, fairy flax <i>Linum catharticum</i>, selfheal <i>Prunella vulgaris</i>, barren strawberry <i>Potentilla sterilis</i>, brambles <i>Rubus fruticosus</i> agg., bush vetch <i>Vicia sepium</i>, common ivy <i>Hedera helix</i>, dove's-foot crane's-bill <i>Geranium molle</i>, field horsetail <i>Equisetum arvense</i>, giant hogweed <i>Heracleum mantegazzianum</i>, great willowherb <i>Epilobium hirsutum</i>, hedge bindweed <i>Calystegia sepium</i>, herb-robert <i>Geranium robertianum</i>, hoary willowherb <i>Epilobium parviflorum</i>, lesser burdock <i>Arctium minus</i>, meadow vetchling <i>Lathyrus pratensis</i>, rosebay willowherb <i>Chamaenerion angustifolium</i>, silverweed <i>Potentilla anserina</i>, smooth hawk's-beard <i>Crepis capillaris</i>, wood avens <i>Geum urbanum</i>, yarrow <i>Achillea millefolium</i>, scarlet pimpernel <i>Anagallis arvensis</i>, columbine species <i>Aquilegia</i> sp., turnip <i>Brassica rapa</i>, bittercress species <i>Cardamine</i> sp., carnation sedge <i>Carex panicea</i>, crested dog's-tail <i>Cynosurus cristatus</i>, common stork's-bill <i>Erodium cicutarium</i>, meadow fescue <i>Schedonorus pratensis</i>, cleavers <i>Galium aparine</i>, bristly oxtongue <i>Helminthotheca echioides</i>, six-rowed barley <i>Hordeum vulgare</i>, nipplewort <i>Lapsana communis</i>, annual meadow-grass <i>Poa annua</i>, knotgrass <i>Polygonum ariculare</i> agg., meadow buttercup <i>Ranunculus acris</i>, common sorrel <i>Rumex acetosa</i>, procumbent pearlwort <i>Sagina procumbens</i>, charlock <i>Sinapsis arvensis</i>, goat's beard <i>Tragopogon pratensis</i>, silver bryum <i>Bryum argenteum</i>, crescent-cup liverwort <i>Lunularia cruciata</i> and pink-fruited thread-moss <i>Pohlia melanodon</i>.</p>
ED5	N/A	No flora species recorded
FL8	N/A	No flora species recorded
FS1	N/A	Amphibious bistort <i>Persicaria amphibia</i> , reed canary-grass <i>Phalaris arundinacea</i> , yellow iris <i>Iris pseudacorus</i> , hogweed <i>Heracleum sphondylium</i> and rosebay willowherb <i>Chamaenerion angustifolium</i> .
FS2	6430	Reed canary-grass <i>Phalaris arundinacea</i> , reed sweet-grass <i>Glyceria maxima</i> , amphibious bistort <i>Persicaria amphibia</i> , bittersweet <i>Solanum dulcamara</i> , branched bur-reed <i>Sparganium erectum</i> , butterbur <i>Petasites hybridus</i> , great willowherb <i>Epilobium hirsutum</i> , hedge bindweed <i>Calystegia sepium</i> , meadowsweet <i>Filipendula ulmaria</i> , water figwort <i>Scrophularia auriculata</i> , water mint <i>Mentha aquatica</i> , wild angelica <i>Angelica sylvestris</i> , wild teasel <i>Dipsacus fullonum</i> , yellow iris <i>Iris pseudacorus</i> and alder <i>Alnus glutinosa</i> .
FW2	N/A	Fool's-water-cress <i>Helosciadium nodiflorum</i> , water mint <i>Mentha aquatica</i> , water horsetail <i>Equisetum fluviatile</i> , common duckweed <i>Lemna minor</i> , filamentous algae, blue water-speedwell <i>Veronica anagallis-aquatica</i> , water-cress <i>Nasturtium officinale</i> , greater water-moss <i>Fontinalis antipyretica</i> .
FW3	N/A	Yellow water-lily <i>Nuphar lutea</i> , amphibious bistort <i>Persicaria amphibia</i> , arrowhead <i>Sagittaria sagittifolia</i> , ivy-leaved duckweed <i>Lemna trisulca</i> , water-plantain <i>Alisma plantago-aquatica</i> , fool's-water-cress <i>Helosciadium nodiflorum</i> , water mint <i>Mentha aquatica</i> , Canadian waterweed <i>Elodea canadensis</i> , New Zealand pigmyweed <i>Crassula helmsii</i> , whorled water-milfoil <i>Myriophyllum verticillatum</i> , common stonewort <i>Chara vulgaris</i> and Nuttall's pondweed <i>Elodea nuttallii</i> .
FW4	N/A	Common duckweed <i>Lemna minor</i> , fool's-water-cress <i>Helosciadium nodiflorum</i> , water horsetail <i>Equisetum fluviatile</i> , wild angelica <i>Anthriscus sylvestris</i> , oat species <i>Avenula</i> sp., water-starwort <i>Callitriche</i> agg., cuckooflower <i>Cardamine pratensis</i> , willowherb species <i>Epilobium</i> sp., tall fescue <i>Schedonorus arundinaceus</i> , meadowsweet <i>Filipendula ulmaria</i> , cut-leaved crane's-bill <i>Geranium dissectum</i> , herb-robert <i>Geranium roberitanum</i> , hogweed <i>Heracleum sphondylium</i> , jointed rush <i>Juncus articulatus</i> , soft-rush <i>Juncus effusus</i> , forget-me-not species <i>Myosotis</i> sp., water-cress <i>Nasturtium officinale</i> agg., poppy species <i>Papaver</i> sp., rough meadow-grass <i>Poa trivialis</i> , creeping buttercup <i>Ranunculus repens</i> , curled dock <i>Rumex crispus</i> , marsh ragwort <i>Jacobaea aquatica</i> and bulrush <i>Typha latifolia</i> .
GA1	N/A	Annual meadow-grass <i>Poa annua</i> , cock's-foot <i>Dactylis glomerata</i> , creeping bent <i>Agrostis stolonifera</i> ,

Habitat code <sup>1</sup>	Annex I habitat	Key species
		crested dog's-tail <i>Cynosurus cristatus</i> , meadow foxtail <i>Alopecurus pratensis</i> , perennial rye-grass <i>Lolium perenne</i> , rough meadow-grass <i>Poa trivialis</i> , Yorkshire-fog <i>Holcus lanatus</i> , broad-leaved dock <i>Rumex obtusifolius</i> , common chickweed <i>Stellaria media</i> , common mouse-ear <i>Cerastium fontanum</i> , common nettle <i>Urtica dioica</i> , creeping buttercup <i>Ranunculus repens</i> , creeping thistle <i>Cirsium arvense</i> , daisy <i>Bellis perennis</i> , dandelion <i>Taraxacum officinale</i> agg., germander speedwell <i>Veronica chamaedrys</i> , greater plantain <i>Plantago major</i> , red clover <i>Trifolium pratense</i> , ribwort plantain <i>Plantago lanceolata</i> , silverweed <i>Potentilla anserina</i> , spear thistle <i>Cirsium vulgare</i> , sticky mouse-ear <i>Cerastium glomeratum</i> , white clover <i>Trifolium repens</i> , field horsetail <i>Equisetum arvense</i> and hybrid fescue <i>X Schedolium loliaceum</i> .
GA2	N/A	Annual meadow-grass <i>Poa annua</i> , common bent <i>Agrostis capillaris</i> , creeping bent <i>Agrostis stolonifera</i> , crested dog's-tail <i>Cynosurus cristatus</i> , perennial rye-grass <i>Lolium perenne</i> , Yorkshire-fog <i>Holcus lanatus</i> , black medick <i>Medicago lupulina</i> , broad-leaved dock <i>Rumex obtusifolius</i> , common ragwort <i>Jacobaea vulgaris</i> , creeping buttercup <i>Ranunculus repens</i> , creeping cinquefoil <i>Potentilla reptans</i> , creeping thistle <i>Cirsium arvense</i> , curled dock <i>Rumex crispus</i> , daisy <i>Bellis perennis</i> , dandelion <i>Taraxacum officinale</i> agg, dove's-foot crane's-bill <i>Geranium molle</i> , greater plantain <i>Plantago major</i> , prickly sow thistle <i>Sonchus asper</i> , ribwort plantain <i>Plantago lanceolata</i> , selfheal <i>Prunella vulgaris</i> , silverweed <i>Potentilla anserina</i> , spear thistle <i>Cirsium vulgare</i> , red clover <i>Trifolium pratense</i> , white clover <i>Trifolium repens</i> , meadow foxtail <i>Alopecurus pratensis</i> , false-brome <i>Brachypodium sylvaticum</i> , soft-brome <i>Bromus hordeaceus</i> , common mouse-ear <i>Cerastium fontanum</i> , short-fruited willowherb <i>Epilobium obscurum</i> , meadow fescue <i>Schedonorus pratensis</i> , red fescue <i>Festuca rubra</i> agg., cut-leaved crane's-bill <i>Geranium dissectum</i> , hogweed <i>Heracleum sphondylium</i> , cat's-ear <i>Hypochaeris radicata</i> , medick <i>Medicago sativa</i> , field forget-me-not <i>Myosotis arvensis</i> , rough meadow-grass <i>Poa trivialis</i> , meadow buttercup <i>Ranunculus acris</i> , brambles <i>Rubus fruticosus</i> agg., groundsel <i>Senecio vulgaris</i> , lesser trefoil <i>Trifolium dubium</i> , common nettle <i>Urtica dioica</i> , common vetch <i>Vicia sativa</i> subsp. <i>segetalis</i> , and bush vetch <i>Vicia sepium</i> .
GS1	N/A	Annual meadow grass <i>Poa annua</i> , cock's-foot <i>Dactylis glomerata</i> , creeping bent <i>Agrostis stolonifera</i> , false brome <i>Brachypodium sylvaticum</i> , false oat-grass <i>Arrhenatherum elatius</i> , meadow foxtail <i>Alopecurus pratensis</i> , perennial rye-grass <i>Lolium perenne</i> , quaking-grass <i>Briza media</i> , red fescue <i>Festuca rubra</i> agg., rough meadow-grass <i>Poa trivialis</i> , soft brome <i>Bromus hordeaceus</i> , sweet vernal-grass <i>Anthoxanthum odoratum</i> , timothy <i>Phleum pratense</i> , Yorkshire-fog <i>Holcus lanatus</i> , hard rush <i>Juncus inflexus</i> , soft rush <i>Juncus effusus</i> , toad rush <i>Juncus bufonius</i> , common sedge <i>Carex nigra</i> , glaucous sedge <i>Carex flacca</i> , hairy sedge <i>Carex hirta</i> , remote sedge <i>Carex remota</i> , cat's ear <i>Hypochaeris radicata</i> , common bird's-foot-trefoil <i>Lotus corniculatus</i> , fairy flax <i>Linum catharticum</i> , lady's bedstraw <i>Galium verum</i> , wild carrot <i>Daucus carota</i> subsp. <i>carota</i> , rough hawkbit <i>Leontodon hispidus</i> , selfheal <i>Prunella vulgaris</i> , yellow-rattle <i>Rhinanthus minor</i> , yellow-wort <i>Blackstonia perfoliata</i> , bee orchid <i>Ophrys apifera</i> , pyramidal orchid <i>Anacamptis pyramidalis</i> , black medick <i>Medicago lupulina</i> , brambles <i>Rubus fruticosus</i> agg., bush vetch <i>Vicia sepium</i> , colt's-foot <i>Tussilago farfara</i> , common mouse-ear <i>Cerastium fontanum</i> , common ragwort <i>Jacobaea vulgaris</i> , creeping buttercup <i>Ranunculus repens</i> , creeping cinquefoil <i>Potentilla reptans</i> , creeping thistle <i>Cirsium arvense</i> , curled dock <i>Rumex crispus</i> , daisy <i>Bellis perennis</i> , dandelion <i>Taraxacum officinale</i> agg., groundsel <i>Senecio vulgaris</i> , common knapweed <i>Centaurea nigra</i> , field horsetail <i>Equisetum arvense</i> , greater plantain <i>Plantago major</i> , hedge mustard <i>Sisymbrium officinale</i> , herb-robert <i>Geranium robertianum</i> , kidney vetch <i>Anhyllis vulneraria</i> , lesser trefoil <i>Trifolium dubium</i> , meadow buttercup <i>Ranunculus acris</i> , meadow vetchling <i>Lathyrus pratensis</i> , oxeye daisy <i>Leucanthemum vulgare</i> , pineappleweed <i>Matricaria discoidea</i> , red bartsia <i>Odontites vernus</i> , red clover <i>Trifolium pratense</i> , ribwort plantain <i>Plantago lanceolata</i> , shepherd's-purse <i>Capsella bursa-pastoris</i> , silverweed <i>Potentilla anserina</i> , slender St. John's-wort <i>Hypericum pulchrum</i> , smooth hawk's-beard <i>Crepis capillaris</i> , spear thistle <i>Cirsium vulgare</i> , square-stalked St. John's-wort <i>Hypericum tetrapterum</i> , trailing tormentil <i>Potentilla anglica</i> , tufted vetch <i>Vicia cracca</i> , yarrow <i>Achillea millefolium</i> and white clover <i>Trifolium repens</i> .
GS2	N/A	Cock's-foot <i>Dactylis glomerata</i> , common bent <i>Agrostis capillaris</i> , false oat-grass <i>Arrhenatherum</i>

Habitat code <sup>1</sup>	Annex I habitat	Key species
		<i>elatus</i> , perennial rye-grass <i>Lolium perenne</i> , red fescue <i>Festuca rubra</i> agg., rough meadow-grass <i>Poa trivialis</i> , sweet vernal-grass <i>Anthoxanthum odoratum</i> , Yorkshire-fog <i>Holcus lanatus</i> , pendulous sedge <i>Carex pendula</i> , hard rush <i>Juncus inflexus</i> , black medick <i>Medicago lupulina</i> , brambles <i>Rubus fruticosus</i> agg., broad-leaved dock <i>Rumex obtusifolius</i> , bush vetch <i>Vicia sepium</i> , colt's-foot <i>Tussilago farfara</i> , common chickweed <i>Stellaria media</i> , common knapweed <i>Centaurea nigra</i> , common nettle <i>Urtica dioica</i> , common ragwort <i>Jacobaea vulgaris</i> , creeping buttercup <i>Ranunculus repens</i> , creeping cinquefoil <i>Potentilla reptans</i> , creeping thistle <i>Cirsium arvense</i> , curled dock <i>Rumex crispus</i> , dandelion <i>Taraxacum officinale</i> agg., goat's-beard <i>Tragopogon pratensis</i> , hedge bindweed <i>Calystegia sepium</i> , hedge woundwort <i>Stachys sylvatica</i> , hogweed <i>Heracleum sphondylium</i> , kidney vetch <i>Anthyllis vulneraria</i> , meadow buttercup <i>Ranunculus acris</i> , meadow vetchling <i>Lathyrus pratensis</i> , nipplewort <i>Lapsana communis</i> , oxeye daisy <i>Leucanthemum vulgare</i> , prickly sowthistle <i>Sonchus asper</i> , red clover <i>Trifolium pratense</i> , ribwort plantain <i>Plantago lanceolata</i> , rosebay willowherb <i>Chamerion angustifolium</i> , silverweed <i>Potentilla anserina</i> , small burnet <i>Sanguisorba minor</i> subsp. <i>balearica</i> , spear thistle <i>Cirsium vulgare</i> , tufted vetch <i>Vicia cracca</i> , white clover <i>Trifolium repens</i> , wild angelica <i>Angelica sylvestris</i> , wild teasel <i>Dipsacus fullonum</i> , winter heliotrope <i>Petasites pyrenaicus</i> , creeping bent <i>Agrostis stolonifera</i> , garlic mustard <i>Alliaria petiolata</i> , meadow foxtail <i>Alopecurus pratensis</i> , scarlet pimpernel <i>Anagallis arvensis</i> , mugwort <i>Artemisia vulgaris</i> , soft-brome <i>Bromus hordeaceus</i> , butterfly-bush <i>Buddleja davidii</i> , glaucous sedge <i>Carex flacca</i> , common mouse-ear <i>Cerastium fontanum</i> , great willowherb <i>Epilobium hirsutum</i> , hoary willowherb <i>Epilobium parviflorum</i> , field horsetail <i>Equisetum arvense</i> , meadow fescue <i>Schedonorus pratensis</i> , tall ramping-fumitory <i>Fumaria bastardii</i> , hedgerow crane's-bill <i>Geranium pyrenacium</i> , common bird's-foot-trefoil <i>Lotus corniculatus</i> , mallow species <i>Malva</i> sp., changing forget-me-not <i>Myosotis discolor</i> , long-headed poppy <i>Papaver dubium</i> , timothy <i>Phleum pratense</i> , cowslip <i>Primula veris</i> , selfheal <i>Prunella vulgaris</i> , common sorrel <i>Rumex acetosa</i> , red campion <i>Silene dioica</i> , charlock <i>Sinapsis arvensis</i> , hedge mustard <i>Sisymbrium officinale</i> , Alexanders <i>Smyrniolum olusatrum</i> , lesser trefoil <i>Trifolium dubium</i> , gorse <i>Ulex europaeus</i> , germander speedwell <i>Veronica chamaedrys</i> and common field-speedwell <i>Veronica persica</i> .
GS4	N/A	Creeping bent <i>Agrostis stolonifera</i> , rough meadow-grass <i>Poa trivialis</i> , Yorkshire-fog <i>Holcus lanatus</i> , hard rush <i>Juncus inflexus</i> , soft rush <i>Juncus effusus</i> , false fox-sedge <i>Carex otrubae</i> , hairy sedge <i>Carex hirta</i> , common fleabane <i>Pulicaria dysenterica</i> , field horsetail <i>Equisetum arvense</i> , great willowherb <i>Epilobium hirsutum</i> , meadowsweet <i>Filipendula ulmaria</i> , silverweed <i>Potentilla anserina</i> , bulrush <i>Typha latifolia</i> , common nettle <i>Urtica dioica</i> , common ragwort <i>Jacobaea vulgaris</i> , creeping cinquefoil <i>Potentilla reptans</i> , curled dock <i>Rumex crispus</i> , meadow buttercup <i>Ranunculus acris</i> , goat willow <i>Salix caprea</i> and grey willow <i>S. cinerea</i> .
WD1	N/A	Alder <i>Alnus glutinosa</i> , ash <i>Fraxinus excelsior</i> , beech <i>Fagus sylvatica</i> , blackthorn <i>Prunus spinosa</i> , elder <i>Sambucus nigra</i> , hazel <i>Corylus avellana</i> , field maple <i>Acer campestre</i> , hawthorn <i>Crataegus monogyna</i> , holly <i>Ilex aquifolium</i> , hornbeam <i>Carpinus betulus</i> , Lombardy-poplar <i>Populus nigra</i> 'Italica', sycamore <i>Acer pseudoplatanus</i> , wych elm <i>Ulmus glabra</i> , Alexanders <i>Smyrniolum olusatrum</i> , bluebell <i>Hyacinthoides non-scripta</i> , brambles <i>Rubus fruticosus</i> agg., cherry laurel <i>Prunus laurocerasus</i> , common nettle <i>Urtica dioica</i> , common ivy <i>Hedera helix</i> , cow parsley <i>Anthriscus sylvestris</i> , false-brome <i>Brachypodium sylvaticum</i> , false oat-grass <i>Arrhenatherum elatius</i> , flowering currant <i>Ribes sanguineum</i> , herb-robert, <i>Geranium robertianum</i> , hogweed <i>Heracleum sphondylium</i> , rosebay willowherb <i>Chamerion angustifolium</i> , Leyland cypress <i>Cupressus x Leylandii</i> , yew <i>Taxus baccata</i> , horse-chestnut <i>Aesculus hippocastanum</i> , soft-shield fern <i>Polystichum setiferum</i> , oak species <i>Quercus</i> sp., dog-rose <i>Rosa canina</i> agg., broad-leaved dock <i>Rumex obtusifolius</i> , rusty willow <i>Salix cinerea</i> subsp. <i>oleifolia</i> , snowberry <i>Symphoricarpos albus</i> and guelder-rose <i>Viburnum opulus</i> .
WD2	N/A	Beech <i>Fagus sylvatica</i> , ivy <i>Hedera helix</i> , lodgepole pine <i>Pinus contorta</i> , poplar species <i>Populus</i> sp. and elder <i>Sambucus nigra</i> .
WD3	N/A	Pine <i>Pinus nigra</i> , oak species <i>Quercus</i> sp., rowan <i>Sorbus aucuparia</i> , sycamore <i>Acer pseudoplatanus</i> , perennial rye-grass <i>Lolium perenne</i> , rough meadow-grass <i>Poa trivialis</i> , brambles <i>Rubus fruticosus</i>

Habitat code <sup>1</sup>	Annex I habitat	Key species
		agg., common ivy <i>Hedera helix</i> , cleavers <i>Galium aparine</i> , dandelion <i>Taraxacum officinale</i> agg. herb-robert <i>Geranium robertianum</i> , smooth sowthistle <i>Sonchus oleraceus</i> , spear thistle <i>Cirsium vulgare</i> and wood avens <i>Geum urbanum</i> .
WD5	N/A	Ash <i>Fraxinus excelsior</i> , beech <i>Fagus sylvatica</i> , birch species <i>Betula</i> sp., common lime <i>Tilia x europaea</i> , copper beech <i>Fagus sylvatica purpurea</i> , field maple <i>Acer campestre</i> , holly <i>Ilex aquifolium</i> , holm oak <i>Quercus ilex</i> , hornbeam <i>Carpinus betulus</i> , horse-chestnut <i>Aesculus hippocastanum</i> , Norway maple <i>Acer platanoides</i> , oak species <i>Quercus</i> sp., small-leaved lime <i>Tilia cordata</i> , sycamore <i>Acer pseudoplatanus</i> , wild cherry <i>Prunus avium</i> , annual meadow-grass <i>Poa annua</i> , common bent <i>Agrostis capillaris</i> , perennial rye-grass <i>Lolium perenne</i> , rough meadow-grass <i>Poa trivialis</i> , Yorkshire-fog <i>Holcus lanatus</i> , brambles <i>Rubus fruticosus</i> agg., cleavers <i>Galium aparine</i> , common ivy <i>Hedera helix</i> , creeping buttercup <i>Ranunculus repens</i> , curled dock <i>Rumex crispus</i> , daisy <i>Bellis perennis</i> , dandelion <i>Taraxacum officinale</i> agg., herb-robert <i>Geranium robertianum</i> , hogweed <i>Heracleum sphondylium</i> , selfheal <i>Prunella vulgaris</i> , smooth sowthistle <i>Sonchus oleraceus</i> , spear thistle <i>Cirsium vulgare</i> , white clover <i>Trifolium repens</i> and wood avens <i>Geum urbanum</i> .
WL1	N/A	Ash <i>Fraxinus excelsior</i> , beech <i>Fagus sylvatica</i> , Lombardy-poplar <i>Populus nigra</i> 'Italica', blackthorn <i>Prunus spinosa</i> , cherry laurel <i>Prunus laurocerasus</i> , common lime <i>Tilia x europaea</i> , copper beech <i>Fagus sylvatica purpurea</i> , elder <i>Sambucus nigra</i> , crack-willow <i>Salix euxina</i> , English elm <i>Ulmus procera</i> , field maple <i>Acer campestre</i> , goat willow <i>Salix caprea</i> , grey willow <i>Salix cinerea</i> , hawthorn <i>Crataegus monogyna</i> , hazel <i>Corylus avellana</i> , horse-chestnut <i>Aesculus hippocastanum</i> , oak species <i>Quercus</i> sp., osier <i>Salix viminalis</i> , poplar species <i>Populus</i> sp., silver birch <i>Betula pendula</i> , sweet chestnut <i>Castanea sativa</i> , sycamore <i>Acer pseudoplatanus</i> , wych elm <i>Ulmus glabra</i> , willow species <i>Salix</i> sp., Leyland cypress <i>Cupressus x Leylandii</i> , Monterey cypress <i>Hesperocyparis macrocarpa</i> , yew <i>Taxus baccata</i> , brambles <i>Rubus fruticosus</i> agg., butterfly-bush <i>Buddleja davidii</i> , cotoneaster species <i>Cotoneaster</i> sp., dog rose <i>Rosa canina</i> agg., garden privet <i>Lucidum ovalifolium</i> , gorse <i>Ulex europaeus</i> , honeysuckle <i>Lonicera periclymenum</i> , New Zealand broadleaf <i>Griselinia littoralis</i> , rose species <i>Rosa</i> sp., snowberry <i>Symphoricarpos albus</i> , false oat-grass <i>Arrhenatherum elatius</i> , false brome <i>Brachypodium sylvaticum</i> , rough meadow-grass <i>Poa trivialis</i> , alexanders <i>Smyrniolus olusatrum</i> , American willowherb, broad-leaved dock <i>Rumex obtusifolius</i> , bush vetch <i>Vicia sepium</i> , cleavers <i>Galium aparine</i> , common ivy <i>Hedera helix</i> , common nettle <i>Urtica dioica</i> , common ragwort <i>Jacobaea vulgaris</i> , cow parsley <i>Anthriscus sylvestris</i> , creeping buttercup <i>Ranunculus repens</i> , creeping thistle <i>Cirsium arvense</i> , curled dock <i>Rumex crispus</i> , field horsetail <i>Equisetum arvense</i> , foxglove <i>Digitalis purpurea</i> , great horsetail <i>Equisetum telmateia</i> , great willowherb <i>Epilobium hirsutum</i> , hedge bindweed <i>Calystegia sepium</i> , hedge mustard <i>Sisymbrium officinale</i> , hedge woundwort <i>Stachys sylvatica</i> , hogweed <i>Heracleum sphondylium</i> , lords-and-ladies <i>Arum maculatum</i> , nipplewort <i>Lapsana communis</i> , perennial sowthistle <i>Sonchus arvensis</i> , primrose <i>Primula vulgaris</i> , red clover <i>Trifolium pratense</i> , remote sedge <i>Carex remota</i> , ribwort plantain <i>Plantago lanceolata</i> , rough hawkbit <i>Leontodon hispidus</i> , spear thistle <i>Cirsium vulgare</i> , tufted vetch <i>Vicia cracca</i> , cock's foot <i>Dactylis glomerata</i> , holly <i>Ilex aquifolium</i> , purple toadflax <i>Linaria purpurea</i> and wild plum <i>Prunus domestica</i> .
WL2	N/A	Alder <i>Alnus glutinosa</i> , ash <i>Fraxinus excelsior</i> , aspen <i>Populus tremula</i> , beech <i>Fagus sylvatica</i> , blackthorn <i>Prunus spinosa</i> , cherry laurel <i>Prunus laurocerasus</i> , common oak <i>Quercus robur</i> , common lime <i>Tilia x europaea</i> , hawthorn <i>Crataegus monogyna</i> , horse-chestnut <i>Aesculus hippocastanum</i> , elder <i>Sambucus nigra</i> , field maple <i>Acer campestre</i> , green alkanet <i>Pentaglottis sempervirens</i> , holly <i>Ilex aquifolium</i> , Leyland cypress <i>Cupressus x Leylandii</i> , Monterey cypress <i>Hesperocyparis macrocarpa</i> , wych elm <i>Ulmus glabra</i> , silver birch <i>Betula pendula</i> , sycamore <i>Acer pseudoplatanus</i> , wild cherry <i>Prunus avium</i> , alexanders <i>Smyrniolus olusatrum</i> , brambles <i>Rubus fruticosus</i> agg., butterfly-bush <i>Buddleja davidii</i> , cherry laurel <i>Prunus laurocerasus</i> , cleavers <i>Galium aparine</i> , common ivy <i>Hedera helix</i> , cow parsley <i>Anthriscus sylvestris</i> , common nettle <i>Urtica dioica</i> , greater plantain <i>Plantago major</i> , groundsel <i>Senecio vulgaris</i> , guelder rose <i>Viburnum opulus</i> , smooth sowthistle <i>Sonchus oleraceus</i> , rose species <i>Rosa</i> sp., wood avens <i>Geum urbanum</i> , hazel <i>Corylus avellana</i> , Cypress tree species <i>Cupressus</i> sp., herb-robert <i>Geranium robertianum</i> , hogweed <i>Heracleum sphondylium</i> , honeysuckle

Habitat code <sup>1</sup>	Annex I habitat	Key species
		<i>Lonicera periclymenum</i> , Scots pine <i>Pinus sylvestris</i> , soft shield-fern <i>Polysichum setiferum</i> , poplar species <i>Populus</i> sp., oak species <i>Quercus</i> sp., dog-rose <i>Rosa canina</i> agg., broad-leaved dock <i>Rumex obtusifolius</i> , elder <i>Sambucus nigra</i> , lime species <i>Tilia</i> sp. and guelder-rose species <i>Viburnum</i> sp.
WS1	N/A	Blackthorn <i>Prunus spinosa</i> , brambles <i>Rubus fruticosus</i> agg., gorse <i>Ulex europaeus</i> , dog-rose <i>Rosa canina</i> agg., butterfly-bush <i>Buddleja davidii</i> , common nettle <i>Urtica dioica</i> and hedge bindweed <i>Calystegia sepium</i> .
WS2	N/A	Birch species <i>Betula</i> sp., oak species <i>Quercus</i> sp., sycamore <i>Acer pseudoplatanus</i> , hawthorn <i>Crataegus monogyna</i> , spindle <i>Euonymus europaeus</i> , blackthorn <i>Prunus spinosa</i> , rusty willow <i>Salix cinerea</i> subsp. <i>oleifolia</i> , and guelder-rose <i>Viburnum opulus</i> .
WS3	N/A	New-Zealand broadleaf <i>Griselinia littoralis</i> , holly <i>Ilex aquifolium</i> and lime species <i>Tilia</i> sp. (cf. <i>henryana</i> )